

Synthesis of *p*-coumaroylquinic acids and analysis of their interconversion

Anggy Lusanna Gutiérrez Ortiz,^a Federico Berti,^a Luciano Navarini,^b Angelo Monteiro,^c Marina Resmini^c and Cristina Forzato^{a,*}

^a*Dipartimento di Scienze Chimiche e Farmaceutiche, Università di Trieste, via L. Giorgieri 1, 34127 Trieste, Italy*

^b*illycaffè S.p.A., via Flavia 143, 34100, Trieste*

^c*School of Biological and Chemical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, United Kingdom*

Abstract

The synthesis of four isomers of *p*-coumaroylquinic acids was performed by esterification of *p*-acetylcoumaroylchloride with a suitable protected (-)-quinic acid. All isomers have been characterized by means of NMR spectroscopy and circular dichroism. Acyl migration was observed in the synthesis of 3-*O-p*-coumaroylquinic acid and 4-*O-p*-coumaroylquinic acid. Calculations on the most stable conformations of all isomers have also been performed to explain the acyl migration observed during the synthesis procedure.

Keywords: chlorogenic acids, coffee, acylation, circular dichroism, dft

Introduction

Chlorogenic acids (CGAs) belong to the family of phenolic compounds often found in plants,^{1,2} that are secondary metabolites involved in defense mechanisms against environmental stress.³ Although CGAs are present in many vegetables⁴ and fruits, like potatoes, pears, apples and berries,⁵ green coffee beans are particularly rich in these compounds,^{6,2} and coffee is in fact the main source of CGAs in the human diet.⁵ Moreover, CGAs can be used as indicators of coffee quality,^{5,7} since the final content of CGAs and their corresponding lactones formed after roasting are responsible for the acidity and bitterness of the beverage.^{6,8} In the last years, some health benefits have also been associated with CGAs and several reports have claimed that CGAs contribute to the prevention of cardiovascular diseases and types 2 diabetes.^{5,9,10,11} CGAs are esters formed between *trans*-cinnamic acids (such as caffeic, ferulic and *p*-coumaric acid) and quinic acid;¹² therefore, depending on the type of cinnamic acid and on which hydroxyl group of the cyclohexane ring in quinic acid is esterified, a great variety of CGAs can be formed, not only as monoesters but also as di- and triesters.¹³ The total content of CGAs in coffee depends on the coffee species (*Coffee arabica* 4-8% and *Coffee canephora* 7-14% of the dry matter basis),^{2,14} but also on the degree of roasting, the agriculture practices as well as the soil composition.³ The most abundant CGA is 5-caffeoylquinic acid **2b** (also called chlorogenic acid, Figure 1), but a total number of 76 CGAs have been isolated and identified in the last few years.¹⁰ *p*-Coumaroylquinic acids (*p*CoQA) are less abundant and for this reason are the less studied¹⁰ so there is a lack of information on their contribution to the aroma of coffee. The experimental procedure for the quantification of CGAs in coffee beans is rather complex since it comprises extraction, separation and purification processes and their identification

* Corresponding author: Cristina Forzato, e-mail: cforzato@units.it, phone: +390405583921

and quantification is usually done by HPLC coupled with mass spectrometry.^{3,6,10,14} For this reason, it is important to have analytical standards to unequivocally determine their presence in coffee. Since Panizzi et al.^{15,16} reported the first synthesis of CGAs, several literature reports have described the synthesis of different isomers of CQA (Sefkow et al.)^{17,18} and FQA (Dokli et al.).¹⁹ In 1961 Haslam et al.²⁰ synthesized for the first time 5-*O-p*-coumaroylquinic acid using a condensation reaction and subsequently in 1964²¹ the same authors synthesized the other three isomers using acyl migration as the synthetic method. It is important to highlight that a different numbering system for the substituents on the cyclohexane ring was adopted, resulting in different names.²² In the same way, Chao-Mie Ma *et al.*²³ carried out the synthesis of 5-*O-p*-coumaroylquinic acid by condensation between quinic acid bisacetonide and *p*-acetylcoumaroylchloride, in order to evaluate its potential antifungal activity. Even though it seems that all the methods described in literature involve the esterification of the quinic acid with an acyl chloride, to protect selectively the hydroxyl groups of the cyclohexane ring, there is still a lack of information about the synthetic route and characterisation data of the different isomers of *p*-coumaroylquinic acid.

In this work we report the synthesis and characterization of four commercially unavailable isomers of *p*-coumaroylquinic acid: 1-*O-p*-coumaroylquinic acid **1**, 5-*O-p*-coumaroylquinic acid **2a**, 3-*O-p*-coumaroylquinic acid **3a**, 4-*O-p*-coumaroylquinic acid **4a** (Figure 1) that were carried out with some modifications of the synthesis route proposed by Sefkow et al.^{17,18} and Dokli et al.¹⁹ The work is complemented by a computational study which provided important evidence to explain the variations in outcome and chemical yields of the different reactions, as a result of the relative stability of the target products and their intermediates, together with side products resulting from interconversions.





Figure 1

Results and discussion

All *p*-CoQAs were synthesized by coupling *p*-acetylcoumaroylchloride^{18,24} with the different targeted (-)-quinic acids containing only one free hydroxyl group and all the other hydroxyl groups protected in different ways. Although there is no report in the literature of quinic acid rings esterified at position 1 occurring in nature,²⁵ this compound was identified as one of the targets. It was envisaged that its availability would allow it to be used as a standard to further confirm the absence of these compounds in natural source. The synthesis of 1-*O-p*-coumaroylquinic acid **1** was carried out according to scheme 1. In the first step the protection of the hydroxyl groups in positions 3,4 and 5 was achieved by a modified literature procedure.^{19,26} Lactone **5** was synthesized in 72% yield and used in the next step without further purification. The second step involved the condensation reaction with *p*-acetylcoumaroylchloride, using DMAP with pyridine in dichloromethane at room temperature. The protected ester **6** was obtained in 57% yield after purification by column chromatography. The deprotection of all hydroxyl groups was performed in acidic conditions using HCl (2N)/ THF (4:1) under stirring for 11 days to give 1-*O-p*-coumaroylquinic acid **1** in 84% yield (Scheme 1).



Scheme 1. Synthesis of 1-*O-p*-coumaroylquinic acid **1**


The same lactone **5** was used as the starting building block also for the synthesis of 5-*O-p*-coumaroylquinic acid **2a**, following scheme 2. The ethyl carboxylate **7**¹⁹ was directly obtained by treatment with sodium ethoxide in ethanol resulting in an opening of the lactone ring of compound **5** and protection of the carboxylic group. ¹H-NMR of the crude product revealed the presence of the ethyl carboxylate **7** although in a mixture with lactone **5** in a 13:1 ratio. The crude product was

esterified following the same protocol as with **1** and purified by column chromatography to give compound **8** in 48% yield. Protection of the hydroxyl group was not necessary since Pooter *et al.* demonstrated that under mild conditions no esterification occurs at the axial C-1 hydroxyl group of quinic acid.²⁷ Deprotection was performed in 6 days in the presence of HCl (2N)/ THF to obtain 5-*O-p*-coumaroylquinic acid **2a** in 77% from the protected ester **8**.

¹H-NMR data are very similar to that of an authentic sample of 5-*O-p*-caffeoylquinic acid **2b** except for the aromatic ring protons, showing the same stereochemistry and conformation of the cyclohexane ring. The most stable conformation of compound **2a**, as well as that of **2b**, is the one with the ester and carboxylic groups in equatorial position, hydroxyl group at C-4 equatorial and hydroxyl group at C-3 axial. In compound **2b** this is clearly evidenced by the coupling constants and W_H of the proton signals at C-3, C-4 and C-5 (see table 1). H-5 resonates at 5.34 with a W_H of 23.2, indicating an axial position while H-3 resonates at 4.16 with a W_H of 10.7 indicating an equatorial position. Compound **2a** is showing almost an identical spectra for the protons of the quinic ring thus demonstrating that both have the same conformation.


Table 1 – ¹H-NMR in CD₃OD at 500MHz

Compound	Ar protons	Vinyl protons	H-3	H-4	H-5	H-2 H-6
2a	7.47 (d, J 8.5), 6.81 (d, J 8.5)	7.62 (d, J 16.0), 6.32 (d, J 16.0)	4.17 (m, W_H 10.7)	3.72 (dd, J ₁ 3.6, J ₂ 8.2)	5.34 (dt, J ₁ 8.9, J ₂ 4.3, W_H)	2.16-2.25 (2H, m), 2.01-2.11 (2H, m)
2b ²⁸	7.05 (d, J 2.9), 6.96 (dd, J ₁ 8.8, J ₂ 2.9), 6.78 (d, J 8.8)	7.56 (d, J 14.7), 6.26 (d, J 14.7)	4.16 (m, W_H 10.7)	3.72 (dd, J ₁ 7.0, J ₂ 3.6)	5.34 (dt, J ₁ 8.9, J ₂ 5.3, W_H 23.2)	2.16-2.24 (2H, m), 2.02-2.10 (2H, m)
3a	7.47 (d, J 8.5), 6.81 (d, J 8.5)	7.67 (d, J 15.9), 6.39 (d, J 15.9)	5.39 (m, W_H 13.7)	3.71 (1H, dd, J ₁ 7.6, J ₂ 2.7)	4.10 (1H, m, W_H 17.8)	2.10-2.20 (3H, m), 1.93-2.02 (1H, m)
3b ²⁸	7.04 (d, J 2.4), 6.94 (dd, J ₁ 8.8, J ₂ 2.4), 6.78 (d, J 8.8)	7.58 (d, J 16.7), 6.31 (d, J 16.7)	5.35 (m, W_H 11.9)	3.65 (dd, J ₁ 8.7, J ₂ 4.3)	4.14 (dt, J ₁ 8.7, J ₂ 4.3, W_H 21.7)	2.11-2.22 (3H, m), 1.93-1.99 (1H, m)
4a	7.49 (d, J 8.6), 6.82 (d, J 8.6)	7.73 (d, J 15.9), 6.45 (d, J 15.9)	4.32 (m)	4.81 (dd, J ₁ 10.0, J ₂ 2.8)	4.32 (m)	2.17-2.22 (2H, m), 2.00-2.10 (2H, m)
4b ²⁸	7.07 (d, J 3.1), 6.96 (dd, J ₁ 9.4, J ₂ 3.1), 6.78 (d, J 9.4)	7.64 (d, J 15.6), 6.37 (d, J 15.6)	4.28 (m)	4.80 (dd, J ₁ 9.6, J ₂ 3.8)	4.28 (m)	2.16-2.22 (2H, m), 1.98-2.08 (2H, m)



Scheme 2. Synthesis of 5-*O*-*p*-coumaroylquinic acid **2a**

3-*p*-coumaroylquinic acid **3a** was synthesized following scheme 3. The carboxyl group of (-)-quinic acid was protected by esterification with MeOH, followed by protection of the hydroxyl groups at positions 4 and 5 using 2,2,3,3-tetramethoxybutane to give the protected methyl quinate **10**^{17,19,29} with 15% overall yield from (-)-quinic acid. Coupling between *p*-acetylcoumaroylchloride and **10** under standard esterification conditions gave the corresponding ester **11** in 74% yield, after purification by column chromatography. Deprotection reaction under acidic conditions by HCl (2N)/ THF (3:1) for 6 days afforded a 4:1 mixture of 3-*O*-*p*-coumaroylquinic acid **3a** and 4-*O*-*p*-coumaroylquinic acid **4a** (62% conversion yield) as determined by ¹H NMR. It is necessary to monitor the reaction by ¹H-NMR since for prolonged time the hydrolysis reaction of the ester bond between quinic acid and *p*-coumaroyl moiety occurs. Compound **4a** could be recognized by ¹H-NMR since a double of doublet at lower field (4.81 ppm) appeared, due to the presence of the acyl group at C-4, together with an overlapped signal of two protons at C-3 and C-5 at 4.32 ppm. Also in this case the ¹H-NMR spectrum of **3a** is very similar to the one of 3-*O*-caffeoylquinic acid **3b** (see table 1) with respect to the most stable conformation too. The W_{H} 17.8 of H-5 clearly shows that it is an axial proton while H-3 is an equatorial proton due to the lower W_{H} (13.7).




CSA: (-)-10-camphorsulfonic acid

Scheme 3. Synthesis of 3-*O*-*p*-coumaroylquinic acid **3a**

4-*O*-*p*-Coumaroylquinic acid **4a** was obtained after protection at positions 5 and 3 of the quinic acid ring following scheme 4. 1,5- γ -Quinide was synthesized from (-)-quinic acid through dehydration in the absence of any solvent, as described by Wolinsky *et al.*³⁰ and the crude product was purified by heating under reflux in ethyl acetate as suggested by Raheem *et al.*³¹ Recrystallizations of the brown sticky residue with EtOH or MeOH as suggested by Wolinsky *et al.* and other literature procedures^{30,2} were not successful since compound **4** was obtained in less than 5% yield. Subsequently, protection with *tert*-butyldimethylsilylchloride (TBS) following a literature procedure^{30,32} gave a mixture of monosilylated isomers in positions 3 and 4 of the cyclohexane ring in a 70:30 ratio (3-OTBDMS) : (4-OTBDMS) with the protection at position 3 in major amount, as determined by ¹H-NMR spectroscopy. Although several eluents were tried in order to separate the two isomers by flash chromatography it was not possible to isolate the 3-OTBDMS isomer as a pure compound so the mixture of the two was used in the next step. Esterification with *p*-acetylcoumaroylchloride, using pyridine as the solvent, as suggested by Sefkow *et al.*¹⁷ and Dokli *et al.*,¹⁹ gave nevertheless only **15** as a pure compound while no esterification at position 3 was observed as confirmed by ¹H-NMR analysis of the crude product. Compound **15** was obtained in 41% yield after purification by column chromatography and subsequently deprotected under acidic conditions HCl (2N)/ THF (3:1) to give a mixture of isomers **3a** and **4a** in a 1:1 ratio (43% of conversion yield from the protected ester). Since the starting compound was only isomer **15**, an acyl

migration from the C-4 to the C-3 of the cyclohexane ring occurred as it was observed by ^1H NMR spectroscopy. This kind of rearrangement was already observed by Haslam et al. in 1964²¹ when isomers 3- and 5-*O-p*-coumaroylquinic acid were obtained from 4-*O-p*-coumaroylquinic acid by treatment with sodium hydrogen carbonate. Although in our case deprotection reaction was carried out in acidic conditions it seems that the same acyl migration occurs probably by formation of the intermediate orthoesters.



Scheme 4. Synthesis of 4-*O-p*-coumaroylquinic acid **4a**

In order to explain the interconversions observed along the syntheses of the esters, we have carried out a computational analysis on the end products and on the main intermediates leading to their formation (Figure 2).




Figure 2: computational analysis of the interconversions between products and between synthetic intermediates. The relative B3LYP/6.31G(d,p) energies are given in Kcal/mol.

The geometries of the products and intermediates were optimized first at the HF/6.31G(d) level, and then further refined with a DFT calculation carried out at the B3LYP/6.31G(d,p) level. The end products **2a**, **4a** and **3a** show slight differences in energy, the most stable being the 5-acyloyl derivative **2a**. This explains why its direct synthesis from compounds **7** and **8** is not affected by any isomerization. Esters **4a** and **3a** are only 1.2 and 2.9 Kcal/mol less stable, respectively. The overall conformation of the three compounds is very similar, with the carboxyl group at position 1 always found in an equatorial conformation. As a consequence, ester **3a** is the only product with the cumaroyl group in an axial conformation, as experimentally observed in the NMR spectra. The occurrence of a 20% **4a** in the synthesis of **3a** from the protected intermediate **9** (Scheme 3) can therefore be explained by the thermodynamically favored intramolecular acyl transfer from **3a** to **4a**, starting upon deprotection of **9**.

The synthesis of **4a**, as outlined in scheme 4, involves more complex interconversions. Protection of the starting 1,5-g-quinide may lead to two different silylated compounds and the 3-protected derivatives is the most abundant in the reaction crude, while compound **12** is the only product deriving from the acylation of such mixture. 3-OTBDMS is actually much more stable than its isomer 4-OTBDMS, by 8.8 Kcal/mol. In quinides, at difference with quinic derivatives, position 4 is in fact axial, and for this reason the 4-protected compound is strongly destabilized and the bulky protecting group forces the quinide to a boat-like conformations. Full equilibration to the most stable 3-derivative is likely to occur easily, via a pentacoordinate silicon intermediate, and this may explain the fully selective transformation into compound **12**, which, by the way, is even more stable than its isomer **12'** (scheme 5). In the subsequent step of the synthesis, compound **12** is deprotected and hydrolyzed, and a 1:1 mixture of products **4a** and **3a** is obtained. As **12** would lead directly to

4a, and this compound is more stable than **3a**, the only explanation for the observed result may be found if deprotection occurs before the ring opening reaction of quinides intermediates **Q4a** and **Q3a** (Figure 2). The relative stability of the two deprotected quinides is in fact reversed with respect to the end products, and **Q3a** is more stable by over 10 Kcal/mol. Interconversion thus happens at the quinide level and not at the product level in this synthetic path, and its outcome is the result of a complex competition between equilibria.

Circular Dichroism

The Circular dichroism spectra of all isomers **1-4a** were registered and a comparison with the one obtained for the commercially available caffeoyl analogues **2b-4b** were made to verify the same absolute configuration of the stereocenters of the two series of compounds. Additionally, two different solvents were used for this study, methanol and acetonitrile, to establish whether hydrogen bonding can modify the spectra.

In figure 3 the circular dichroism spectra of all compounds in methanol are reported. The CD spectra of the *p*-coumaroylquinic acids **2a-4a** and that of the corresponding caffeoyl analogues **2b-4b** are very similar, indicating that they must have the same absolute configuration of the chiral centers. Furthermore, the same behavior is qualitatively observed for all compounds in both solvents (methanol and acetonitrile) used as it can be noticed comparing figure 3 with figure 4 indicating that the distributions of conformers are quite the same in both solvents.

Compounds **2a,b-3a,b** present a double Cotton effect, with a positive band in the range 290-340nm and a negative band in the range 200-220nm while compounds **4a** and **4b** have both bands negative. To note that **3a** and **3b** present also a third positive band in the range 220-260nm.




Figure 3 – Circular dichroism spectra of compounds **1-4** in MeOH




Figure 4 - Circular dichroism spectra of compounds **1-4** in MeCN

In conclusion, all four isomers of *p*-coumaroylquinic acids were synthesized and characterized. The quantification of these compounds is very important in the area of coffee analysis and they will be used as standards to evaluate a range of coffee matrices of different origins.

The interconversion occurring between the isomers and during their synthesis have been explained on the basis of the relative stability of the isomers and of the intermediates leading to them.

Experimental section

General Methods

All reagents and solvents were purchased from Sigma-Aldrich and used without further purification. Dichloromethane was dried over CaCl_2 . Caffeoylquinic acids **2b-4b** were purchased from Phytolab. Esterification reactions were performed under argon atmosphere. Thin layer chromatographic (TLC) were performed on Merck silica gel 60 F₂₅₄ silica gel plates. ^1H and ^{13}C NMR spectra were recorded with a Varian 500 spectrometer (residual solvent peaks, $\delta = 7.26$ ppm for CDCl_3 and 3.31 ppm for CD_3OD , were used as the internal standard). Electrospray Ionization (ESI) mass spectrometry measurements were performed with an Esquire 400 (Bruker-Daltonics) spectrometer. HRMS-ESI were obtained with a Waters Xevo Q-ToF spectrometer in negative mode.

Infrared spectra (IR) were recorded with an Avatar 320-IR FTIR (ThermoNicolet). Optical rotations were recorded on a Jasco P2000 polarimeter at the wavelength of sodium D band ($\lambda = 589$) using a quartz cell of 1dm length. Circular dichroism spectra were recorded on a Jasco J-710

spectropolarimeter. Melting points were measured with a Sanyo Gallenkamp apparatus and were uncorrected.

p-Acetylcoumaroylchloride

Acetic anhydride was added (4.66g, 45.69mmol) at 0°C to a suspension of *p*-coumaric acid (5g, 30,46mmol) and DMAP (93mg, 0.76 mmol) in pyridine (10mL). The reaction was stirred for 3h at room temperature and then poured onto crushed ice. After acidification with aq. HCl (pH<2) acetyl *p*-coumaric acid was obtained as a white solid which was filtered and washed with water (93% yield). Oxalyl chloride was added at -5 ° C to a suspension of acetyl *p*-coumaric acid (1g, 4.85mmol) in toluene (17mL) containing two drops of DMF and the reaction was stirred at -5 ° C for 2h and then overnight at room temperature. Solvent was removed under reduced pressure to afford *p*-acetylcoumaroylchloride as a yellow solid in 95% yield. NMR data were in accordance with the literature data.²⁴

3,4-O-Isopropylidene -1,5-quinic lactone **5**

2,2dimethoxypropane (4.87g, 46.83 mmol) was added to a suspension of quinic acid (3g, 15.61 mmol) and *p*-toluenesulfonic acid (216 mg, 1.15 mmol) in acetone (60mL) and the mixture was heated under reflux for 2 h. After cooling neutralization with NaHCO₃ (5%) was performed and the mixture was stirred for 1h at room temperature. The reaction mixture was sequentially extracted with CH₂Cl₂ (three times, 20 mL at time) and washed with water (two times, 20 mL at time). The organic layer was dried with Na₂SO₄ and the solvent was removed under reduced pressure. Lactone **5** was obtained as a white solid in 72% yield and was used in the next step without further purification. NMR data were in accordance with the literature.²⁶

1-Acetyl *p*-coumaroyl-3,4-*O*-Isopropylidene quinide **6**

3,4-*O*-isopropylidene-1,5-quinic lactone **5** (500 mg, 2.33mmol) was suspended in CH₂Cl₂ (20 mL), DMAP (86 mg, 0.7 mmol), pyridine (0.47 mL, 4.66mmol) and *p*-acetylcoumaroylchloride (783 mg, 3.49mmol) were added. The mixture was stirred 24h at room temperature. The reaction mixture was diluted with CH₂Cl₂ and sequentially extracted with 1 M aqueous HCl solution (three times, 10 mL at time), NaHCO₃ (5%) (10 mL) and brine (10mL). The organic layer was dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by column chromatography on silica gel (diethyl ether/CH₂Cl₂ = 1/1) to afford ester **6** (57%) as a colorless solid. ¹H NMR (500 MHz, CDCl₃) δ 7.72 (1H, d, *J* = 16.0 Hz, CH=CH), 7.55 (2H, d, *J* = 8.6 Hz, Ar), 7.14 (2H, d, *J* = 8.6 Hz, Ar), 6.41 (1H, d, *J* = 16.0 Hz, CH=CH), 4.82 (1H, dd, *J* = 6.5, 2.5 Hz, H-4), 4.59 (1H, dt, *J* = 2.19, 6.9, H-5), 4.36 (1H, m, H-3), 3.11 (1H, m, H-6), 2.65 (1H, apparent d, H-6), 2.53 (1H, ddd, *J* = 14.0, 6.5, 2.3 Hz, H-2ax), 2.45 (1H, dd, *J* = 14.5, 3.2 Hz, H-2eq), 2.31 (3H, s, CH₃CO), 1.54 (3H, s, CH₃), 1.35 (3H, s, CH₃); ¹³C NMR (500 MHz, CDCl₃) δ 173.65 (s, COO), 169.21 (s, COO), 164.99 (s, COO), 152.59 (s, Ar), 145.71 (d, CH=CH), 131.86 (s, Ar), 129.59 (d, Ar), 122.37 (d, Ar), 117.09 (d, CH=CH), 110.14 (s, C(CH₃)₂), 76.39 (s, C-1), 75.57 (d, C-5), 72.64 (d, C-4), 71.33 (d, C-3), 35.82 (t, C-2), 30.87 (t, C-6), 27.15 C₁₈ (q, C(CH₃)₂), 24.50 (q, C(CH₃)₂), 21.29 (q, CH₃CO).

1-O-p-coumaroylquinic acid **1**

Ester **6** (500mg, 1.24mmol) was dissolved in a mixture of THF (10 mL) and aq. 2M HCl (40mL) and the yellowish solution formed was stirred for 11 days at room temperature. The solution was

saturated with solid NaCl and then extracted with EtOAc (3*20 mL). The organic layer was dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. 1-*p*-coumaroylquinic acid **1** was obtained as a colorless solid in 84% from the corresponding protected ester **6**. M.p. 130-135°C; ¹H NMR (500 MHz, CD₃OD) δ 7.61 (1H, d, *J* = 15.9 Hz, CH=CH), 7.45 (2H, d, *J* = 8.6 Hz, Ar), 6.81 (2H, d, *J* = 8.6 Hz, Ar), 6.35 (1H, d, *J* = 15.9 Hz, CH=CH), 4.15 (1H, q, *J* = 4.3, Hz, H-5), 4.06 (1H, dt, *J* = 9.1, 3.6 Hz, H-3), 3.48 (1H, dd, *J* = 8.3, 3.3 Hz, H-4), 2.57 (1H, m, H-6), 2.44 (1H, m, H-2), 2.21 (dd, *J* = 14.9, 3.5 Hz, H-6), 1.91 (1H, dd, *J* = 13.8, 8.5 Hz, H-2); ¹³C NMR (500 MHz, CD₃OD) δ 174.91 (s, COO), 167.55 (s, COO), 160.79 (s, Ar), 146.40 (d, CH=CH), 130.67 (d, Ar), 126.73 (s, Ar), 116.30 (d, Ar), 114.97 (d, CH=CH), 80.95 (s, C-1), 75.77 (d, C-4), 69.13 (d, C-5), 67.40 (d, C-3), 39.40 (t, C-2), 35.38 (t, C-6); IR (nujol): $\tilde{\nu}$ = 3582.61, 3358.97, 2950, 1693.99, 1631.07, 1170.67, 1113.35, 831.61 cm⁻¹. MS (ESI⁺): *m/z* [M+Na] = 361.0; HRMS (ESI⁺): [M-H] = 337.092 (calculated: 337.092345); [α]_D²⁰ = +5.1 (*c* 1.10, MeOH) (lit. ^{26,33}); [α]_D²² = -5.0 (*c* 2, MeOH)); UV (MeOH): ε₃₁₄ = 84200.

Ethyl-3,4-*O*-Isopropylidene quinate **7**

A suspension of crude lactone **5** (1 g, 4.67mmol) in absolute EtOH (30 mL) was treated with NaOEt (12.71mg, 0.19 mmol) dissolved in EtOH (160 μL). The brownish solution was stirred at room temperature for 2 h and then stored at -20°C for 24h. After quenching the unreacted NaOEt by addition of acetic acid (13 μL) the solvent was removed under reduced pressure at 30 °C. The residue showed to be a mixture of lactone **5** and ester **7** in a ratio 13:1 determined by ¹H NMR analysis. This crude mixture was used without further purification in the next step.^{26,33}

Ethyl-5-*O*-acetyl-*p*-coumaroyl-3,4-*O*-Isopropylidene quinate **8**

To a solution of ethyl-3,4-*O*-Isopropylidene quinate **7** (500 mg, 1.92mmol), DMAP (35mg, 0.15mmol) and pyridine (6 mL) in CH₂Cl₂ (25mL), *p*-acetylcoumaroylchloride (645.18 mg, 2.88mmol) was added. The mixture was stirred 24h at room temperature and acidified with aq. HCl 1M (pH 2-3) and then extracted with CH₂Cl₂ (three times, 50 mL at time). The organic layer was dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The brownish residue was purified by column chromatography on silica gel (diethyl ether/CH₂Cl₂ = 1/1) to afford ester **8** in 48% yield as a colorless solid. ¹H NMR (500 MHz, CDCl₃) δ 7.69 (1H, d, *J* = 15.9 Hz, CH=CH), 7.53 (2H, d, *J* = 8.6 Hz, Ar), 7.13 (2H, d, *J* = 8.6 Hz, Ar), 6.40 (1H, d, *J* = 15.9 Hz, CH=CH), 5.49 (1H, dt, *J* = 11.7, 4.5 Hz, H-5), 4.55 (1H, dt, *J*₁ = 3.7, *J*₂ = 5.6, H-3), 4.28 – 4.20 (3H, m, OCH₂ + H-4), 2.31 (3H, CH₃CO), 2.32 – 2.28 (2H, m, H-2), 2.25 (1H, dd, *J* = 13.2, 4.4 Hz, H-6_{eq}), 1.94 (1H, dd, *J* = 13.3, 11.3 Hz, H-6_{ax}), 1.60 (s, C(CH₃)₂), 1.38 (s, C(CH₃)₂), 1.30 (3H, t, *J* = 7.2 Hz, CH₃CH₂); ¹³C NMR (500 MHz, CDCl₃) δ 174.48 (s, COO), 169.27 (s, COO), 166.03 (s, COO), 152.27 (s, Ar), 144.19 (d, CH=CH), 132.24 (s, Ar), 129.37 (d, Ar), 122.30 (d, Ar), 118.29 (d, CH=CH), 109.76 (s, C(CH₃)₂), 77.05 (d, C-3), 75.65 (s, C-1), 73.81 (d, C-4), 71.11 (d, C-5), 62.36 (t, CH₂CH₃), 37.13 (t, C-6), 34.56 (t, C-2), 28.17 (q, C(CH₃)₂), 26.01 (q, C(CH₃)₂), 21.30 (q, CH₃CO), 14.28 (q, CH₃CH₂).

5-*O*-*p*-coumaroylquinic acid **2a**

Ethyl 1-acetyl *p*-coumaroyl-3,4-*O*-isopropylidene quinate **12** (290mg, 0.65mmol) was dissolved in a mixture of THF (10 mL) and aq. 2M HCl (40mL) and the solution was stirred for 6 days at room temperature. After saturation with solid NaCl the mixture was extracted with EtOAc (3*30 mL) and the organic phase was dried over anhydrous Na₂SO₄. Evaporation of the solvent gave 5-*O*-*p*-coumaroylquinic acid as a colorless solid in 77% yield from the corresponding protected ester **12**.

M.p. 215-218°C (lit. ^{Error! Bookmark not defined.} 247-248°C); IR (nujol): $\tilde{\nu}$ = 3582.67, 3302.38, 2917.48, 1687.13, 1633.37, 1170.30, 1080.85, 825.27 cm⁻¹; ¹H NMR is in accordance with literature data.²³ ¹³C NMR (126 MHz, CD₃OD): δ 177.02 (s, C-7), 168.61 C₈ (s), 161.28 C₁₄ (s), 146.68 C₁₀ (d), 131.18 C_{12,12'} (d), 127.23 C₁₁ (s), 116.80 C_{13,13'} (d), 115.33 C₉(d), 76.15 C₁ (s), 73.41 C₄ (d), 72.00 C₅ (d), 71.15 C₃ (d), 38.77 C₂ (t), 38.22 C₆ (t).; MS (ESI⁺): *m/z* [M+Na]: 361.4; $[\alpha]_D^{20}$ = -39.5 (c 0.79, MeOH) [lit. ^{Error! Bookmark not defined.} $[\alpha]_D^{20}$ = -53.6 (c 1.04, MeOH)]. UV (MeOH): ϵ_{315} = 70000.

4,5-*O*-(2',3'-Dimethoxybutane-2',3'-diyl)-1,3-dihydroxycyclohexanecarboxylic acid methyl ester 10

To a suspension of quinic acid (1g, 5.20mmol) in MeOH (30mL), (-)-10-camphorsulfonic acid (15 mg, 0.065mmol) was added and the mixture was refluxed for 15 h under Ar atmosphere. Methyl quinate **9** so obtained was added with 2,2,3,3-tetramethoxybutane (1.01 g, 5,7mmol), trimethylorthoformate (2.6mL, 0.024mmol) and (-)-10-camphorsulfonic acid (12 mg, 0.052mmol) and the mixture was refluxed again. After 15 h the mixture was cooled and NaHCO₃ (0,1 g) was added. Solution was concentrated under reduced pressure and the orange suspension was partitioned between EtOAc (30mL) and saturated aqueous NaHCO₃(30mL). The aqueous layer was extracted with EtOAc (30mL) and the organic layer was dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. Recrystallization of the brownish residue from EtOAc and hexane (1:5, v/v) afforded **10** in 27% yield as an orange oil. NMR data were in accordance with the literature^{19,29}

3-Acetyl-*p*-coumaroyl-4,5-*O*-(2',3'-Dimethoxybutane-2',3'-diyl)-1-hydroxycyclohexanecarboxylic acid methyl ester 11

4,5-*O*-(2',3'-Dimethoxybutane-2',3'-diyl)-1,3-dihydroxycyclohexanecarboxylic acid methyl ester **10** (122 mg, 0.38mmol) was suspended in CH₂Cl₂ (20 mL) and DMAP (4,17 mg, 0.034 mmol), pyridine (320 μ L, 4.03mmol) and *p*-acetylcoumaroylchloride (128mg, 0.57mmol) were added. The mixture was stirred 24h at room temperature and then acidified with aq. HCl 1M (pH 2-3). After extraction with CH₂Cl₂ (three times, 30 mL at time) the organic layer was dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The brownish residue was purified by column chromatography on silica gel (diethyl ether/CH₂Cl₂ = 1/1) to afford ester **11** in 74% yield as a colorless solid. ¹H NMR (500 MHz, CDCl₃) δ 7.71 (1H, d, *J* = 15.8 Hz, CH=CH), 7.57 (2H, d, *J* = 8.6 Hz, Ar), 7.14 (2H, d, *J* = 8.6 Hz, Ar), 6.47 (1H, d, *J* = 15.9 Hz, CH=CH), 5.38 (1H, q, *J* = 9.1 Hz, H-3), 4.45 (dt, *J* = 10.2, 5.6 Hz, H-5), 3.79 (3H, s, COOCH₃), 3.71 (1H, dd, *J* = 9.9, 3.2 Hz, H-4), 3.31 (3H, s, OCH₃), 3.27 (3H, s, OCH₃), 2.29 (3H, s, CH₃CO), 2.28 (2H, m, H-2 + H-6), 2.14 (1H, dd, *J* = 15.7, 3.2 Hz, H-2), 2.04 (1H, m, H-6); ¹³C NMR (500 MHz, CDCl₃) δ 175.62 (s, COO), 169.26 (s, COO), 166.61 (s, COO), 152.19 (s, Ar), 144.19 (d, CH=CH), 132.45 (s, Ar), 129.47 (d, Ar), 122.21 (d, Ar), 118.79 (d, CH=CH), 100.29 (s, C(OCH₃)), 99.73 (s, C(OCH₃)), 74.78 (s, C-1), 71.38 (d, C-4), 70.02 (d, C-3), 62.94 (d, C-5), 53.39 (q, CH₃COO), 48.3 (q, OCH₃), 48.14 (q, OCH₃), 38.91 (t, C-6), 36.81 (t, C-2), 21.3 (q, CH₃CO), 18.03 (q, CH₃C(OCH₃)), 17.81 (q, CH₃C(OCH₃)).

3-*O*-*p*-coumaroylquinic acid 3a

3-Acetyl-*p*-coumaroyl-4,5-*O*-(2',3'-dimethoxybutane-2',3'-diyl)-1-hydroxycyclohexanecarboxylic acid methyl ester **11** (35mg, 0,069mmol) was dissolved in a mixture of THF (0.5 mL) and aq. 2M HCl (1.5mL) and the solution was stirred for 6 days at room temperature. After saturation with solid NaCl the mixture was extracted with EtOAc (3*20 mL) and the organic phase dried over anhydrous Na₂SO₄. Evaporation of the solvent gave a yellowish solid 62% yield which was defined to be a mixture of 3-*p*-coumaroylquinic acid **3a** and 4-*p*-coumaroylquinic acid **4a** in a ratio 8:2 as determined from ¹H-NMR. The crude was purified by semi-preparative RP-HPLC on a

Phenomenex Gemini C18 5 μm 10 x 250 mm column, using a gradient of H₂O+0,1% formic (A) acid and MeOH+0,1% (B), (20 min A 80% B 20%, from 20 to 90 min increase of B until A 40% B 60%, from 90 to 110 min A 5% B 95%, from 110 to 125 min A 95% B 5%) at a flow rate of 2 mL/min. The elution was monitored with an UV/vis detector λ 325nm and the fractions corresponding to each peak were collected and kept at -80°C and then freeze dried and analyzed by ¹H NMR. 3-*O-p*-coumaroylquinic acid **3a** (3mg) was obtained as a white solid. M.p. 192-194°C [lit. ^{Error! Bookmark not defined.} 194°C]; IR (nujol): $\tilde{\nu}$ =3582.64, 3381.37, 2921.16, 1694.22,1631.26, 1171.87, 1019.74, 831.37 cm^{-1} ; ¹H NMR (500 MHz, CD₃OD) δ 7.67 (1H, d, J = 15.9 Hz, H-10), 7.47 (2H, d, J = 8.5 Hz, H-12), 6.81 (2H, d, J = 8.3 Hz, H-13), 6.39 (1H, d, J = 15.9 Hz, H₉), 5.39 (1H, m, W_{H} 13.7, H-3), 4.10 (1H, m, W_{H} 17.8, H-5), 3.71 (1H, dd, J = 7.6, 2.7 Hz, H-4), 2.20 – 1.93 (4H, m, H-2+H-6); ¹³C NMR (500 MHz, CD₃OD) δ 177.59 (s, COO), 168.91(s, COO), 161.15 (s, Ar), 146.43 (d, CH=CH), 131.09 (d, Ar), 127.39 (s, Ar), 116.79 (d, Ar), 115.85 (d, CH=CH), 76.42 (s, C-1), 74.22 (d, C-4), 72.61 (d, C-3), 68.93 (d, C-5), 36.93 (t, C-2), 36.22 (t, C-6). MS (ESI⁺): m/z [M+Na]: 361.0 [α]_D²⁰ = 2.2 (c 0.12 MeOH) [lit. [α]_D²⁰ = -5.6 (c 0.6, MeOH); UV (MeOH): ϵ_{314} =73000.

1,5- γ -Quinide

Quinic acid (3g, 15.61mmol) was heated in an open flask at 220°C for 90 min. The brown sticky residue was refluxed with EtOAc (60 mL) for 4h and then the solution was cooled to room temperature. The solvent was removed under pressure to give 1,5- γ -quinide as a colorless solid in 85% yield. NMR data were in accordance with the literature.^{28,29}

3-tert-Butyldimethylsiloxy-1,4-dihydroxy-cyclohexane-1,5-carbolactone **13** and **14**

TBSi-Cl (1.31 g, 8.68 mmol) was added to a stirred solution of 1,5- γ -quinide (1.31 g, 7.55 mmol) and imidazole (1,9 g, 28 mmol) in anhydrous DMF (14 mL) at 0 °C. The mixture was stirred at 0 °C for 30 min followed by 1 h at room temperature and then poured into water (50 mL) and extracted with EtOAc (50 ml) and diethyl ether (40 mL). The organic layer was washed with water (3 \times 100 mL), dried over Na₂SO₄ and concentrated under reduced pressure to give a white solid in 57% yield containing esters **13** and **14** in a ratio 7:3. The crude was used in the next step without further purification. NMR data were in accordance with the literature.^{31,32}

4-Acetyl-*p*-coumaroyl-3-tert-butyldimethylsiloxy-1-hydroxycyclohexane-1,5-carbolactone **15**

To a solution of 3-tert-Butyldimethylsiloxy-1,4-dihydroxy-cyclohexane-1,5-carbolactone, as a mixture of **13** and **14**, (500 mg, 1.74mmol) and DMAP (32 mg, 0.26 mmol) in pyridine (15 mL), *p*-acetylcoumarylchloride (700mg, 3.12mmol) was added. The mixture was stirred 24h at room temperature and then poured onto crush ice. Successively CH₂Cl₂ (20mL) was added. The mixture was acidified with aq. HCl 1M (pH 2-3) and then extracted with CH₂Cl₂ (three times, 30 mL at time). The organic layer was dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The brownish residue was purified by column chromatography on silica gel (diethyl ether/CH₂Cl₂ = 1/1) to afford the only ester **15**, in 41% yield, as a colorless solid. ¹H NMR (500 MHz, CDCl₃) δ 7.72 (1H, d, J = 15.9 Hz, CH=CH), 7.57 (2H, d, J = 8.6 Hz, Ar), 7.15 (2H, d, J = 8.6 Hz, Ar), 6.46 (1H, d, J = 15.9 Hz, CH=CH), 5.43 (1H, t, J = 4.8 Hz, H-4), 4.88 (1H, dt, J = 12.6, 5.3 Hz, H-5), 4.03 (1H, dt, J = 10.7, 4.6 Hz, H-3), 2.55 (1H, d, J = 11.8 Hz, H-6), 2.43 (1H, dd, J = 11.8, 5.8 Hz, H-6), 2.32 (3H, s, CH₃CO), 2.11 (2H, apparent d, H-2), 0.81 (9H, s, C(CH₃)₃), 0.06 (3H, s, CH₃Si), 0.03 (3H, s, CH₃Si); ¹³C NMR (500 MHz, CDCl₃) δ 177.48 (s, COO), 169.27 (s, COO), 165.61 (s, COO), 152.50 (s, Ar), 145.15 (d, CH=CH), 131.98 (s, Ar), 129.53 (d, Ar),

122.38 (d, Ar), 117.40 (d, CH=CH), 74.42 (d, C-5), 72.15 (s, C-1), 66.68 (d, C-4), 66.06 (d, C-3), 41.14 (t, C-2), 37.64 (t, C-6), 25.71 (q, C(CH₃)₃), 21.28 (q, CH₃CO), 18.05 (s, C(CH₃)₃), -4.92 (q, (CH₃)₂Si).

4-*O-p*-coumaroylquinic acid **4a**

4-*O*-acetyl-*p*-coumaroyl-3-tert-butyl-dimethylsiloxy-1-hydroxycyclohexane-1,5-carbolactone **15** (332mg, 0.7mmol) was dissolved in a mixture of THF (5 mL) and aq. 2M HCl (15mL) and the solution was stirred for 6 days at room temperature. After saturation with solid NaCl the mixture was extracted with EtOAc (3*50 mL) and the organic phase was dried over Na₂SO₄. Evaporation of the solvent gave a yellowish solid in 43% yield which was a mixture of 3-*p*-coumaroylquinic acid **3a** and 4-*p*-coumaroylquinic acid **4a** as determined by ¹H-NMR. The crude was purified by semi-preparative RP-HPLC on a Phenomenex Gemini C18 5 μm 10 x 250 mm column, using a gradient of H₂O+0,1% formic (A) acid and MeOH+0,1% (B) (20 min A 80% B 20%, from 20 to 90 min increase of B until A 40% B 60%, from 90 to 110 min A 5% B 95%, from 110 to 125 min A 95% B 5%) at a flow rate of 2 mL/min. A total of 4 run were performed each one injecting 15 mg of crude. The elution was monitored with UV/vis detector λ 325nm and the fractions corresponding to each peak were collected and kept at -80⁰C and then freeze dried and analyzed by ¹H NMR. 4-*p*-coumaroylquinic acid **4a** (5mg) was obtained as a white solid. M.p. 179-182⁰C [lit. ^{Error! Bookmark not defined.} 192-193⁰C]; IR (nujol): $\tilde{\nu}$ = 3580, 3382.60, 2952.03, 1689.11, 1604.93, 1172.21, 1024.40.85, 830.63 cm⁻¹; ¹H NMR (500 MHz, CD₃OD) δ 7.73 (1H, d, *J* = 15.9 Hz, CH=CH), 7.49 (2H, d, *J* = 8.6 Hz, Ar), 6.82 (2H, d, *J* = 8.6 Hz, Ar), 6.45 (1H, d, *J* = 15.9 Hz, CH=CH), 4.81 (1H, dd, *J* = 10.0, 2.8 Hz, H-4), 4.32 (2H, m, H-3 + H-5), 2.22 (4H, m, H-2 + H-6); ¹³C NMR (126 MHz, CD₃OD) δ 177.97 (s, COO), 168.97 (s, COO), 161.25 (s, Ar), 146.73 (d, CH=CH), 131.16 (d, Ar), 127.31 (s, Ar), 116.82 (d, Ar), 115.44 (d, CH=CH), 79.26 (d, C-4), 76.95 (s, C-1), 69.65 (d, C-5), 65.69 (d, C-3), 42.64 (t, C-6), 38.49 (t, C-2). MS (ESI⁺): *m/z* [M+Na]: 361.0 [α]_D²⁰ = -28.3 (*c* 0.3, MeOH) [lit. ^{Error! Bookmark not defined.} [α]_D²⁰ = -47.3 (*c* 1.4, MeOH)]; UV (MeOH): ε₃₁₆ = 63200.

Calculations

Preliminary Molecular Mechanics calculations and HF optimizations were performed using the Spartan 14 package³⁴ which was installed on an Antec P193 V3, with two six core AMD opteron Processor 2427 2.20GHz, 4 GB RAM, 1 TB physical memory, and 64-bit Windows 7 Enterprise as operating system. Convergence criteria for geometry optimization were set as follow: energy 1.0x10⁻⁶ hartrees, gradient tolerance 3x10⁻⁴ hartrees, distance tolerance 1.2x10⁻³ Å. The DFT simulations were performed on the same machine with the Schrodinger suite of programmes using the B3LYP functional³⁵ and a localized 6-31G(d,p) basis set.

Acknowledgements

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 642014.

We are grateful to Dr. Luca Zuppiroli of the Dipartimento di Chimica Industriale "Toso Montanari", Università di Bologna, for the HRMS analysis of compound **1**.

References

- ¹ Farah, A.; De Paulis, T.; Moreira, D.; Trugo L.; Martin, P. *J. Agric. Food Chem.* **2005**, *53*, 1505-1513.
- ² Sinisi, V.; Boronova, K.; Colomban, S.; Navarini, L.; Berti, F.; Forzato, C. *Eur. J. Org. Chem.* **2014**, 1321-1326.
- ³ Monteiro, M.; Farah, A. *Food Chem.* **2012**, *134*, 611-614.
- ⁴ Clifford, M.; Knight, S.; Surucu, B.; Kuhnert, N. *J. Agric. Food Chem.* **2006**, *54*, 1957-1969.
- ⁵ Olthof, M.; Hollman, P.; Katan, M.; *J. Nutrition* **2001**, *131*(1), 66-71.
- ⁶ Perrone, D.; Farah, A.; Donangelo, C.; De Paulis, T.; Martin, P. *Food Chem.* **2008**, *106*, 859-867.
- ⁷ Farah, A.; Monteiro, R. C.; Calado, V.; Franca, A.; Trugo, L. C. *Food Chem.*, **2006**, *98*, 373-380.
- ⁸ Frank, O.; Zehentbauer, G.; Hofmann, T. *Food Res. Technol.* **2006**, *222*, 492-508.
- ⁹ Farah, A.; Monteiro M.; Donangelo, C.; Lafay, S. *J. Nutrition* **2008**, *138*, 2309-2315.
- ¹⁰ Mills, C.; Concha, M.; Mottram, D.; Gibson, G.; Spencer, J. *Food Chem.* **2013**, *141*, 3335-3340.
- ¹¹ Ludwig, I.; Sanchez, L.; Caemmerer, B.; Kroh, L. *Food Res. Int.* **2012**, *48*(1), 57-64.
- ¹² Clifford, M.; Knight, S.; Surucu, B.; Kuhnert, N. *J. Agric. Food Chem.* **2006**, *54*, 1957-1969.
- ¹³ Clifford, M.; *J. Sci. Food Agric.* **2000**, *80*, 1033-1043.
- ¹⁴ Ky, C.; Noirrot, M.; Hamon, S. *J. Agric. Food Chem.* **1997**, *45*, 786-790.
- ¹⁵ Panizzi, L.; Scarpati, M.; Oriente, G. *Gazz. Chim. Ital.* **1956**, *86*, 913-922.
- ¹⁶ Panizzi, L.; Scarpati, M.; Oriente, G. *Experientia* **1955**, *11*, 383-384.
- ¹⁷ Sefkow, M.; Kelling, A.; Schilde, U.; *Eur. J. Org. Chem.* **2001**, 2735-2742.
- ¹⁸ Sefkow, M. *Eur. J. Org. Chem.* **2001**, 1137-1141.
- ¹⁹ Dokli, I.; Navarini, L.; Hamersak, Z. *Tetrahedron: Asymmetry*, **2013**, *24*, 785-790.
- ²⁰ Haslam, E.; Haworth, R.; Makinson, G. *J. Chem. Soc.* **1961**, 5153-5156.
- ²¹ Haslam, E.; Makinson, G.; Naumann, G.; Cunningham, J. *J. Chem. Soc.*, **1964**, 2137-2146.
- ²² In the present work the IUPAC numbering system is consistently applied (i.e. in the cyclohexane ring the C atom bearing the OH on the same side as the carboxylic acid, is identified as position 5) (IUPAC Commission on the Nomenclature of Organic Chemistry (CNOC) and IUPAC-IUB Commission on Biochemical Nomenclature (CBN), *Biochem. J.* **1976**, *153*, 23-31).
- ²³ Ma, C.; Kully, M.; Khan, J.; Hattori, M.; Daneshtalab, M. *Bioorg. Med. Chem.* **2007**, *15*, 6830-6833.
- ²⁴ Criton, M.; Le Mellay-Hamon, V. *Biol. Pharm. Bull.* **2011**, *34*, 420-425.
- ²⁵ Jaiswal, R.; Patras, M.; Eravuchira, P.; Kuhnert, N. *J. Agric. Food Chem.* **2010**, *58*, 8722-8737.
- ²⁶ Rohloff, J.; Kent, K.; Postich, M.; Becker, M.; Chapman, H.; Kelly, D.; Lew, W.; Louie, M.; McGee, L.; Prisbe, E.; Schultze, L.; Yu, R.; Zhang, L. *J. Org. Chem.* **1998**, *63*, 4545-4550.
- ²⁷ De Pooter, H.; De Brucker, J.; van Sumere, C. F. *Bull. Soc. Chim. Belge*, **1976**, *85*, 663-671.
- ²⁸ Choi, Y. H.; Kim, H. K.; Linthorst, H. J. M.; Hollander, J. G.; Lefeber, A. W. M.; Erkelens, C.; Nuzillard, J.-M.; Verpoorte, R. *J. Nat. Prod.* **2006**, *69* (5), 742-748.
- ²⁹ Montchamp, J.; Tian, F.; Hart, M.; Frost, J. *J. Org. Chem.* **1996**, *61*, 3897-3899.
- ³⁰ Wolinsky, J.; Novak, R.; Vasileff, R., *J. Org. Chem.*, **1964**, *29*, 3596-3598.
- ³¹ Raheem, K.; Botting, N.; Williamson, G.; Barron, D. *Tetrahedron Lett.* **2011**, *52*, 7175-7177.
- ³² Glebocka, A.; Sicinski, R.; Plum, L.; Clagett-Dame, M.; De Luca, H. *J. Med. Chem.* **2006**, *49*, 2009-2020.
- ³³ Federspiel, M.; Fisher, R.; Hennig, M. et al, *Org. Process Res. & Dev.* **1999**, *3*, 266-274.
- ³⁴ Hehre, W. J. *A Guide to Molecular Mechanics and Quantum Chemical Calculations*; Wavefunction, Inc., **2003**.
- ³⁵ Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648-5652.