



QUEEN MARY
AND WESTFIELD COLLEGE
UNIVERSITY OF LONDON

Department of Computer Science

Research Report No. RR-00-03

ISSN 1470-5559

August 2000

Relative Definability of Boolean Functions via Hypergraphs

Antonio Bucciarelli

Pasquale Malacaria

Relative Definability of Boolean Functions via Hypergraphs

Antonio Bucciarelli

*Dipartimento di Scienze dell'Informazione, Università di Roma "La Sapienza",
via Salaria 113, 00198 Roma, Italy*

Pasquale Malacaria

*Department of Computing, Imperial College of Science Technology and Medicine,
180 Queen's Gate London SW7 2AZ, UK*

The aim of this work is to show how hypergraphs can be used as a *systematic* tool in the classification of continuous boolean functions according to their *degree of parallelism*. Intuitively f is "less parallel" than g if it can be defined by a sequential program using g as its only free variable. It turns out that the poset induced by this preorder is (as for the degrees of recursion) a sup-semilattice.

Although hypergraphs have already been used in [6] as a tool for studying degrees of parallelism, no general result relating the former to the latter has been proved in that work. We show that the sup-semilattice of degrees has a categorical counterpart: we define a category of hypergraphs such that every object "represents" a monotone boolean function; finite co-products in this category correspond to lubs of degrees. Unlike degrees of recursion, where every set has a recursive upper bound, monotone boolean functions may have no sequential upper bound. However the ones which do have a sequential upper bound can be nicely characterised in terms of hypergraphs. These subsequential functions play a major role in the proof of our main result, namely that f is less parallel than g if there exists a morphism between their associated hypergraphs.

1 Introduction

In this paper we will consider first-order continuous functions of type $\mathcal{B}^n \rightarrow \mathcal{B}$ where \mathcal{B} is the flat domain of boolean values $\{\perp, \text{tt}, \text{ff}\}$. Tuples of boolean values are ordered component-wise. Note that continuous functions of this type are just monotone functions.

Given two continuous functions f and g , we say that f is *less parallel than* g ($f \leq_{\text{par}} g$) if there exists a closed PCF-term M such that $\llbracket M \rrbracket g = f$ (where $\llbracket M \rrbracket$ denotes the interpretation of M in the standard Scott model [17])¹.

A *degree of parallelism* is a class of the equivalence relation associated with the preorder \leq_{par} . Two functions in the same class will be called *equiparallel*. The degree of a given continuous function f will be denoted by $[f]$.

We will use sometimes the expression f is *g -definable* for $f \leq_{\text{par}} g$.

The study of degrees of parallelism was pioneered by Sazonov and Tracktembrot, [16,21] who singled out some finite subsets of degrees.

In order to study \leq_{par} we introduce a category of hypergraphs. Continuous functions will be projected on the objects of this category, and hypergraph morphisms will be witnesses of \leq_{par} relations.

An informal way of gradually describing the passage from function to hypergraph is the following:

Any function is a set of pairs (argument,value): its graph.

Monotone functions on finite posets can be represented by a set of pairs (minimal argument, value): their trace (for a formal definition of trace see the next section).

In the hypergraph representations the arity of the function and the actual content of minimal arguments are forgotten. The vertexes of the hypergraph stand for minimal arguments, and the edges encode a partial information on the actual content of such minimal arguments. The values of the encoded function are recorded by coloring the vertexes.

Consider for instance the n -ary logical connective that outputs tt if all its arguments are tt and is undefined otherwise. Then the hypergraph associated to any such function is the same for all n , namely the hypergraph with a unique vertex and no arcs. Indeed any hypergraph represents infinitely many functions whereas traces are in a one-to-one correspondence with (monotone) functions.

A natural question is hence how faithful the hypergraph representation is. This question is indeed twofold, namely:

- Which properties of functions are characterised in terms of hypergraphs?

¹ Actually, $\llbracket M \rrbracket g = f$ is an abbreviation for $\llbracket M \rrbracket (\text{curry}(g)) = \text{curry}(f)$, since PCF does not have product types. We will use this abbreviation throughout the paper.

- Is it the case that two functions having the same hypergraph are equiparallel?

Concerning the first questions the results in this paper are summarised in the following table (rows stand for type of the function, column for hypergraph properties characterising that type of function)²:

Functions	Hypergraphs			
	<i>functional</i>	<i>no hyperarcs</i>	<i>no binary hyperarcs</i>	<i>only monochromatic hyperarcs</i>
<i>continuous</i>	Yes	No	No	No
<i>stable</i>	Yes	No	Yes	No
<i>sequential</i>	Yes	Yes	Yes	Yes
<i>subsequential</i>	Yes	No	No	Yes

So, for instance, a function f is stable if and only if the hypergraph H_f associated to it is functional and all its hyperarcs have at least three elements.

Concerning the second questions let us consider an example which gives some evidence of the fact that the question itself is non-trivial:

Example 1: Let us consider, for $n \in \omega$, $n \geq 1$ the monotone functions $f_n, g_n : \mathcal{B}^n \rightarrow \mathcal{B}$ defined by the following traces:

$$\text{tr}(f_n) = \{(v, \text{tt}), (\sigma^1(v), \text{tt}), \dots, (\sigma^{n-1}(v), \text{tt})\}$$

$$\text{tr}(g_n) = \{(w, \text{tt}), (\sigma^1(w), \text{tt}), \dots, (\sigma^{n-1}(w), \text{tt})\}$$

where $v = (\text{tt}, \underbrace{\perp, \dots, \perp}_{n-1})$, $w = (\perp, \underbrace{\text{tt}, \dots, \text{tt}}_{n-1})$, and $\sigma((b_1, \dots, b_{n-1}, b_n)) = (b_n, b_1, \dots, b_{n-1})$.

i.e. f_n is the function that outputs tt if it has at least one tt in its n arguments whereas g_n outputs tt if it has at least $n - 1$ tt among its arguments.

² Stable and sequential functions are introduced in section 2. Functional hypergraphs in section 3; monochromatic hypergraphs and subsequential functions in section 4.

For a given n the maps f_n and g_n are represented by the same hypergraph, namely the complete hypergraph of order n (that is the hypergraph in which all but singletons subsets of vertices are hyperarcs). Hence there is a trivial morphism (namely the identity) between the hypergraphs of f_n and g_n . However the PCF term M_n defining f_n in terms of g_n has at least $n - 1$ “nested” calls of g_n .

For example for $n = 3$ we have

$$f_3 = \lambda xyz. g_3(x g_3(\text{tt } y z) \text{tt})$$

and for $n = 4$

$$f_4 = \lambda xyzw. g_4(x g_4(y g_4(\text{tt } \text{tt } z w) \text{tt } \text{tt}) \text{tt } \text{tt})$$

■

The moral is that if we could prove that hypergraphs isomorphisms *reflect* equivalence of degrees (i.e. that functions whose hypergraphs are isomorphic are equiparallel) then we would have a simple and effective tool for the study of degrees. We will indeed prove such a result as a corollary of our main result: hypergraphs morphisms reflect \leq_{par} relations.

1.1 Related works

The study of degrees of parallelism was pioneered by Sazonov and Tracktembrot [16,21] who singled out some finite subposets of degrees. Some results on degrees are corollary of well known facts: for instance Plotkin’s full abstraction result for PCF+por implies that this poset has a top. The bottom of degrees is the set of PCF-definable functions which is fully characterised by the notion of sequentiality (in any of its formulations). Moreover Sieber’s *sequentiality relations* [18] provide a characterization of first-order degrees of parallelism and this characterization is effective: given f and g one can decide if $f \leq_{\text{par}} g$, and recently Stoughton [19] has implemented an algorithm which solves this decision problem.

Recently, Loader has shown that the PCF-definability problem, i.e. the problem of deciding if a given continuous function is PCF-definable, is undecidable [12]. As a consequence, the relation \leq_{par} is undecidable in general (at higher-order), since, if g is PCF-definable and f continuous, then f is PCF-definable if and only if $f \leq_{\text{par}} g$.

Hypergraphs for the study of degrees were first introduced in [6] where an infinite subposet of degrees was pointed out. However no precise connection between hypergraphs and monotone functions was established there. The definition of functional hypergraphs bears striking resemblance to Ehrhard's definition of *parallel* hypercoherence [8] and indeed we owe him the condition [H2'] in section 3.

2 The upper semi-lattice of degrees

Throughout this paper, we will often define boolean functions via their *trace*. The notion of trace of a function has been defined by Berry [4] and Girard [9] in the framework of stable semantics of λ -calculi. For first-order, monotone boolean functions traces are particularly easy to define. In the next paragraphs we sketch the isomorphism between traces and boolean functions, without proofs.

A $(n$ -ary) *trace* is a set $T \subseteq \mathcal{B}^n \times (\mathcal{B} \setminus \{\perp\})$ satisfying the following conditions:

- If $(w_1, b_1), (w_2, b_2) \in T$ and $w_1 \uparrow w_2$ then $b_1 = b_2$.
- If $w \in \pi_1(T)$ and $w < v$ then $v \notin \pi_1(T)$.

A n -ary trace T univoquely determines the function $f_T : \mathcal{B}^n \rightarrow \mathcal{B}$ defined by:

$$f_T(v) = \bigvee \{b \in \mathcal{B} \mid \exists w \leq v (w, b) \in T\}$$

Given a monotone function $f : \mathcal{B}^n \rightarrow \mathcal{B}$, the *trace of f* is defined by

$$\text{tr}(f) = \{(v, b) \mid v \in \mathcal{B}^n, b \in \mathcal{B}, b \neq \perp, f(v) = b \text{ and } \forall v' < v f(v') = \perp\}.$$

Traces are in one-to-one correspondence with monotone functions. It is easy to check that, given a trace T and a monotone function g , $\text{tr}(f_T) = T$ and $f_{\text{tr}(g)} = g$.

In order to introduce the first remark on degrees we recall the parallel or function *por* defined by

$$\text{por}(x, y) = \begin{cases} \text{tt} & \text{if } x = \text{tt} \text{ or } y = \text{tt} \\ \text{ff} & \text{if } x = \text{ff} \text{ and } y = \text{ff} \\ \perp & \text{otherwise.} \end{cases}$$

Fact 2 *The poset of degrees of parallelism is a sup semilattice with a bottom element (the set of PCF-definable functions) and a top element (the equivalence class of parallel or).*

Proof: The set of PCF-definable functions is the \perp of degrees by definition, whereas the fact that por is the \top of degrees, is a corollary of Plotkin's definability result [15]³.

Given $f : \mathcal{B}^n \rightarrow \mathcal{B}$ and $g : \mathcal{B}^m \rightarrow \mathcal{B}$, we define $h : \mathcal{B}^k \rightarrow \mathcal{B}$ such that $[h] = [f] \vee [g]$. Without loss of generality, let us suppose that there exists $l \geq 0$ such that $m = n - l$. Then we set $k = n + 1$, and let h be the unique function from \mathcal{B}^k to \mathcal{B} such that:

$$\begin{aligned} \text{tr}(h) = & \{((\text{tt}, x_1, \dots, x_n), b) \mid ((x_1, \dots, x_n), b) \in \text{tr}(f)\} \cup \\ & \{((\text{ff}, \underbrace{\perp, \dots, \perp}_l, x_1, \dots, x_m), b) \mid ((x_1, \dots, x_m), b) \in \text{tr}(g)\}. \end{aligned}$$

In order to prove that $[h] = [f] \vee [g]$ we have first to show that $f \leq_{\text{par}} h$ and $g \leq_{\text{par}} h$. It is easy to check that $h(\text{tt}, x_1, \dots, x_n) = f(x_1, \dots, x_n)$, and thus

$$\llbracket \lambda d \lambda x_1 \dots x_n. d \text{tt } x_1 \dots x_n \rrbracket h = f$$

and that $h(\text{ff}, y_1, \dots, y_l, x_1, \dots, x_m) = g(x_1, \dots, x_m)$, and thus

$$\llbracket \lambda d \lambda x_1 \dots x_m. d \text{ff } \underbrace{\perp \dots \perp}_l x_1 \dots x_m \rrbracket h = g.$$

Moreover, let $h' : \mathcal{B}^j \rightarrow \mathcal{B}$ be such that $f, g \leq_{\text{par}} h'$, i.e. such that there exist M, N : $\llbracket M \rrbracket h' = f$ and $\llbracket N \rrbracket h' = g$. Then it is again easy to check that

$$\llbracket \lambda d \lambda x_1 \dots x_k. \text{if } x_1 \text{ then } M g x_2 \dots x_k \text{ else } N g x_{l+2} \dots x_k \rrbracket h' = h$$

Hence $[h] = [f] \vee [g]$. ■

Given f, g as above the function h given in the proof of the proposition will be denoted by $f + g$.

The set of monotone functions which can be computed by sequential, purely functional programs is the \perp of the hierarchy of degrees, and it has been the object of a considerable amount of research. We end this section with a short overview of some of these works, pointing out some notions and results used in the rest of the paper.

³ Actually in Plotkin's original proof a *parallel if* function is used instead of *por*. For the interdefinability of the parallel "if" and "or" see [20].

The Full Abstraction problem for PCF led to the definition of classes of functions which are more constrained than the continuous ones; in particular, as we will see, stable [3] and strongly stable [5] functions have a nice characterisation in term of hypergraphs.

A continuous function $f : \mathcal{B}^n \rightarrow \mathcal{B}$ is *stable* if for all $v_1, v_2 \in \mathcal{B}^n$, if v_1 and v_2 are bounded then $f(v_1 \wedge v_2) = f(v_1) \wedge f(v_2)$ (or equivalently if for all distinct $v_1, v_2 \in \pi_1(\text{tr}(f))$, v_1 and v_2 are unbounded.)

A subset $A = \{v_1, \dots, v_k\}$ of \mathcal{B}^n is *linearly coherent* (or simply *coherent*) if

$$\forall j \ 1 \leq j \leq n \ (\perp \in \pi_j(A) \text{ or } \#\pi_j(A) = 1)$$

where $\#X$ denotes the cardinality of the set X (we use this notation throughout the paper). The set $\pi_j(A) = \{v_1^j, \dots, v_k^j\}$ is the j -th *component* of A . A subset A of \mathcal{B} is coherent if either it contains \perp or it is a singleton.

Example 3: Consider the sets $A, B \subseteq \mathcal{B}^3$ defined by

$$A = \{(\text{tt}, \text{tt}, \perp), (\text{tt}, \text{ff}, \perp), (\text{ff}, \perp, \text{tt}), (\text{ff}, \perp, \text{ff})\}$$

$$B = \{(\perp, \text{tt}, \text{ff}), (\text{ff}, \perp, \text{tt}), (\text{tt}, \text{ff}, \perp)\}$$

A is not coherent, since its first component does not contain \perp nor it is a singleton. B is coherent since all its components do contain \perp . A is the set of minimal points of the if-then-else function, which is PCF-definable; B is the set of minimal points of the so called Berry function, which is stable but not PCF-definable. ■

The set of coherent subsets of \mathcal{B}^n (resp. \mathcal{B}) is denoted $\mathcal{C}(\mathcal{B}^n)$ (resp. $\mathcal{C}(\mathcal{B})$).

Coherent sets play an important role in our description of monotone functions via hypergraphs: the vertexes of the hypergraph associated to a function f stand for the minimal points of f (i.e. the elements of the first projection of the trace of f), and a set $\{v_1, \dots, v_k\}$ of vertexes is an arc if and only if the set of the corresponding minimal points of f is coherent. We will often use the following simple properties of traces and coherence:

Fact 4 – *If $A \in \mathcal{C}(\mathcal{B}^n)$ and B is an Egli-Milner lower bound of A (that is if $\forall x \in A \exists y \in B \ y \leq x$, $\#B \leq \#A$ and $\forall y \in B \exists x \in A \ y \leq x$) then $B \in \mathcal{C}(\mathcal{B}^n)$.*

- If $f : \mathcal{B}^n \rightarrow \mathcal{B}$ is a monotone function, $A \subseteq \mathcal{B}^n$, and $f(A) \subseteq \mathcal{B} \setminus \{\perp\}$, then there exists an Egli-Milner lower bound B of A such that $B \subseteq \pi_1(\text{tr}(f))$, $\#B \leq \#A$ and $f(B) = f(A)$.

The first item is easy to check (a proof can be found in [5]); the second one is an immediate consequence of the definition of trace.

Definition 5 A continuous function $f : \mathcal{B}^n \rightarrow \mathcal{B}$ is linearly strongly stable (or simply strongly stable) if for any $A \in \mathcal{C}(\mathcal{B}^n)$

- $f(A) \in \mathcal{C}(\mathcal{B})$.
- $f(\wedge A) = \wedge f(A)$.

Example 6: Let us see how strong stability rules out the Berry function $g : \mathcal{B}^3 \rightarrow \mathcal{B}$ defined by

$$\text{tr}(g) = \{((\perp, \text{tt}, \text{ff}), \text{tt}), ((\text{ff}, \perp, \text{tt}), \text{tt}), ((\text{tt}, \text{ff}, \perp), \text{tt})\}$$

As we have seen in example 3 the set B of minimal points of g is coherent, but $\wedge g(B) = \text{tt} \neq g(\wedge(B)) = \perp$. Hence g is not strongly stable. ■

Even though the model of strongly stable functions is not fully abstract for PCF, i.e. there exist strongly stable functionals which are not PCF-definable, see [5], strong stability does capture the notion of sequentiality, or PCF-definability, at first-order. In the following proposition “sequential” stands for “Kahn-Plotkin sequential” [11], “Milner sequential” [13] or “Vuillemin sequential” [22], since all these notions coincide for first-order functions.

Proposition 7 Let $f : \mathcal{B}^n \rightarrow \mathcal{B}$ be a monotone function. The following are equivalent:

- f is strongly stable.
- f is PCF-definable.
- f is sequential.

A proof can be found in [6] and in [2]. The original proof of “sequential \Leftrightarrow PCF-definable” is in [4].

Actually there exist several alternative characterizations of the notion of PCF-definability for first-order functions, for instance Sieber’s logically sequential functions [18] and Colson-Ehrhard’s hereditarily sequential ones [7]. Of course any fully abstract model of PCF [1], [10],[14] provides *a fortiori* a characterization of PCF-definability for monotone, first order functions.

3 Hypergraphs and monotone functions

Definition 8 A colored hypergraph $H = (V_H, A_H, C_H)$ is given by a finite set V_H of vertices, a set $A_H \subseteq \{A \subseteq V_H \mid \#A \geq 2\}$ of (hyper)arcs and a coloring function $C_H : V_H \rightarrow \{\text{black}, \text{white}\}$.

As a first approximation a map between two hypergraphs is a set-theoretic map from vertices to vertices which preserves hyperarcs; concerning colours, several notions are possible: one extreme is to ask for the preservation of colours; on the other hand a more liberal requirement is to say that the images of “adjacent” vertices of different colours have different colours (think of “adjacent” as “being in the same hyperarc”).

Formally we consider two notions of morphisms on hypergraphs:

A *weak* morphism from a hypergraph H to a hypergraph H' is a function $m : V_H \rightarrow V_{H'}$ such that:

- For all $A \subseteq V_H$, if $A \in A_H$ then $m(A) \in A_{H'}$.
- for all $X \in A_H$, if $x, x' \in X$ and $C_H(x) \neq C_H(x')$ then $C_{H'}(m(x)) \neq C_{H'}(m(x'))$.

A *strong* morphism is more restrictive on colours: we require that for all $x \in V_H$, $C_H(x) = C_{H'}(m(x))$.

A *sub-hypergraph* H' of a hypergraph H has as set of vertices $V_{H'}$ a subset of V_H and as hyperarcs those of H whose vertices belong to H' . Colours are given by restriction.

Note that set theoretical inclusions are both weak and strong morphisms with this notion of sub-hypergraph.

We will restrict our attention on a particular class of hypergraphs which turns out to be in a very precise relationship with monotone functions.

A *functional hypergraph* is an hypergraph H such that:

- H1 : If $\{x, y\} \in A_H$ then $C_H(x) = C_H(y)$.
- H2 : If $X \subseteq V_H$, such that $\#X \geq 2$, is not a hyperarc then there exists a partition X_1, X_2 of X such that for all $Y \subseteq X$ if $Y \cap X_1 \neq \emptyset, Y \cap X_2 \neq \emptyset$ then Y is not a hyperarc.

Condition [H2] can be equivalently and more synthetically expressed as follows:

- H2' : If X_1, X_2 are hyperarcs and $X_1 \cap X_2 \neq \emptyset$ then $X_1 \cup X_2$ is an hyperarc.

Lemma 9 *The conditions [H2] and [H2'] above are equivalent.*

Proof: [H2] \Rightarrow [H2'] is easy to prove. Conversely let $X \subseteq V_H$ be such that $\#X \geq 2$ and $X \notin A_H$. If there is no hyperarc included in X , then any partition satisfies [H2]. Otherwise let $Y \subset X$ be a maximal hyperarc included in X , i.e. a (*a fortiori* proper) subset of X such that $Y \in A_H$ and for all $Z \subseteq X$, if $Z \in A_H$ then $\#Z \leq \#Y$. By [H2'] and by maximality of Y we have that for all $Z \subseteq X$, if $Z \cap Y \neq \emptyset$ and $Z \cap (X \setminus Y) \neq \emptyset$ then $Z \notin A_H$. Hence, the partition $Y, X \setminus Y$ satisfies [H2]. \blacksquare

It is trivial to check that a sub-hypergraph of a functional hypergraph is functional.

We are now ready to define our categories of interest: $\mathcal{SH}, \mathcal{WH}$

object $\mathcal{SH} = \text{object } \mathcal{WH} = \text{Functional Hypergraphs.}$

arrows $\mathcal{SH} = \text{Strong Morphisms.}$

arrows $\mathcal{WH} = \text{Weak Morphisms.}$

(it is trivial indeed to check that in both cases we have a category).

Definition 10 *Let $f : \mathcal{B}^n \rightarrow \mathcal{B}$ be the n -ary function defined by $\text{tr}(f) = \{(v_1, b_1), \dots, (v_k, b_k)\}$. The hypergraph H_f is defined by*

- $V_{H_f} = \{1, 2, \dots, k\}$.
- $A_{H_f} = \{\{i_1, i_2, \dots, i_l\} \subseteq V_{H_f} \mid l \geq 2 \text{ and } \{v_{i_1}, v_{i_2}, \dots, v_{i_l}\} \in \mathcal{C}(\mathcal{B}^n)\}$.
- $C_{H_f}(i) = \text{if } b_i \text{ then white else black.}$

Example 11: Consider the Berry function $g : \mathcal{B}^3 \rightarrow \mathcal{B}$ defined in example 6 and the parallel-or function $\text{por} : \mathcal{B}^2 \rightarrow \mathcal{B}$ defined in section 2, whose traces are respectively

$$\text{tr}(g) = \{((\perp, \text{tt}, \text{ff}), \text{tt}), ((\text{ff}, \perp, \text{tt}), \text{tt}), ((\text{tt}, \text{ff}, \perp), \text{tt})\}$$

$$\text{tr}(\text{por}) = \{((\perp, \text{tt}), \text{tt}), ((\text{tt}, \perp), \text{tt}), ((\text{ff}, \text{ff}), \text{ff})\}$$

We have:

$$H_g = (\{1, 2, 3\}, \{\{1, 2, 3\}\}, C_{H_g}(1) = C_{H_g}(2) = C_{H_g}(3) = \text{white})$$

$$H_{\text{por}} = (\{1, 2, 3\}, \{\{1, 2\}, \{1, 2, 3\}\},$$

$$C_{H_{\text{por}}}(1) = C_{H_{\text{por}}}(2) = \text{white}, C_{H_{\text{por}}}(3) = \text{black}$$

The map $\alpha : H_g \rightarrow H_{\text{por}}$ defined by $\alpha(1) = \alpha(2) = 1$, $\alpha(3) = 2$ is a (strong) morphism. ■

Proviso 12: The vertexes of H_f are in one-to-one correspondence with $\pi_1(\text{tr}(f))$. We could have turned this correspondence into an identity, by stipulating that $V_{H_f} = \pi_1(\text{tr}(f))$. However, since we will prove that whenever H_f and H_g are (weakly or strongly) isomorphic, f and g are equiparallel, and since hypergraph isomorphisms are clearly independent from vertexes' names, we do prefer to keep this identity implicit. Nevertheless in several proofs of the following sections, given H_f we will need to explicitly refer to minimal points of f (i.e. to elements of $\pi_1(\text{tr}(f))$). Formally, given a functional hypergraph H , there exists a family of functions $\{h_f\}_{f \in \{g \mid H_g = H\}} : V_H \rightarrow \bigcup_{n \in \omega} \mathcal{B}^n$ such that $h_f(V_H) = \pi_1(\text{tr}(f))$.

For the sake of simplicity we will omit h_f whenever possible, and in particular we will feel free of considering the vertexes of H_f as if they were labelled by $\pi_1(\text{tr}(f))$.

Also, in definition 10, the hypergraph H_f associated to f is defined up to (strong) isomorphism, since the order of $\text{tr}(f)$'s elements is not determined. We could introduce a canonical numbering of the elements of \mathcal{B}^n to overcome this problem, but again, since we will show eventually that (even weak) isomorphisms reflect equality of degree of parallelism, it is satisfactory for us to work with hypergraphs defined up to isomorphisms. ■

We can observe that for any monotone function $f : \mathcal{B}^n \rightarrow \mathcal{B}$, the hypergraph H_f is functional: the requirement H1 is satisfied by H_f since if two minimal points v_1, v_2 of f are coherent, then they are bounded (note that this is true only for binary sets), hence $f(v_1) = f(v_2)$. H2 is verified as well, since if a set $A = \{v_1, \dots, v_k\}$, $k \geq 2$ of minimal points of f is not coherent, then there exists $1 \leq j \leq n$ such that the j -th component $\{v_1^j, \dots, v_k^j\}$ of A is $\{\text{tt}, \text{ff}\}$. Hence the partition of $\{1, \dots, k\}$ given by $\{\{i \mid v_i^j = \text{tt}\}, \{i \mid v_i^j = \text{ff}\}\}$ satisfies H2. Actually the converse does hold, too:

Proposition 13 *Given an hypergraph H there exists a monotone function $f : \mathcal{B}^n \rightarrow \mathcal{B}$, for some n , such that H_f is strongly isomorphic to H if and only if H is a functional hypergraph.*

Proof: The function F_H associated to a functional hypergraph $H = (V_H, A_H, C_H)$ is defined as follows: $F_H : \mathcal{B}^n \rightarrow \mathcal{B}$ where $n = \#V_H + \#\overline{A_H}$ with

$$\overline{A_H} = \{B \subseteq V_H \mid \#B \geq 2 \text{ and } B \notin A_H\}.$$

The trace of F_H has $m = \#V_H$ elements. We fix enumerations v_1, \dots, v_m for the set V_H and B_1, \dots, B_l for the set $\overline{A_H}$. For all $B_i \in \overline{A_H}$ let (B_i^1, B_i^2) a partition of B_i satisfying the condition [H2] (at least one such partition does exist, since H is functional).

The i -th element of $\text{tr}(F_H)$ is then defined as follows:

$$((\underbrace{\perp, \dots, \perp}_{i-1}, \text{tt}, \underbrace{\perp, \dots, \perp}_{m-i}, b_i^1, \dots, b_i^l), c_i)$$

where

$$b_i^j = \begin{cases} \text{tt} & \text{if } v_i \in B_j^1 \\ \text{ff} & \text{if } v_i \in B_j^2 \\ \perp & \text{otherwise} \end{cases}$$

and

$$c_i = \begin{cases} \text{tt} & \text{if } C_H(v_i) = \text{white} \\ \text{ff} & \text{if } C_H(v_i) = \text{black}. \end{cases}$$

We leave to the reader to check that F_H is a monotone function whose hypergraph is (strongly isomorphic to) H . ■

It is easy to see that the function F_{H_f} bears in general no resemblance with f for example if $f = \text{por} : \mathcal{B}^2 \rightarrow \mathcal{B}$ then $F_{H_f} : \mathcal{B}^5 \rightarrow \mathcal{B}$. The function F_H associated with a functional hypergraph H is not uniquely specified, since it depends on the choice of the partitions (B_i^1, B_i^2) , $1 \leq i \leq l$ in the construction above.

We end this section with a nice property of the categories $\mathcal{SH}, \mathcal{WH}$.

Proposition 14 $\mathcal{SH}, \mathcal{WH}$ have coproducts.

Proof: Let us define the binary coproducts: given H, H' let H'' be the hypergraph given by the disjoint union of vertices of H, H' , the disjoint union of hyperarcs of H, H' and the disjoint union of the colouring maps of H, H' . Then H'' is a functional hypergraph (condition H1 is trivial and condition H2 is trivially checked as well by using $H2'$).

The inclusion maps h (resp h') from H (resp H') to H'' provide the injections. Finally is easy to see that any pair of maps f, f' from H (resp H') to H''' factorize through H'' , both in \mathcal{SH} and in \mathcal{WH} . ■

Note that categorical coproduct and l.u.b. of degrees are related in the following sense:

Fact 15 *The coproduct $H_f \oplus H_g$ (in both categories $\mathcal{SH}, \mathcal{WH}$) is isomorphic the hypergraph of $f + g$.*

Proof: By definition the trace of $f + g$ has $l + r$ elements with l (resp r) being the number of element in the trace of f (resp g); this means that H_{f+g} has as vertices the disjoint union of vertices of H_f, H_g . By the definition of trace of $f + g$ is also clear that the colouring map of H_{f+g} is the disjoint union of the maps in H_f, H_g .

The only thing we are left to check is hence the hyperarcs. Again by definition of trace of $f + g$ and by definition of coherence it is easy to check that a coherent subset of trace of f (resp of trace of g) is a coherent subset $\text{tr}(f + g)$. For the opposite direction note that by the definition of coherence a coherent subset of $\text{tr}(f + g)$ cannot contain elements from both $\text{tr}(f)$ and $\text{tr}(g)$ (again by definition of $\text{tr}(f + g)$ because of the first argument). This implies that the hyperarcs of H_{f+g} are indeed the disjoint union of the hyperarcs of H_f and H_g . ■

3.1 Relating hypergraphs and degrees

First we can observe how clearly hypergraphs classify PCF-definable and stable functions versus general monotone functions.

Fact 16 *Let $f : \mathcal{B}^n \rightarrow \mathcal{B}$ be a continuous function: f is stable if and only if H_f has no binary hyperarcs. It is strongly stable if and only if H_f has no hyperarcs.*

Proof: Let us prove the statement concerning strongly stable functions: given $f : \mathcal{B}^n \rightarrow \mathcal{B}$, if H_f has a hyperarc $A = \{v_1, \dots, v_k\}$ (see proviso 12), then by definition $\{v_1, \dots, v_k\} \in \mathcal{C}(\mathcal{B}^n)$. Now either all the vertexes of A have the same colour in H_f , and hence $f(\bigwedge A) < \bigwedge f(A)$, or they have not, hence $f(A) \notin \mathcal{C}(\mathcal{B})$. In both cases f is not strongly stable.

Conversely if H_f has no hyperarc, let $A \in \mathcal{C}(\mathcal{B}^n)$ be such that $\perp \notin f(A)$ (otherwise $f(A) \in \mathcal{C}(\mathcal{B})$ and $f(\bigwedge A) = \bigwedge f(A)$ holds trivially). By fact 4, there exists an Egli-Milner lower bound B of A such that $B \subseteq \pi_1(\text{tr}(f))$ and $f(A) = f(B)$. Since B is coherent and H_f has no hyperarc, $\#B = 1$, hence $f(A) \in \mathcal{C}(\mathcal{B})$ and $f(\bigwedge A) = \bigwedge f(A)$, since it is easy to see that $\bigwedge A$ is above the element of B .

The proof of the statement concerning stable functions is a particular case of the one above, with $k = 2$ (one needs here that $\#B \leq \#A$, in fact 4). ■

Hypergraphs have already been used in [6] in order to show that the poset of degrees is highly non-trivial; in particular it contains both infinite (ascending and descending) chains and infinite anti-chains. Bucciarelli defined a class of hypergraphs as follows.


Definition 17 *Given two natural numbers $m \geq n \geq 3$, let $H(n, m)$ be the hypergraph defined by:*

$$H(n, m) = (\{1, 2, \dots, m\}, \{A \subseteq \{1, 2, \dots, m\} \mid \#A \geq n\}, \text{for all } i \ C(i) = \text{white})$$

It is easy to check that the $H(n, m)$'s are functional hypergraphs. Let's call \mathcal{SH}' the full subcategory of \mathcal{SH} whose objects are (strongly isomorphic to) the $H(n, m)$. The main result of [6] is then:

Proposition 18 *Let f, g be such that H_f, H_g are objects of \mathcal{SH}' ; then $\mathcal{SH}'(H_f, H_g) \neq \emptyset$ iff $f \leq_{\text{par}} g$.*

In the following picture, $f(n, m)$ stands for a function such that $H_{f(n, m)}$ is weakly isomorphic to $H(n, m)$ (a canonical choice for the $f(n, m)$'s is presented in [6]), and arrows denote \leq_{par} relations:



4 Subsequential functions

A monotone function $f : \mathcal{B}^n \rightarrow \mathcal{B}$ is *subsequential* if it is extensionally upper bounded by a strongly stable function. As shown in proposition 20 subsequential functions correspond to hypergraphs with monochromatic hyperarcs and to functions preserving linear coherence. Such a class of functions admits hence a natural characterisation in order theoretic, graph theoretic and algebraic terms. Moreover, thanks to their properties subsequential functions will be an important combinatorial tool in our work.

Lemma 19 *Let $\{B_x\}_{x \in X}$ (X a non-empty set of indices) be such that $\forall x \in X, B_x \in \mathcal{C}(\mathcal{B}^n)$ and $A = \{\bigwedge B_x \mid x \in X\} \in \mathcal{C}(\mathcal{B}^n)$. Then $\bigcup_{x \in X} B_x \in \mathcal{C}(\mathcal{B}^n)$.*

Proof: Suppose that $Y = \bigcup_{x \in X} B_x \notin \mathcal{C}(\mathcal{B}^n)$; then there exists a component $1 \leq j \leq n$ and a partition (Y_1, Y_2) of Y such that for all $y_1 \in Y_1, (y_1)^j = \text{tt}$ and for all $y_2 \in Y_2, (y_2)^j = \text{ff}$.

It is easy to see that $\forall x \in X, B_x \subseteq Y_1$ or $B_x \subseteq Y_2$; hence if $a = \bigwedge B_x$ we get

- $a^j = \text{tt}$ if $B_x \subseteq Y_1$.
- $a^j = \text{ff}$ if $B_x \subseteq Y_2$.

We hence deduce a non-trivial partition (A_1, A_2) of A such that $a \in A_1$ iff $a^j = \text{tt}$ and $a \in A_2$ iff $a^j = \text{ff}$. This is a contradiction since $A \in \mathcal{C}(\mathcal{B}^n)$. ■

Proposition 20 *Let $f : \mathcal{B}^n \rightarrow \mathcal{B}$ be a monotone function. The following are equivalent:*

- 1 For all $A \subseteq \text{tr}(f)$, if $\pi_1(A) \in \mathcal{C}(\mathcal{B}^n)$ then $\pi_2(A) \in \mathcal{C}(\mathcal{B})$ ⁴
- 2 For all $A \in \mathcal{C}(\mathcal{B}^n)$, $f(A) \in \mathcal{C}(\mathcal{B})$. (i.e. f preserves the linear coherence of \mathcal{B}^n .)
- 3 f is subsequential.
- 4 If $X \in A_{H_f}$ then for all $x, y \in X, C_{H_f}(x) = C_{H_f}(y)$ (i.e. X is monochromatic).

Proof:

1 \Rightarrow 2: Let $A \in \mathcal{C}(\mathcal{B}^n)$ be such that $\perp \notin f(A)$ (otherwise $f(A) \in \mathcal{C}(\mathcal{B})$). By fact 4 there exists $B \subseteq \text{tr}(f)$ such that $\pi_1(B)$ is an Egli-Milner lower bound of A , and $\pi_2(B) = f(A)$. Since $\pi_1(B)$ is coherent (fact 4) we are done.

⁴ Since by definition of trace $\perp \notin \pi_2(A)$, $\pi_2(A) \in \mathcal{C}(\mathcal{B})$ if and only if $\pi_2(A)$ is a singleton

2 \Rightarrow 3: We have to define a strongly stable upper bound of f . Let $\bar{f} : \mathcal{B}^n \rightarrow \mathcal{B}$ be the function defined as follows:

$$\bar{f}(x) = \bigvee_{A \in \mathcal{C}(\mathcal{B}^n), x \geq \bigwedge A} (\bigwedge_{y \in A} f(y)).$$

First of all we have to show that \bar{f} is a function, i.e. that, given $x \in \mathcal{B}^n$, if $A, B \in \mathcal{C}(\mathcal{B}^n)$ are such that $x \geq \bigwedge A, \bigwedge B$, then $\bigwedge f(A)$ and $\bigwedge f(B)$ are bounded (this is sufficient since \mathcal{B} is clearly a *coherent* bounded complete cpo, i.e. any set of pairwise bounded boolean values is bounded, and hence has a l.u.b.). If A and B are as above, let us suppose, without loss of generality, that $\bigwedge f(A) = \mathbf{tt}$ and $\bigwedge f(B) = \mathbf{ff}$. Since $C = \{\bigwedge A, \bigwedge B\}$ is Egli-Milner smaller than $\{x\}$, which is coherent, C is coherent (see fact 4), hence by lemma 19 $A \cup B \in \mathcal{C}(\mathcal{B}^n)$. Since $\bigwedge f(A) = \mathbf{tt}$ and $\bigwedge f(B) = \mathbf{ff}$ we conclude that $f(A \cup B) = \{\mathbf{tt}, \mathbf{ff}\} \notin \mathcal{C}(\mathcal{B})$, hence f does not preserve $\mathcal{C}(\mathcal{B}^n)$. Since we know that f does preserve $\mathcal{C}(\mathcal{B}^n)$, we conclude that \bar{f} is well defined.

Moreover \bar{f} is clearly monotone, and it is an upper bound of f since for any $x \in \mathcal{B}^n$, $\{x\} \in \mathcal{C}(\mathcal{B}^n)$.

In order to prove that \bar{f} is strongly stable, given $A \in \mathcal{C}(\mathcal{B}^n)$, let us prove that (1) $\bar{f}(A) \in \mathcal{C}(\mathcal{B})$ and (2) $\bar{f}(\bigwedge A) = \bigwedge \bar{f}(A)$.

(1) If $\perp \in \bar{f}(A)$ then $\bar{f}(A) \in \mathcal{C}(\mathcal{B})$. Let us suppose that $\perp \notin \bar{f}(A)$. In this case, by definition of \bar{f} , for any $x \in A$ there exists $B_x \in \mathcal{C}(\mathcal{B}^n)$ such that $\bigwedge B_x \leq x$ and $\bigwedge f(B_x) > \perp$. Since $\{\bigwedge B_x \mid x \in A\}$ vis Egli-Milner smaller than A , we conclude as above by fact 4 and lemma 19, that $\bigcup_{x \in A} B_x \in \mathcal{C}(\mathcal{B}^n)$. Hence $f(\bigcup_{x \in A} B_x) \in \mathcal{C}(\mathcal{B})$. Now since for all $x \in A$ $\bar{f}(x) = \bigwedge f(B_x) > \perp$, we have $\bar{f}(A) = \{\bigwedge f(B_x) \mid x \in A\} = f(\bigcup_{x \in A} B_x) \in \mathcal{C}(\mathcal{B})$ and we are done.

(2) Since \bar{f} is monotone, $\bar{f}(\bigwedge A) \leq \bigwedge \bar{f}(A)$. Let $\bigwedge \bar{f}(A) = b > \perp$, and for any $x \in A$ let B_x be as above, that is $B_x \in \mathcal{C}(\mathcal{B}^n)$, $\bigwedge B_x \leq x$ and $\bigwedge f(B_x) = b > \perp$. Again we have that $D = \bigcup_{x \in A} B_x \in \mathcal{C}(\mathcal{B}^n)$. Moreover $\bigwedge(D) \leq \bigwedge A$, since for any x in A , $\bigwedge B_x \leq x$, hence by definition of \bar{f} , $\bar{f}(\bigwedge A) \geq \bigwedge f(D) = b$, and we are done.

3 \Rightarrow 4: If $X \in A_{H_f}$ and $x, y \in X$ are such that $C_{H_f}(x) \neq C_{H_f}(y)$ then we can find a subset A of $\text{tr}(f)$ such that $\pi_1(A) \in \mathcal{C}(\mathcal{B}^n)$ and $\pi_2(A) \notin \mathcal{C}(\mathcal{B})$; it is clear then that any extensional upper bound of f will not preserve the coherence on $\pi_1(A)$ and henceforth will not be strongly stable.

4 \Rightarrow 1: Immediate by definition of H_f .

■

We can observe that Berry's function g is subsequential, whereas por is not (see example 11).

Given a set $A = \{v_1, \dots, v_k\} \subseteq \mathcal{B}^n$, there exist in general a number of functions whose minimal points are exactly the elements of A . For instance, if the v_i are pairwise unbounded, there exist 2^k such functions. The following lemma states that, among these functions, the subsequential ones are those whose degree of parallelism is minimal.

Lemma 21 *Let $f, g : \mathcal{B}^n \rightarrow \mathcal{B}$ be such that g is subsequential and $\pi_1(\text{tr}(f)) = \pi_1(\text{tr}(g))$. Then $g \leq_{\text{par}} f$.*

Proof: Let M be a PCF term which defines the sequential upper bound \bar{g} of g , defined as in proposition 20.

Let us define $g_0 : \mathcal{B}^n \rightarrow \mathcal{B}$ by

$$g_0 = [\lambda f \lambda x_1 \dots x_n. \text{if } f x_1 \dots x_n \text{ then } M x_1 \dots x_n \text{ else } M x_1 \dots x_n] f$$

If we prove that $g_0 = g$ we are done. Let $\bar{a} = (a_1, \dots, a_n) \in \mathcal{B}^n$, and suppose $g(\bar{a}) = b \neq \perp$; then $f(\bar{a}) \neq \perp$ and $\bar{g}(\bar{a}) = b$. Hence $g_0(\bar{a}) = b$. Conversely if $g_0(\bar{a}) = b \neq \perp$ then $f(\bar{a}) \neq \perp$ and hence $g(\bar{a}) \neq \perp$ as well. Since $g(\bar{a}) \leq \bar{g}(\bar{a}) = b$, we get $g(\bar{a}) = b = g_0(\bar{a})$ and we are done. \blacksquare

Our main result of section 5 is that, if there exists a morphism $\alpha : H_f \rightarrow H_g$, then $f \leq_{\text{par}} g$. The following lemma introduces a key notion towards that result, namely the one of *slice function*. The idea is the following: in order to reduce $f : \mathcal{B}^m \rightarrow \mathcal{B}$ to $g : \mathcal{B}^n \rightarrow \mathcal{B}$ we start by transforming the minimal points of f into the ones of g . This amounts to defining a function from \mathcal{B}^m to \mathcal{B}^n , that we describe as a set of functions $f_1, \dots, f_n : \mathcal{B}^m \rightarrow \mathcal{B}$. If these functions are g -definable, then we can already g -define a function which converges if and only if f converges, namely

$$h = \lambda x_1 \dots x_m. g(f_1 \bar{x}) \dots (f_n \bar{x})$$

and we are left with the problem of forcing h to agree with f whenever it converges.

For the time being we show that, if the f_i 's are defined via a hypergraph morphism $\alpha : H_f \rightarrow H_g$, then they are subsequential, hence "relatively simple".

Lemma 22 *Let $f : \mathcal{B}^m \rightarrow \mathcal{B}$, $g : \mathcal{B}^n \rightarrow \mathcal{B}$ be monotone functions and $\alpha : H_f \rightarrow H_g$ be a weak morphism. For $1 \leq i \leq n$ let $f_i : \mathcal{B}^m \rightarrow \mathcal{B}$ be the function*

defined by ⁵

$$\text{tr}(f_i) = \{(v, \alpha(v)^i) \mid v \in \pi_1(\text{tr}(f)), \alpha(v)^i \neq \perp\}$$

Then for all $A \subseteq \text{tr}(f_i)$, if $\pi_1(A) \in \mathcal{C}(\mathcal{B}^m)$ then $\pi_2(A) \in \mathcal{C}(\mathcal{B})$.

(we will call f_i the i th-slice of f following f and α)

Note that the f_i 's, $1 \leq i \leq n$, defined above are such that for all $v \in V_{H_f}$ $\alpha(v)^i = f_i(v)$.

Proof: It is easy to see that the f_i 's are well defined. Let A be a subset of $\text{tr}(f_i)$ such that $\pi_1(A)$ is coherent. If $\#A = 1$ then $\pi_2(A) \in \mathcal{C}(\mathcal{B})$ holds trivially. Otherwise, by definition of f_i we know that for any $v \in \pi_1(A)$, $\alpha(v)^i \neq \perp$. Moreover $\alpha(\pi_1(A)) \in \mathcal{C}(\mathcal{B}^n)$, since α preserves hyperarcs. Hence we conclude that for all $v, v' \in \pi_1(A)$, $\alpha(v)^i = \alpha(v')^i$, i.e. that $\pi_2(A) = \{\alpha(v)^i \mid v \in \pi_1(A)\} \in \mathcal{C}(\mathcal{B})$.

■


By proposition 20 and lemma 22 we get:

Corollary 23 *Let $f : \mathcal{B}^m \rightarrow \mathcal{B}$, $g : \mathcal{B}^n \rightarrow \mathcal{B}$ be monotone functions and $\alpha : H_f \rightarrow H_g$ be a weak morphism. All the slices of g following f and α are subsequential.*

Example 24: Berry's function g , defined in example 6, is por-definable, as is any other monotone function. Let us define a morphism $\alpha : H_g \rightarrow H_{\text{por}}$, and see how the construction of the two slices of por following g and α provides directly a way of constructing the PCF-term defining g with respect to por. Let $v_1 = (\perp, \text{tt}, \text{ff})$, $v_2 = (\text{ff}, \perp, \text{tt})$ and $v_3 = (\text{tt}, \text{ff}, \perp)$ be the minimal points of g and $w_1 = (\perp, \text{tt})$, $w_2 = (\text{tt}, \perp)$ and $w_3 = (\text{ff}, \text{ff})$ those of por. It is easy to check that the function $\alpha : V_{H_g} \rightarrow V_{H_{\text{por}}}$ defined by $\alpha(v_1) = \alpha(v_2) = w_1$ and $\alpha(v_3) = w_2$ is a (strong) morphism from H_g to H_{por} .

The morphism α defines the map from $\pi_1(\text{tr}(g))$ into $\pi_1(\text{tr}(\text{por}))$ shown in the following picture:

⁵ see proviso 12.



The corresponding slice functions $f_1, f_2 : \mathcal{B}^3 \rightarrow \mathcal{B}$ are then defined by:

$$\text{tr}(f_1) = \{((\text{tt}, \text{ff}, \perp), \text{tt})\}$$

$$\text{tr}(f_2) = \{((\perp, \text{tt}, \text{ff}), \text{tt}), ((\text{ff}, \perp, \text{tt}), \text{tt})\}$$

Both f_1 and f_2 are sequential, hence PCF-definable. For example the following terms M_1, M_2 define f_1, f_2 respectively:

$$M_1 = \lambda x y z. \text{if } x \text{ then (if } y \text{ then } \perp \text{ else tt) else } \perp$$

$$M_2 = \lambda x y z. \text{if } z \text{ then (if } x \text{ then } \perp \text{ else tt) else (if } y \text{ then tt else } \perp)$$

The pair (M_1, M_2) realizes a sequential transformation of the minimal points of g onto (some of) the minimal points of f . This allows to construct a term M defining g with respect to por as follows:

$$M = \lambda f \lambda x y z f(M_1 x y z) (M_2 x y z)$$

It is easy to check that $\llbracket M \rrbracket_{\text{por}} = g$.

■

The theorem of the following section generalizes the situation above: we show that, given a (weak) morphism $\alpha : H_f \rightarrow H_g$, the slices of g following f and α are g -definable (even if in general they are not sequential), and this is enough to construct a PCF-term which g -defines f .

5 Hypergraph morphisms and degrees

Theorem 25 *Let $f : \mathcal{B}^l \rightarrow \mathcal{B}$, $g : \mathcal{B}^m \rightarrow \mathcal{B}$ be monotone functions. If $\mathcal{WH}(H_f, H_g) \neq \emptyset$ then $f \leq_{\text{par}} g$.*

Proof:

Let $\alpha : H_f \rightarrow H_g$ be a weak morphism. We prove the theorem by induction on $k = \#\text{tr}(f)$.

If $k = 1$ f is sequential (strongly stable), hence PCF-definable, and $f \leq_{\text{par}} g$ holds trivially.

Suppose now $k = n + 1$; we reason by cases on the structure of H_f :

- $V_{H_f} \notin A_{H_f}$: this means that there exists a sequentiality index for f , that is a component of $\pi_1(\text{tr}(f))$ which is not a singleton and which does not contain \perp ; let i be such a component. Define

$$M = \lambda g \lambda \bar{x}. \text{if } x_i \text{ then } M_{\text{tt}} g \bar{x} \text{ else } M_{\text{ff}} g \bar{x}$$

where M_ρ , $\rho = \text{tt}, \text{ff}$, is the term g -defining the sub-function f_ρ of f such that $\pi_i(\pi_1(\text{tr}(f_\rho))) = \{\rho\}$. The terms M_ρ do exist by inductive hypothesis: $\#\text{tr}(f_\rho) < \#\text{tr}(f)$, and $\mathcal{WH}(H_{f_\rho}, H_g) \neq \emptyset$ since the restriction of α to H_{f_ρ} is a morphism.

It is easy to check that M g -defines f .

- $V_{H_f} \in A_{H_f}$:
Let f_i , $1 \leq i \leq m$, be the i th-slice of g following f and α , and now define \hat{f}_i as

$$\hat{f}_i = \begin{cases} f_i & \text{if } \#\text{tr}(f_i) < \#\text{tr}(f) \\ \lambda \bar{x}. v \text{ for } v \in \pi_2(\text{tr}(f_i)) & \text{otherwise} \end{cases}$$

The \hat{f}_i 's are well defined, since if $\#\text{tr}(f_i) = \#\text{tr}(f)$ then $\#\pi_2(\text{tr}(f_i)) = 1$, V_{H_f} being a hyperarc and f_i subsequential.

Let us prove that the \hat{f}_i 's are g -definable. The only case to be checked is $\hat{f}_i = f_i$ in the previous definition, since $\lambda \bar{x}. v$ is PCF-definable.

Since the f_i 's are subsequential, by lemma 21 $f_i \leq_{\text{par}} f^i$, where $\text{tr}(f^i) = \{v \in \text{tr}(f) \mid \pi_1(v) \in \pi_1(\text{tr}(f_i))\}$. Now $\#\text{tr}(f^i) < \#\text{tr}(f)$, and, as above, $\mathcal{WH}(H_{f^i}, H_g) \neq \emptyset$. Hence by inductive hypothesis $f^i \leq_{\text{par}} g$, and finally $f_i \leq_{\text{par}} g$ by transitivity of \leq_{par} . Let M_i be a term g -defining \hat{f}_i .

Before constructing a term M g -defining f let us prove that we can already g -define a "convergence test" for f , i.e. that for all $\bar{x} = (x_1, \dots, x_l) \in \mathcal{B}^l$

$$f(\bar{x}) \neq \perp \Leftrightarrow g([M_1]g\bar{x}, \dots, [M_m]g\bar{x}) \neq \perp$$

The direction \Rightarrow is trivial, since the \hat{f}_i 's are upper bounds of the f_i 's, hence if there exists $v \in \pi_1(\text{tr}(f))$ such that $v \leq \bar{x}$, then $([M_1]g\bar{x}, \dots, [M_m]g\bar{x}) \geq \alpha(v)$.

For the opposite direction, let us suppose that $f(\bar{x}) = \perp$, and hence for all $v \in \pi_1(\text{tr}(f))$, $\bar{x} \not\geq v$. By definition of the \hat{f}_i 's we know that for all $w \in \alpha(V_{H_f})$, $([M_1]g\bar{x}, \dots, [M_m]g\bar{x}) \leq w$, since, under the hypothesis $f(\bar{x}) = \perp$, we have that for all $1 \leq j \leq m$, for all $b \in \{ \text{tt}, \text{ff} \}$ $[M_j]g\bar{x} = b$ implies $\hat{f}_j = \lambda\bar{x}$. b implies for all $w \in \alpha(V_{H_f})$, $w^j = b$.

Since V_{H_f} is a hyperarc, we know that $\#\alpha(V_{H_f}) \geq 2$, and by minimality of the elements of $\pi_1(\text{tr}(g))$ we conclude that for all $w \in \pi_1(\text{tr}(g))$ $([M_1]g\bar{x}, \dots, [M_m]g\bar{x}) \not\geq w$, and hence $g([M_1]g\bar{x}, \dots, [M_m]g\bar{x}) = \perp$.

We can now conclude the proof, again by case reasoning on the structure of H_f :

- V_{H_f} is a monochromatic hyperarc (w.l.o.g. assume that all vertices are white). Then it is easy to check that the term

$$M = \lambda g \lambda \bar{x}. \text{ if } g(M_1 g \bar{x}) \dots (M_m g \bar{x}) \text{ then tt else tt}$$

g -defines f .

- V_{H_f} is not monochromatic: we first note that in this case

$$\forall x, y \in V_{H_f} \quad C(x) = C(y) \Leftrightarrow C(\alpha(x)) = C(\alpha(y))$$

i.e. α acts as the identity or the “negation” on colours (the “ \Leftarrow ” direction follows directly from the definition of weak morphism; as for “ \Rightarrow ”, remark that, since V_{H_f} is a polychromatic hyperarc, if $C(x) = C(y)$, then there exists $z \in V_{H_f}$ such that $C(z) \neq C(x)$. Since it must be $C(\alpha(z)) \neq C(\alpha(x))$ and $C(\alpha(z)) \neq C(\alpha(y))$, the result follows). We define then

$$M = \lambda g \lambda \bar{x}. \epsilon(g(M_1 g \bar{x}) \dots (M_m g \bar{x}))$$

where ϵ is the boolean identity or the boolean negation according to how α acts on colours. Then again it is easily checked that M g -defines f . ■

In the following example, we “run” the proof of the theorem in order to construct a PCF-term which defines f_3 relatively to g_3 , these functions being defined in the example 1.

Example 26:

Since $H_{f_3} = H_{g_3} =$

$$(\{1, 2, 3\}, \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}, C(1) = C(2) = C(3) = \text{white})$$

we can choose $id : H_{f_3} \rightarrow H_{g_3}$ as morphism. The corresponding transformation of $\pi_1(\text{tr}(f_3))$ onto $\pi_1(\text{tr}(g_3))$ is then:

$$(\text{tt}, \perp, \perp) \longrightarrow (\perp, \text{tt}, \text{tt})$$

$$(\perp, \text{tt}, \perp) \longrightarrow (\text{tt}, \perp, \text{tt})$$

$$(\perp, \perp, \text{tt}) \longrightarrow (\text{tt}, \text{tt}, \perp)$$

The slice functions are hence defined by:

$$\text{tr}(f'_1) = \text{tr}(\hat{f}'_1) = \{((\perp, \text{tt}, \perp), \text{tt}), ((\perp, \perp, \text{tt}), \text{tt})\}$$

$$\text{tr}(f'_2) = \text{tr}(\hat{f}'_2) = \{((\text{tt}, \perp, \perp), \text{tt}), ((\perp, \perp, \text{tt}), \text{tt})\}$$

$$\text{tr}(f'_3) = \text{tr}(\hat{f}'_3) = \{((\text{tt}, \perp, \perp), \text{tt}), ((\perp, \text{tt}, \perp), \text{tt})\}$$

The f'_i 's being non-sequential, we have to re-run our proof in order to define them relatively to g_3 . Let us consider f''_1 . The following picture represents a morphism $\alpha' : H_{f''_1} \rightarrow H_{g_3}$:

$$(\perp, \text{tt}, \text{tt})$$

$$(\perp, \text{tt}, \perp) \longrightarrow (\text{tt}, \perp, \text{tt})$$

$$(\perp, \perp, \text{tt}) \longrightarrow (\text{tt}, \text{tt}, \perp)$$

The corresponding slice functions are

$$f''_1 = f'_1 \neq \hat{f}''_1 = \lambda \bar{x} \text{tt}$$

$$\text{tr}(f''_2) = \text{tr}(\hat{f}''_2) = \{((\perp, \perp, \text{tt}), \text{tt})\}$$

$$\text{tr}(f''_3) = \text{tr}(\hat{f}''_3) = \{((\perp, \text{tt}, \perp), \text{tt})\}$$

Now the f_i'' s are trivially g_3 -definable (their traces are singletons). The corresponding terms are $M_1 = \lambda h \lambda \bar{x} \text{ tt}$, $M_2 = \lambda h \lambda \bar{x}$ if x_3 then tt else \perp , $M_3 = \lambda h \lambda \bar{x}$ if x_2 then tt else \perp .

The term M g_3 -defining f_1' is hence:

$$M = \lambda h \lambda \bar{x} \text{ if } h (M_1 h \bar{x}) (M_2 h \bar{x}) (M_3 h \bar{x}) \text{ then } \text{tt} \text{ else } \text{tt}$$

By eliminating redundant conditional statements (and with some abuse of notation) we obtain the following definition of f_1' :

$$f_1' = \lambda \bar{x} g_3(\text{tt}, x_3, x_2)$$

similar constructions allow us to obtain the terms g_3 -defining f_2' and f_3' , and finally we get (again with some simplifications)

$$f_3 = \lambda x_1 x_2 x_3 g_3(g_3(\text{tt}, x_3, x_2), g_3(x_3, \text{tt}, x_1), g_3(x_2, x_1, \text{tt}))$$

We can observe that this construction leads to a term which is more complex than the one showed in example 1.

■

We can of course remark that:

Corollary 27 *If H_f and H_g are strongly (or weakly) isomorphic, then $[f] = [g]$.*

This corollary answers to a question asked in the introduction: functions having the same hypergraph are equiparallel.

Another remark concerns subsequential functions: if H_f has monochromatic hyperarcs then any function $\alpha : V_{H_f} \rightarrow V_{H_g}$ which preserves hyperarcs is a weak morphism. Hence:

Corollary 28 *Let \mathcal{F} be the forgetful functor from colored hypergraph to hypergraph, and let $\alpha : \mathcal{F}(H_f) \rightarrow \mathcal{F}(H_g)$ be a hypergraph morphism. If f is subsequential then $f \leq_{\text{par}} g$.*

6 Conclusion

We have seen several properties relating the poset of degrees and a category of hypergraphs: Concerning the objects of this category we have shown how one can naturally characterize basic properties of boolean functions in term of hypergraphs. Concerning the arrows we have shown that hypergraph morphisms reflect \leq_{par} relations. Moreover, when a morphism $\alpha : H_f \rightarrow H_g$ does exist, we can extract from the proof of theorem 25 a PCF-term which defines f relatively to g .

One natural question at this point is whether hypergraph morphisms preserve \leq_{par} relations, i.e. whether whenever $f \leq_{\text{par}} g$, $\mathcal{WH}(H_f, H_g)$ is non-empty. The answer is no; for example, consider:

Example 29: Let $f_3 : \mathcal{B}^3 \rightarrow \mathcal{B}$ be the function defined in example 1. Its hypergraph is:

$$H_{f_3} = (\{1, 2, 3\}, \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}, C(1) = C(2) = C(3) = \text{white})$$

It is easy to see that there exists no (even weak) morphism $m : H_{f_3} \rightarrow H_{\text{por}}$. Nevertheless $f_3 \leq_{\text{par}} \text{por}$, since for instance

$$f_3 = \llbracket M \rrbracket_{\text{por}}$$

where

$$M = \lambda f \lambda x_1 x_2 x_3. \text{ if } f(f(x_1, x_2))x_3 \text{ then tt else } \perp$$

■

Although the notions of hypergraph morphism presented here are too weak in order to get a completeness result we do believe that hypergraph representation does retain enough information on functions in order to achieve such completeness. The price to pay seems to be the use of more involved notions than (weak or strong) hypergraphs morphisms.

References

- [1] S. Abramsky, R. Jagadeesan, P. Malacaria. *Full abstraction for PCF (Extended Abstract)*. Proc. of TACS 94, Lecture Notes in Computer Science 789, Springer, 1994.
- [2] R. Amadio, P.-L. Curien. *Selected Domains*. To appear.

- [3] G. Berry. *Stable models of typed lambda-calculi*. Proc. 5th Int. Coll. on Automata, Languages and Programming, Lecture Notes in Computer Science 62, Springer, 1978.
- [4] G. Berry. *Modèles complètement adéquats et stables des lambda-calculs typés*. Thèse de Doctorat d'Etat, Université Paris 7, 1979.
- [5] A. Bucciarelli, T. Ehrhard. *Sequentiality in an extensional framework*. Information and Computation, Volume 110, Number 2, 1994.
- [6] A. Bucciarelli. *Degrees of Parallelism in the Continuous Type Hierarchy*. To appear in Theoretical Computer Science.
- [7] L. Colson, T. Ehrhard. *On strong stability and higher-order sequentiality*. Proc. 9th Symp. on Logic in Computer Science, IEEE press, 1994.
- [8] T. Ehrhard. *Parallel and Serial Hypercoherences*. Manuscript 1995.
- [9] J-Y. Girard. *The system F of variable types fifteen years later*. Theoretical Computer Science 45, 159-192, 1986.
- [10] J.M.E. Hyland, L. Ong. *On full abstraction for PCF: I, II and III* (preliminary version, september 1995).
- [11] G. Kahn and G. Plotkin. *Domaines Concrets*. Rapport IRIA-LABORIA 336, 1978, republished in the special issue of Theoretical Computer Science dedicated to Professor C. Böhm's 70th birthday, 1993.
- [12] R. Loader. *Finitary PCF is not decidable*. Unpublished notes, available at <http://info.ox.ac.uk/loader>, 1996.
- [13] R. Milner. *Models of LCF*. Computer Science Departement Memo. AIM-186/CS 332, Stanford University, 1973.
- [14] P. O'Hearn, J. Riecke. *Kripke Logical Relations and PCF*. To appear in Information and Computation.
- [15] G. Plotkin. *LCF considered as a programming language*. Theoretical Computer Science 5, 223-256, 1977.
- [16] V. Y. Sazonov. *Degrees of Parallelism in Computations*. Proc. Conference on Mathematical Foundations of Computer Science, Lecture Notes in Computer Science 45, Springer, 1976.
- [17] D. Scott. *A type theoretic alternative to OWHY, CUCH, ISWIM*. Theoretical Computer Science 121, 411-440, 1993 (manuscript circulating since 1969).
- [18] K. Sieber. *Reasoning about Sequential Functions via Logical Relations*. Proc. LMS Symposium on Applications of Categories in Computer Science, M. Fourman, P. Johnstone, A. Pitts eds, LMS Lecture Note Series 177, Cambridge University Press, 1992.
- [19] A. Stoughton. *Mechanizing Logical Relations*. Proc. Ninth International Conference on Mathematical Foundations of Programming Semantics, Lecture Notes in Computer Science 802, Springer, 1994.

- [20] A. Stoughton. *Interdefinability of parallel operations in PCF* Theoretical Computer Science 79, 357-358, 1991.
- [21] M.B. Trakhtenbrot. *On Representation of Sequential and Parallel Functions*. In Proc. Fourth Symp. on Mathematical Foundations of Computer Science, Lecture Notes in Computer Science 32, Springer, 1975.
- [22] J. Vuillemin. *Proof Techniques for Recursive Programs*. Ph.D. Thesis, Stanford University, 1973.