

The effect of ethnicity and genetic ancestry on the epidemiology, clinical features and outcome of Systemic Lupus Erythematosus

Myles J. Lewis,^{1*} Ali S. Jawad¹

¹Centre for Experimental Medicine and Rheumatology, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, EC1M 6BQ, UK.

*Corresponding author

Email: myles.lewis@qmul.ac.uk

Tel. +44 20 7882 3305

Fax +44 20 7882 8250

Keywords: systemic lupus erythematosus, lupus nephritis, ethnicity, ethnic groups, ancestry, genetics, epidemiology, incidence, prevalence, autoantibodies

Conflict of interest statement: both authors declare that they have no conflicts of interest.

Abstract

In this in-depth review, we examine the worldwide epidemiology of Systemic lupus erythematosus (SLE) and summarise current knowledge on the influence of race/ethnicity on clinical manifestations, disease activity, damage accumulation and outcome in SLE.

Susceptibility to SLE has a strong genetic component and trans-ancestral genetic studies have revealed a substantial commonality of shared genetic risk variants across different genetic ancestries which predispose to the development of SLE. The highest increased risk of developing SLE is observed in black individuals (incidence 5- to 9-fold increased, prevalence 2- to 3-fold increased), with an increased risk also observed in South Asians, East Asians and other non-white groups, compared to white individuals. Black, East Asian, South Asian and Hispanic SLE individuals tend to develop more severe disease with a greater number of manifestations and accumulate damage from lupus more rapidly. Increased genetic risk burden in these populations, associated with increased autoantibody reactivity in non-white SLE, may explain the more severe lupus phenotype. Even after taking into account socio-economic factors, race/ethnicity remains a key determinant of poor outcome such as end-stage renal failure and mortality in SLE. Community measures to expedite diagnosis through increased awareness in at-risk racial/ethnic populations and ethnically personalised treatment algorithms may help in future to improve long term outcomes in SLE.

Introduction

Over recent years our understanding of Systemic Lupus Erythematosus (SLE) has advanced at the level of its epidemiology, genetic susceptibility and depth of understanding of molecular mechanisms underlying its pathogenesis. These advances have been achieved through national and international collaboration between SLE researchers and clinicians, in conjunction with the harnessing of post-genome era technology. Early studies on the influence of ethnicity on the incidence and prevalence of SLE are thoroughly described in a key review by Danchenko et al [1], and a subsequent analysis of SLE epidemiological studies including the effect of ethnicity on clinical manifestations was last comprehensively conducted by Borchers et al [2]. This current review will concentrate on highlighting and contrasting additional information from subsequent studies.

Worldwide epidemiology of SLE

Reliable studies of incidence and prevalence of SLE remain few and far between, due to problems of cohorts being incomplete or drawn from populations which are not closed or for which the boundaries are unclear. In many countries a lack of high quality census data has prohibited many cohort studies from reporting accurate incidence and prevalence figures, even where consistent patient registries have been built up over a number of years. In countries with more extensive public health systems, accurate SLE patient registries have in the past provided important information on incidence and prevalence including studies from Denmark, Finland, Norway, Sweden and UK [3-7]. More recent studies have benefited from improved definitions of SLE diagnosis based first on the 1997 American College of Rheumatology (ACR) classification criteria for SLE, and the more recent 2012 Systemic Lupus Collaborating Clinics (SLICC) revision of the ACR criteria. However few studies have employed capture-recapture technique to improve validity. Two recent large and comprehensive studies from Michigan and Georgia cohorts in the US [8, 9], showed remarkable consistency and reported the incidence of SLE as 5.5 and 5.6 per 100,000 respectively and the prevalence of SLE as 72.8 and 73.0 per 100,000, based on data from

2002-2004. Comparisons of studies across the world shows a wide range of prevalence from 25-28 per 100,000 in Northern Ireland, Finland and Denmark [3, 7, 10] to 149 in Alaska [11] and 159 in Puerto Rico [12]. These studies are summarised in Figures 1 and 2. Direct comparisons between countries are problematic, because of a clear increase in both the incidence and prevalence of SLE over time, as shown in Fig. 3. A recent large UK study reported a relatively stable overall incidence of 4.9 per 100,000 per year over the time period 1999-2012, while the prevalence of SLE rose steadily from 65.0 in 1999 to 97.0 per 100,000 in 2012 [13]. Comparison of studies across the world from the 1970s through to present day (Fig. 2) illustrate that the prevalence of SLE has risen over time with studies in the 1970s reaching a maximum prevalence of 40 per 100,000, while several large studies performed since 2000 have reported point prevalence of 100 per 100,000 or more [12-15]. The incidence of SLE has not risen over time to the same extent, but there are still a significant number of recent studies from UK, USA and Taiwan with incidence rates ranging from 4.8 to 7.2 per 100,000, a level not seen in early studies. This rise in prevalence is a feature of a number of studies and presumably reflects improved diagnosis over time as well as improved survival rates, alongside the lifelong chronicity of the illness. Although the SLICC modification of the ACR classification criteria for SLE has not yet been used by any of the currently reported epidemiological studies, it is likely to increase the diagnostic rate for SLE further [16].

Defining race, ethnicity and ancestry in clinical studies

Although race and ethnicity are terms that are commonly used in medical studies, their definition is complex. The concept “race” originally arose to group individuals by biological differences such as skin colour, with associated social and culture differences. However the term ran into difficulties when describing increasingly mixed populations such as in the USA. A later concept of “ethnicity” emphasised cultural, social and religious qualities, as opposed to genetic ancestry. However both race and ethnicity are fluid, and perceived racial or ethnic groups change over time. Genetic studies often use the term “ancestry”, by which individuals are categorised according to the origins of their ancestors. Ancestry is usually defined

geographically (e.g. East Asian, sub-Saharan African, northern European), and self-reported ancestry tends to correlate reasonably well with genetically-determined ancestry. However ancestry is a multi-dimensional continuum with endless possible sub-refinements (e.g. Norwegian versus Scandinavian) [17]. In practice all three terms of race, ethnicity and ancestry are interlinked. US government guidelines for reporting race and ethnicity include 5 minimum categories of race (American Indian/ Alaska Native (AIAN), Asian, Black/African American, Hawaiian/Pacific Islander and white), while Hispanic/Latino individuals are categorised separately under ethnicity, referring to people of Mexican, South or Central American or other Spanish culture, regardless of race [18]. “Amerindian” is a broad ancestral term which refers to indigenous people of the Americas, including Central or South America. Genetic analyses reveal that Hispanic individuals display ancestral genetic diversity, with Hispanics from Texas showing a significant proportion of Amerindian ancestral genes, whereas African and white European ancestral genes are more common in Hispanics from Puerto Rico [19, 20]. The Mestizo, studied under the ethnic grouping “Hispanic”, are individuals of combined European and Amerindian (usually indigenous Mexican/central American) ancestry. “Asian” is an ambiguous term since in the UK it usually refers to individuals from India, Pakistan, Bangladesh or neighbouring regions, while in the USA it usually refers to individuals of Chinese, Korean or Japanese origin. Hence in this article we distinguish between South Asian and East Asian, although these terms are generally not used in the source studies.

Effect of race and ethnicity on incidence and prevalence of SLE

A consistent finding across epidemiological studies is that SLE is significantly more common in black individuals and specific races and ethnicities compared to white individuals. This finding has been repeatedly confirmed in numerous studies across the USA through to modern day [8, 9, 21-23]. Similar findings were reported in UK studies based in Birmingham and Nottingham [4, 24], which showed 5- to 9-fold increased incidence and 5- to 10-fold increase in prevalence of SLE in Afro-Caribbeans, as well as a 1.2- to 6-fold increase in

incidence and 2- to 2.4-fold increase in prevalence in South Asian individuals compared to white individuals. Further attempts to define ethnic groups at risk of SLE are complicated by lack of racial and genetic homogeneity for certain ethnicities. SLE in Hispanic individuals has been studied extensively in the LUMINA (Lupus in Minorities: Nature vs. nurture) cohort studies [20]. In Canada, a 2-fold increase in incidence and prevalence of SLE was observed in native American Indian individuals compared to whites [25], and a later study observed that American Indian and Alaska Native individuals demonstrated an incidence of 7.4 per 100,000 and prevalence of 178 per 100,000 of SLE, which are some of the highest rates recorded and comparable to African Americans [11]. Although substantial information is available about SLE in African ancestry individuals living in the Americas and Europe, there are currently no accurate studies of the incidence or prevalence of SLE in Africa, the Middle East or South Asia. A recent UK study, which is one of few to subdivide South Asian ethnicity into subgroups, found that Indian ethnicity is associated with higher incidence (9.9 per 100,000) and prevalence of SLE (193 per 100,000), although not as high as in Afro-Caribbeans (incidence 31.5 and prevalence 518 per 100,000). The incidence (10.0 per 100,000) of SLE was higher in UK Pakistanis, although the prevalence was not increased (143 per 100,000) [13]. The incidence and prevalence of SLE in UK Bangladeshi individuals was apparently not increased compared to whites. However a problem with this study is the high percentage of individuals with 'unknown' or 'unclassified' ethnicity. Subgroup analysis of the Michigan Lupus registry suggests that the incidence of SLE in individuals of Arab or Chaldean origin was 2.1-fold increased compared to white individuals [26].

Early studies anecdotally suggested a low prevalence of SLE in Africa. Although there remains a lack of high quality epidemiology studies from African countries, data on migrants shows that the incidence and prevalence of SLE is high in Africans, Caribbeans and South Asians who have migrated to the UK [4, 27]. Recent evidence suggests that SLE and lupus nephritis are a significant health burden in Africa [28, 29]. An interesting study showed that rates of SLE autoantibody positivity were similar in a cohort of healthy Gullah

individuals (African Americans from the Sea Islands of South Carolina of low genetic admixture) compared to a genetically similar population of women from Sierra Leone [30].

No studies directly comparing diverse populations have reported statistically significant variance from the gender bias towards female individuals being affected by SLE. In studies across multiple populations the percentage of females affected ranged from 88.3 to 96.2%, giving a female:male gender ratio centred around 9:1, ranging from 7.5:1 in Curaçao to 23:1 in Oman and 25:1 in Hispanic individuals in a USA cohort [31-33].

Genetics, ancestry and SLE

SLE is known to have a strong genetic link with a heritability of ~66%. The twin concordance rate for SLE is 24% in monozygotic twins compared to 2% in dizygotic twins [34]. SLE has a sibling recurrence risk ratio (λ_s) of 29 compared to 5.3 in rheumatoid arthritis (RA) [35]. Since the advent of genome wide association studies (GWAS), at least 52 genetic loci with strong evidence of association with susceptibility to SLE have been confirmed [36-41]. Although cohort size is a major factor, which has increased the power of these studies to find genetic associations, the veracity of these large scale studies is also underpinned by radical improvements in computer science in data storage, quality control, statistical and analytical techniques. The region of the genome with by far the strongest association with susceptibility to SLE is the major histocompatibility complex (MHC), located on chromosome 6. Prior to GWAS scale studies the 8.1 ancestral haplotype has long been known to be associated with endocrine, gastrointestinal, haematologic/immunologic autoimmune diseases and sarcoidosis through to multiple connective tissue diseases including SLE. This MHC haplotype extends to include *HLA-A1*01:01*, *B*08:01*, *C4AQ0*, *C4B1*, *DRB1*03:01*, *DQA1*05:01*, *DQB1*02:01* (*A1::DQ2* in older nomenclature), encompassing two significant class III genetic variants, complement *C4A* null as well as the *TNF-308A* allele. Major confounding factors affecting MHC genetic association studies are long-range linkage disequilibrium (LD) and duplications of regions leading to copy number variation (CNV) and pseudogenes. Only the largest most recent genetic studies using techniques including trans-

ancestral mapping have sufficient power to dissect which genes within the MHC are responsible for disease susceptibility rather than finding associations with the whole MHC block. The highest HLA association with SLE has been observed for *HLA-DRB1*03:01*, with a weaker effect observed for *HLA-DRB1*15:01* [42]. However although this study including multiple cohorts from different countries including the USA, the individuals included were predominantly of European ancestry. Trans-ancestral mapping of UK, Spanish and Filipino SLE patients showed differential MHC signals according to ancestry [43]. Independent association of *HLA-DRB1*08:01* was seen in Spanish SLE but not in other groups, in addition to known association with *HLA-DRB1*03:01* and *HLA-DRB1*15:01*. SNPs with the best odds ratio in Spanish SLE clustered around the DNA repair gene *MSH5*. The top SNP in Filipino SLE, rs9271366, located between *HLA-DRB1* and *HLA-DQA1*, may represent association with *HLA-DRB1*15:02*. The risk haplotype with the top 3 independent MHC SNPs was also more common in Filipino SLE, consistent with the increased prevalence of SLE in East Asian population. The association of *HLA-DRB1*03:01* with SLE is stronger in anti-Ro and anti-La antibody positive SLE, suggesting that this HLA haplotype promotes break of tolerance to Ro and La autoantigens, independently of its association with SLE [44]. Interestingly a study of black women with SLE could not confirm association of SLE with *HLA-DRB1*03:01* and *HLA-DRB1*15:01* in this population, although of the four independently associated SNPs, two are located in MHC class II region consistent with other studies in Europeans and East Asians [38, 42]. A relatively small study suggested that *HLA-DRB1*15:03* and *HLA-DRB1*08* were more frequent in African American and Hispanic SLE individuals respectively [45]. These dissimilarities are in keeping with the heterogeneity of the extended MHC locus and will require larger studies to fully dissect differences between ancestries in terms of MHC region mediated susceptibility to SLE. By comparison, in rheumatoid arthritis five key amino acid residues in HLA-DR β 1, HLA-B and HLA-DP β 1 are able to explain almost completely the genetic susceptibility to RA conferred by the HLA region [46]. Trans-ancestral studies in RA show that these amino acids confer shared effects in Asian individuals [47]. Similar in-depth trans-ancestral analyses of the MHC including

HLA amino acid variation in SLE are currently lacking. Genes encoding complement C4 and TNF-alpha are also located within the MHC region. A study of northern European (UK) and southern European (Spanish) SLE populations showed that C4 CNV was not an independent risk factor for SLE susceptibility in either population [48]. A meta-analysis suggested that the *TNF* promoter -308A/G polymorphism may increase risk of SLE in European populations, but not in Asian populations [49].

Outside of the MHC, recent GWAS studies have superseded older candidate gene studies and have shown over 40 genes associated with SLE in European populations reaching genome-wide significance ($p < 5 \times 10^{-8}$) [36]. Comparison with GWAS in Han Chinese individuals initially suggested some differences in gene associations [38], with associations found in *RASGRP3*, *ETS1* and *WDFY4/LRRC18* which had not been previously identified in Europeans and association with *IKZF1* and *SLC15A4* for which early European data was only suggestive of association. These differences are largely due to altered risk variant frequency with apparently East Asian specific variants being more common in East Asians than in Europeans and vice versa. However as GWA studies become larger and larger, some risk variants which are rarer in certain populations are likely to be revealed as part of a commonality of genetic susceptibility across multiple ancestries, and this has proven to be the case for all the above genes except *RASGRP3*, which remains the only gene associated with Chinese SLE not to be replicated in Europeans at time of writing [36]. A fresh meta-analysis of European and Chinese GWAS data [41] confirmed substantial commonality in shared risk variants between the two populations. This is similar to trans-ancestral GWAS in RA which showed >80% of the heritability attributed to 101 RA risk loci was shared between Europeans and Asians [50]. Overall, risk variant frequencies were higher in Chinese individuals than in Europeans in keeping with the higher prevalence of SLE in Chinese individuals, suggesting a greater SLE genetic risk burden in East Asia. GWAS in Amerindian SLE identified association with *IRF5*, *ITGAM*, *STAT4*, *TNIP1*, *NCF2* and *IRAK1*, which were all previously identified in European SLE GWAS [36]. However it was striking that the association with *IRF5* (interferon response factor-5) was stronger than the HLA association in

this cohort. This study also identified rs4917385 in proximity of *USMG5* with eQTL evidence that this SNP affects *USMG5* expression as an Amerindian specific susceptibility locus. A candidate gene study in African American individuals suggested *MECP2*, *MBL2* and *PXK* as European only-associated SLE susceptibility genes [51]. Results of an ongoing GWAS in African ancestry SLE are not yet available, but will further inform our understanding of ancestral differences in genetic susceptibility to SLE once available. Trans-ancestral fine mapping studies of individuals genes are rare but informative. They illustrate that although the most strongly associated SNPs vary with ancestry, common risk haplotypes can be identified across different ancestries for example for *TNFSF4* (which encodes OX40L, an activatory ligand for OX40 receptor on T lymphocytes) [52] and the *IL2/IL21* locus [53]. Functional genetic studies have also begun to elucidate the immunological mechanisms of SLE susceptibility variants in ancestrally diverse populations [54, 55].

A critical component of modern genetic studies is the use of ancestry-informative markers (AIMs), which measure genetic admixture to more accurately account for the effect of ancestry when studying genetic susceptibility to SLE. AIMs are necessary to exclude genetic variants whose apparent association with SLE is merely due to a confounding association with ancestral grouping. AIMs can be also used to probe for interaction between specific genes and ancestry in development of SLE in different ancestral groups. Using this method, Amerindian admixture was modestly correlated with increased numbers of risk alleles in a study of 16 confirmed genetic susceptibility loci [56].

Racial differences in gene expression have been reported, with high expression of *TLR9* and DNA methyltransferase *DNMT3A* observed in African American SLE compared to other ethnicities, associated with higher serum levels of TNF and IL-6 [57, 58].

Effect of ethnicity on clinical features and disease activity

Many studies have examined whether ethnicity affects clinical phenotype in SLE. However these studies are complicated by the exceptional heterogeneity of disease manifestations in SLE: both diversity of organ involvement and heterogeneity between individuals. Differences

in reporting methods for clinical features as well as accurate recording of cumulative occurrence of manifestations add to the complexity of comparing studies. Previous reviews have compared clinical features across studies in different countries [2, 20]. Rates of ACR criteria manifestations were uniformly lower in Cervera and colleagues' study of 1000 predominantly white Europeans [59] compared to the GLADEL study by Pons-Estel et al and the LUMINA cohort by Alarcon et al [31]. Rates of clinical manifestations in a study of Hong Kong Chinese fell in between these studies [60]. These differences are perhaps more likely to be due to methodology than actual differences between ethnicities. Studies which specifically aimed to compare different races/ethnicities over the same time period and similar geographical areas are more likely to be accurate and representative. Thus it is generally accepted that individuals of African or Caribbean ancestry are more frequently affected by discoid rash (20-34%) compared to white Europeans (11-12%) [31, 61, 62], while photosensitivity may be more common in Europeans (81%) and individuals from Puerto Rico (90%) who have some European admixture compared to African Americans or Hispanics in Texas (both 56%) [20, 31, 63]. In UK studies, Afro-Caribbean individuals with SLE tended to develop renal disease earlier in the course of their illness and develop renal disease more frequently [64, 65]. Similar findings are reported in non-Europeans in France and in African Americans in the USA, Canada and Latin America [25, 61, 63, 66-68]. High levels of renal involvement are found in South Africa [69], Martinique [70], in Arab populations in Tunisia and Saudi Arabia [71, 72], and Asian populations including Chinese, Malaysians, Filipinos and Indians [60, 73-76]. A systematic review of glomerulonephritis in Africa identified lupus nephritis as one of the most common causes of secondary glomerular disease [28].

Individuals of African ancestry and Hispanics from Texas show more severe renal involvement with greater rates of progression to end-stage renal disease [66, 67], although Hispanic SLE patients from Texas developed more severe disease than those in Puerto Rico [77]. Renal disease is also more frequent in mestizo populations (mainly from Guatemala, Mexico and Peru) with SLE [61]. Rates of neuropsychiatric SLE (NPSLE) varied widely from cohort to cohort in older studies, with a range of 14% to 91%, rendering direct

comparisons between studies problematic. However the SLICC inception cohort has reported several studies in this area looking at individual NPSLE manifestations. In this cohort, seizure occurrence was 2.0-fold increased in African and 1.6-fold increased in Hispanic SLE compared to white individuals using multivariate analysis, with a tendency for seizures to be lower in frequency in Asian SLE individuals [78]. Mood disorders were lower in Asian SLE individuals, confirmed in South Korean patients [79], however there was no effect of ethnicity/race on the prevalence of lupus headache [80]. However the SLICC cohort made no distinction between South and East Asia. Ethnicity/race has not been shown to predispose to thrombotic events [81, 82].

Some studies have suggested that non-European SLE individuals show a higher frequency of positivity for specific autoantibodies such as anti-Sm [83]. Using a bead-based autoantibody assay African-American and Hispanic SLE patients showed positivity for a greater number of autoantibodies concomitantly: 53% of African American and 34% of Hispanic SLE individuals demonstrated 4 or more autoantibody specificities compared to 19% of European SLE individuals [84]. Anti-chromatin, anti-Sm, anti-RNP, anti-dsDNA positivity was higher in African-American SLE and to a lesser extent Hispanic SLE compared to European SLE in this study. Other studies have confirmed the higher rate of anti-Sm and anti-RNP positivity in non-European ancestry SLE [20, 85, 86], but not all studies have confirmed this [61]. On the other hand anti-Ro and anti-La positivity tends to be similar across ethnicities.

Impact of ethnicity on disease activity, damage accumulation and mortality

Consistent with these trends that ethnicity/race influences rates of skin, renal, neuropsychiatric and immunologic manifestations, ethnicity impacts disease activity as a whole, with non-white individuals showing higher SLEDAI scores compared to whites both at baseline and over time [45, 87, 88]. Beyond the observation that clinical manifestations may be both more common and of greater severity in black and Hispanic individuals and to a varying degree Asian individuals, ethnicity has a knock-on effect on damage accumulation

and outcome over time. Doubling of creatinine and progression to end-stage renal failure (ESRF) is more common in African American and Hispanic individuals with lupus nephritis [89], and presentation with end-stage renal disease or rapid progression to ESRF in newly diagnosed SLE patients is significantly more common in African Americans [90]. Large studies observing more than 12000 lupus nephritis patients in the US ESRF registry have shown that African American SLE patients who reach end-stage renal failure have higher mortality rates than non-African populations, although one study suggested this effect may be related to median income [91] and a second study showed higher levels of cardiovascular events in the African American group [92]. Current data suggests that different ethnicities show similar allograft failure rates following renal transplantation for lupus nephritis [93]. In the SLICC cohort, African American SLE individuals showed the highest rates of progression of SLICC/ACR damage index (SDI), followed by Hispanic individuals, with the rate of progression in Asian SLE lower than for white individuals [94]. Other key factors predictive of damage accrual are age, disease activity including the number of flares and corticosteroid use [94, 95]. Black and Hispanic SLE individuals also respond more slowly to treatment and show slower falls in disease activity over time compared to white patients [96]. Increased rates of damage accumulation in African ancestry SLE is of major concern, because it has been consistently shown that damage score at baseline and damage accrual rates are a major predictor of mortality in SLE and consistent with this mortality is higher in African individuals with SLE [97-99]. A meta-analysis of mortality rates in SLE showed comparable rates between Asian studies in China and Taiwan compared to Europe and North America [100].

Interestingly there is some evidence of differential response to therapy according to ethnic group, although few studies have examined this issue systematically. African and Hispanic SLE patients showed greater response to rituximab compared to other groups [101]. Black and mixed-race SLE individuals showed poorer response to intravenous cyclophosphamide [102, 103]. Future randomised controlled trials (RCT) which include genotype data on study participants will be better placed to dissect these issues by measuring

genetic admixture instead of self-reported race/ ethnicity, especially in the case of mixed-race individuals who may be incorrectly categorised.

Socio-economic factors

In many countries where private health care predominates, including the US, poorer access to health care is self-evidently linked to worse outcome and morbidity. Thus a number of studies have attempted to distinguish the effect of ethnicity from socio-economic factors such as employment status, annual income, education level and social support network. General consensus from these studies is that although socio-economic factors play a definite role in worse outcome especially in SLE individuals with renal disease [91], ethnicity still acts as an independent risk factor for increased prevalence and severity of disease and worse outcome [45, 65, 67, 104]. This effect of ethnicity above and beyond socio-economic factors also remains true in the UK, where health care is primarily publically funded [105]. This evidence supports the notion that genetic-determined ancestry and possibly environmental factors related to ethnicity are involved in biological processes which increase SLE prevalence and disease severity.

Comparison with other autoimmune diseases

These effects of race/ethnicity, which have been strongly observed for SLE as described above, are specific to SLE. Although the incidence and prevalence of RA shows variation in different countries, there is a lack of evidence for a significant effect of race/ ethnicity in RA [106]. Similarly the systemic vasculitides, which themselves can be difficult to classify, vary in incidence and prevalence in different countries [107], although specific types of vasculitis show well known regional distributions, for example Behçet's disease in Turkey and the Silk Road [108], and Takayasu's disease in the Far East. It is worth noting that multiple sclerosis (MS) shows inversion in terms of race/ethnicity in comparison to SLE, since MS is substantially more common in white individuals compared to black or South Asian populations, consistent with the geographical distribution of white populations [109, 110].

Conclusions

The overall message from an extensive body of literature is consistent: ancestry, race and ethnicity together have a major effect on the way in which SLE manifests. Black, Asian and Hispanic SLE individuals tend to develop more severe disease, a greater number of manifestations and accumulate damage from lupus more rapidly. Even after taking into account socio-economic factors, race/ethnicity remains a key determinant of poor outcome such as end-stage renal failure and mortality in SLE. Ongoing genetic studies suggest an increased genetic risk burden in these populations associated with increased autoantibody reactivity in non-European SLE, which may partly explain the more severe lupus phenotype. Subgroup analyses from RCTs have shown that different ethnic groups respond differentially to therapeutic interventions. Community measures to improve early diagnosis through increased awareness in at-risk racial/ethnic communities, together with more ethnically personalised treatment algorithms may be required in future to reduce the severity of SLE in high risk populations and thus improve long term outcome.

Funding statement

This work was supported by Arthritis Research UK. MJL is the recipient of an Arthritis Research UK Clinician Scientist Fellowship (19631).

References

- 1 Danchenko N, Satia JA, Anthony MS. Epidemiology of systemic lupus erythematosus: a comparison of worldwide disease burden. *Lupus* 2006;15(5):308-18.
- 2 Borchers AT, Naguwa SM, Shoenfeld Y, Gershwin ME. The geoepidemiology of systemic lupus erythematosus. *Autoimmunity reviews* 2010;9(5):A277-87.
- 3 Helve T. Prevalence and mortality rates of systemic lupus erythematosus and causes of death in SLE patients in Finland. *Scand. J. Rheumatol.* 1985;14(1):43-6.
- 4 Johnson AE, Gordon C, Palmer RG, Bacon PA. The prevalence and incidence of systemic lupus erythematosus in Birmingham, England. Relationship to ethnicity and country of birth. *Arthritis Rheum.* 1995;38(4):551-8.
- 5 Stahl-Hallengren C, Jonsen A, Nived O, Sturfelt G. Incidence studies of systemic lupus erythematosus in Southern Sweden: increasing age, decreasing frequency of renal manifestations and good prognosis. *J. Rheumatol.* 2000;27(3):685-91.
- 6 Eilertsen GO, Becker-Merok A, Nossent JC. The influence of the 1997 updated classification criteria for systemic lupus erythematosus: epidemiology, disease presentation, and patient management. *J. Rheumatol.* 2009;36(3):552-9.
- 7 Lastrup H, Voss A, Green A, Junker P. Occurrence of systemic lupus erythematosus in a Danish community: an 8-year prospective study. *Scand. J. Rheumatol.* 2009;38(2):128-32.
- 8 Lim SS, Bayakly AR, Helmick CG, Gordon C, Easley KA, Drenkard C. The incidence and prevalence of systemic lupus erythematosus, 2002-2004: The Georgia Lupus Registry. *Arthritis Rheumatol.* 2014;66(2):357-68.
- 9 Somers EC, Marder W, Cagnoli P, et al. Population-based incidence and prevalence of systemic lupus erythematosus: the Michigan Lupus Epidemiology and Surveillance program. *Arthritis Rheumatol.* 2014;66(2):369-78.
- 10 Gourley IS, Patterson CC, Bell AL. The prevalence of systemic lupus erythematosus in Northern Ireland. *Lupus* 1997;6(4):399-403.
- 11 Ferucci ED, Johnston JM, Gaddy JR, et al. Prevalence and incidence of systemic lupus erythematosus in a population-based registry of American Indian and Alaska Native people, 2007-2009. *Arthritis Rheumatol.* 2014;66(9):2494-502.

- 12 Molina MJ, Mayor AM, Franco AE, Morell CA, Lopez MA, Vila LM. Prevalence of systemic lupus erythematosus and associated comorbidities in Puerto Rico. *Journal of clinical rheumatology : practical reports on rheumatic & musculoskeletal diseases* 2007;13(4):202-4.
- 13 Rees F, Doherty M, Grainge M, Davenport G, Lanyon P, Zhang W. The incidence and prevalence of systemic lupus erythematosus in the UK, 1999-2012. *Ann. Rheum. Dis.* 2016;75(1):136-41.
- 14 Furst DE, Clarke AE, Fernandes AW, Bancroft T, Greth W, Iorga SR. Incidence and prevalence of adult systemic lupus erythematosus in a large US managed-care population. *Lupus* 2013;22(1):99-105.
- 15 Yeh KW, Yu CH, Chan PC, Horng JT, Huang JL. Burden of systemic lupus erythematosus in Taiwan: a population-based survey. *Rheumatol. Int.* 2013;33(7):1805-11.
- 16 Ines L, Silva C, Galindo M, et al. Classification of Systemic Lupus Erythematosus: Systemic Lupus International Collaborating Clinics Versus American College of Rheumatology Criteria. A Comparative Study of 2,055 Patients From a Real-Life, International Systemic Lupus Erythematosus Cohort. *Arthritis Care Res (Hoboken)* 2015;67(8):1180-5.
- 17 Race, Ethnicity and Genetics Working Group. The use of racial, ethnic, and ancestral categories in human genetics research. *Am J Hum Genet* 2005;77(4):519-32.
- 18 National Institute of Health. NIH policy and guidelines for the inclusion of women and minorities as subjects in clinical research – Amended, October, 2001. 2001. http://grants.nih.gov/grants/funding/women_min/guidelines_amended_10_2001.htm Accessed 18/7/2016.
- 19 Alarcon GS, Beasley TM, Roseman JM, et al. Ethnic disparities in health and disease: the need to account for ancestral admixture when estimating the genetic contribution to both (LUMINA XXVI). *Lupus* 2005;14(10):867-8.
- 20 Gonzalez LA, Toloza SM, McGwin G, Jr., Alarcon GS. Ethnicity in systemic lupus erythematosus (SLE): its influence on susceptibility and outcomes. *Lupus* 2013;22(12):1214-24.
- 21 Chakravarty EF, Bush TM, Manzi S, Clarke AE, Ward MM. Prevalence of adult systemic lupus erythematosus in California and Pennsylvania in 2000: estimates obtained using hospitalization data. *Arthritis Rheum.* 2007;56(6):2092-4.
- 22 Hochberg MC. The incidence of systemic lupus erythematosus in Baltimore, Maryland, 1970-1977. *Arthritis Rheum.* 1985;28(1):80-6.
- 23 McCarty DJ, Manzi S, Medsger TA, Jr., Ramsey-Goldman R, LaPorte RE, Kwok CK. Incidence of systemic lupus erythematosus. Race and gender differences. *Arthritis Rheum.* 1995;38(9):1260-70.
- 24 Hopkinson ND, Doherty M, Powell RJ. Clinical features and race-specific incidence/prevalence rates of systemic lupus erythematosus in a geographically complete cohort of patients. *Ann. Rheum. Dis.* 1994;53(10):675-80.

- 25 Peschken CA, Esdaile JM. Systemic lupus erythematosus in North American Indians: a population based study. *J. Rheumatol.* 2000;27(8):1884-91.
- 26 Housey M, DeGuire P, Lyon-Callo S, et al. Incidence and prevalence of systemic lupus erythematosus among Arab and Chaldean Americans in southeastern Michigan: the Michigan Lupus Epidemiology and Surveillance Program. *Am. J. Public Health* 2015;105(5):e74-9.
- 27 Molokhia M, McKeigue PM, Cuadrado M, Hughes G. Systemic lupus erythematosus in migrants from west Africa compared with Afro-Caribbean people in the UK. *Lancet* 2001;357(9266):1414-5.
- 28 Okpechi IG, Ameh OI, Bello AK, Ronco P, Swanepoel CR, Kengne AP. Epidemiology of Histologically Proven Glomerulonephritis in Africa: A Systematic Review and Meta-Analysis. *PloS one* 2016;11(3):e0152203.
- 29 Hodkinson B, Mapiye D, Jayne D, Kalla A, Tiffin N, Okpechi I. The African Lupus Genetics Network (ALUGEN) registry: standardized, prospective follow-up studies in African patients with systemic lupus erythematosus. *Lupus* 2016;25(3):325-30.
- 30 Gilkeson G, James J, Kamen D, et al. The United States to Africa lupus prevalence gradient revisited. *Lupus* 2011;20(10):1095-103.
- 31 Alarcon GS, McGwin G, Jr., Petri M, et al. Baseline characteristics of a multiethnic lupus cohort: PROFILE. *Lupus* 2002;11(2):95-101.
- 32 Al-Maini MH, El-Ageb EM, Al-Wahaibi SS, Al-Farsi Y, Richens ER. Demographic, autoimmune, and clinical profiles of patients with systemic lupus erythematosus in Oman. *Rheumatol. Int.* 2003;23(4):186-91.
- 33 Nossent JC. Systemic lupus erythematosus on the Caribbean island of Curacao: an epidemiological investigation. *Ann. Rheum. Dis.* 1992;51(11):1197-201.
- 34 Deapen D, Escalante A, Weinrib L, et al. A revised estimate of twin concordance in systemic lupus erythematosus. *Arthritis Rheum.* 1992;35(3):311-8.
- 35 Alarcon-Segovia D, Alarcon-Riquelme ME, Cardiel MH, et al. Familial aggregation of systemic lupus erythematosus, rheumatoid arthritis, and other autoimmune diseases in 1,177 lupus patients from the GLADEL cohort. *Arthritis Rheum.* 2005;52(4):1138-47.
- 36 Bentham J, Morris DL, Cunninghame Graham DS, et al. Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus. *Nat. Genet.* 2015;47(12):1457-64.
- 37 Gateva V, Sandling JK, Hom G, et al. A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus. *Nat. Genet.* 2009;41(11):1228-33.
- 38 Han JW, Zheng HF, Cui Y, et al. Genome-wide association study in a Chinese Han population identifies nine new susceptibility loci for systemic lupus erythematosus. *Nat. Genet.* 2009;41(11):1234-7.

- 39 Harley JB, Alarcon-Riquelme ME, Criswell LA, et al. Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXX, KIAA1542 and other loci. *Nat. Genet.* 2008;40(2):204-10.
- 40 Hom G, Graham RR, Modrek B, et al. Association of systemic lupus erythematosus with C8orf13-BLK and ITGAM-ITGAX. *N. Engl. J. Med.* 2008;358(9):900-9.
- 41 Morris DL, Sheng Y, Zhang Y, et al. Genome-wide association meta-analysis in Chinese and European individuals identifies ten new loci associated with systemic lupus erythematosus. *Nat. Genet.* 2016.
- 42 Rioux JD, Goyette P, Vyse TJ, et al. Mapping of multiple susceptibility variants within the MHC region for 7 immune-mediated diseases. *Proc. Natl. Acad. Sci. U. S. A.* 2009;106(44):18680-5.
- 43 Fernando MM, Freudenberg J, Lee A, et al. Transancestral mapping of the MHC region in systemic lupus erythematosus identifies new independent and interacting loci at MSH5, HLA-DPB1 and HLA-G. *Ann. Rheum. Dis.* 2012;71(5):777-84.
- 44 Morris DL, Fernando MM, Taylor KE, et al. MHC associations with clinical and autoantibody manifestations in European SLE. *Genes and immunity* 2014;15(4):210-7.
- 45 Reveille JD, Moulds JM, Ahn C, et al. Systemic lupus erythematosus in three ethnic groups: I. The effects of HLA class II, C4, and CR1 alleles, socioeconomic factors, and ethnicity at disease onset. LUMINA Study Group. *Lupus in minority populations, nature versus nurture. Arthritis Rheum.* 1998;41(7):1161-72.
- 46 Raychaudhuri S, Sandor C, Stahl EA, et al. Five amino acids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis. *Nat. Genet.* 2012;44(3):291-6.
- 47 Okada Y, Kim K, Han B, et al. Risk for ACPA-positive rheumatoid arthritis is driven by shared HLA amino acid polymorphisms in Asian and European populations. *Hum. Mol. Genet.* 2014;23(25):6916-26.
- 48 Boteva L, Morris DL, Cortes-Hernandez J, Martin J, Vyse TJ, Fernando MM. Genetically Determined Partial Complement C4 Deficiency States Are Not Independent Risk Factors for SLE in UK and Spanish Populations. *Am J Hum Genet* 2012;90(3):445-56.
- 49 Lee YH, Harley JB, Nath SK. Meta-analysis of TNF-alpha promoter -308 A/G polymorphism and SLE susceptibility. *Eur. J. Hum. Genet.* 2006;14(3):364-71.
- 50 Okada Y, Wu D, Trynka G, et al. Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature* 2014;506(7488):376-81.
- 51 Sanchez E, Comeau ME, Freedman BI, et al. Identification of novel genetic susceptibility loci in African American lupus patients in a candidate gene association study. *Arthritis Rheum.* 2011;63(11):3493-501.
- 52 Manku H, Langefeld CD, Guerra SG, et al. Trans-ancestral studies fine map the SLE-susceptibility locus TNFSF4. *PLoS genetics* 2013;9(7):e1003554.

- 53 Hughes T, Kim-Howard X, Kelly JA, et al. Fine-mapping and transethnic genotyping establish IL2/IL21 genetic association with lupus and localize this genetic effect to IL21. *Arthritis Rheum.* 2011;63(6):1689-97.
- 54 Deng Y, Zhao J, Sakurai D, et al. MicroRNA-3148 modulates allelic expression of toll-like receptor 7 variant associated with systemic lupus erythematosus. *PLoS genetics* 2013;9(2):e1003336.
- 55 Lewis MJ, Vyse S, Shields AM, et al. UBE2L3 Polymorphism Amplifies NF-kappaB Activation and Promotes Plasma Cell Development, Linking Linear Ubiquitination to Multiple Autoimmune Diseases. *Am J Hum Genet* 2015;96(2):221-34.
- 56 Sanchez E, Webb RD, Rasmussen A, et al. Genetically determined Amerindian ancestry correlates with increased frequency of risk alleles for systemic lupus erythematosus. *Arthritis Rheum.* 2010;62(12):3722-9.
- 57 Lyn-Cook BD, Xie C, Oates J, et al. Increased expression of Toll-like receptors (TLRs) 7 and 9 and other cytokines in systemic lupus erythematosus (SLE) patients: ethnic differences and potential new targets for therapeutic drugs. *Mol. Immunol.* 2014;61(1):38-43.
- 58 Wiley KL, Treadwell E, Manigaba K, Word B, Lyn-Cook BD. Ethnic differences in DNA methyltransferases expression in patients with systemic lupus erythematosus. *J. Clin. Immunol.* 2013;33(2):342-8.
- 59 Cervera R, Khamashta MA, Font J, et al. Morbidity and mortality in systemic lupus erythematosus during a 5-year period. A multicenter prospective study of 1,000 patients. *European Working Party on Systemic Lupus Erythematosus. Medicine (Baltimore).* 1999;78(3):167-75.
- 60 Mok CC, To CH, Ho LY, Yu KL. Incidence and mortality of systemic lupus erythematosus in a southern Chinese population, 2000-2006. *J. Rheumatol.* 2008;35(10):1978-82.
- 61 Pons-Estel BA, Catoggio LJ, Cardiel MH, et al. The GLADEL multinational Latin American prospective inception cohort of 1,214 patients with systemic lupus erythematosus: ethnic and disease heterogeneity among "Hispanics". *Medicine (Baltimore).* 2004;83(1):1-17.
- 62 Kamen DL, Barron M, Parker TM, et al. Autoantibody prevalence and lupus characteristics in a unique African American population. *Arthritis Rheum.* 2008;58(5):1237-47.
- 63 Sanchez E, Rasmussen A, Riba L, et al. Impact of genetic ancestry and sociodemographic status on the clinical expression of systemic lupus erythematosus in American Indian-European populations. *Arthritis Rheum.* 2012;64(11):3687-94.
- 64 Alba P, Bento L, Cuadrado MJ, et al. Anti-dsDNA, anti-Sm antibodies, and the lupus anticoagulant: significant factors associated with lupus nephritis. *Ann. Rheum. Dis.* 2003;62(6):556-60.
- 65 Hopkinson ND, Jenkinson C, Muir KR, Doherty M, Powell RJ. Racial group, socioeconomic status, and the development of persistent proteinuria in systemic lupus erythematosus. *Ann. Rheum. Dis.* 2000;59(2):116-9.

- 66 Huong DL, Papo T, Beaufils H, et al. Renal involvement in systemic lupus erythematosus. A study of 180 patients from a single center. *Medicine (Baltimore)*. 1999;78(3):148-66.
- 67 Alarcon GS, McGwin G, Jr., Petri M, et al. Time to renal disease and end-stage renal disease in PROFILE: a multiethnic lupus cohort. *PLoS medicine* 2006;3(10):e396.
- 68 Hanly JG, O'Keeffe AG, Su L, et al. The frequency and outcome of lupus nephritis: results from an international inception cohort study. *Rheumatology (Oxford)*. 2016;55(2):252-62.
- 69 Tikly M, Burgin S, Mohanlal P, Bellingan A, George J. Autoantibodies in black South Africans with systemic lupus erythematosus: spectrum and clinical associations. *Clin. Rheumatol.* 1996;15(3):261-5.
- 70 Deligny C, Thomas L, Dubreuil F, et al. [Systemic lupus erythematosus in Martinique: an epidemiologic study]. *Rev. Med. Interne* 2002;23(1):21-9.
- 71 Khanfir MS, Houman MH, Cherif E, et al. TULUP (TUNISIAN LUPUS): a multicentric study of systemic lupus erythematosus in Tunisia. *International journal of rheumatic diseases* 2013;16(5):539-46.
- 72 Al Arfaj AS, Khalil N, Al Saleh S. Lupus nephritis among 624 cases of systemic lupus erythematosus in Riyadh, Saudi Arabia. *Rheumatol. Int.* 2009;29(9):1057-67.
- 73 Malaviya AN, Chandrasekaran AN, Kumar A, Shamar PN. Systemic lupus erythematosus in India. *Lupus* 1997;6(9):690-700.
- 74 Wang F, Wang CL, Tan CT, Manivasagar M. Systemic lupus erythematosus in Malaysia: a study of 539 patients and comparison of prevalence and disease expression in different racial and gender groups. *Lupus* 1997;6(3):248-53.
- 75 Pan Q, Li Y, Ye L, et al. Geographical distribution, a risk factor for the incidence of lupus nephritis in China. *BMC nephrology* 2014;15:67.
- 76 Villamin CA, Navarra SV. Clinical manifestations and clinical syndromes of Filipino patients with systemic lupus erythematosus. *Modern rheumatology / the Japan Rheumatism Association* 2008;18(2):161-4.
- 77 Vila LM, Alarcon GS, McGwin G, Jr., et al. Early clinical manifestations, disease activity and damage of systemic lupus erythematosus among two distinct US Hispanic subpopulations. *Rheumatology (Oxford)*. 2004;43(3):358-63.
- 78 Hanly JG, Urowitz MB, Su L, et al. Seizure disorders in systemic lupus erythematosus results from an international, prospective, inception cohort study. *Ann. Rheum. Dis.* 2012;71(9):1502-9.
- 79 Hanly JG, Su L, Urowitz MB, et al. Mood Disorders in Systemic Lupus Erythematosus: Results From an International Inception Cohort Study. *Arthritis Rheumatol.* 2015;67(7):1837-47.

- 80 Hanly JG, Urowitz MB, O'Keefe AG, et al. Headache in systemic lupus erythematosus: results from a prospective, international inception cohort study. *Arthritis Rheum.* 2013;65(11):2887-97.
- 81 Burgos PI, McGwin G, Jr., Reveille JD, Vila LM, Alarcon GS. Factors predictive of thrombotic events in LUMINA, a multi-ethnic cohort of SLE patients (LXXII). *Rheumatology (Oxford).* 2010;49(9):1720-5.
- 82 Kaiser R, Cleveland CM, Criswell LA. Risk and protective factors for thrombosis in systemic lupus erythematosus: results from a large, multi-ethnic cohort. *Ann. Rheum. Dis.* 2009;68(2):238-41.
- 83 Arroyo-Avila M, Santiago-Casas Y, McGwin G, Jr., et al. Clinical associations of anti-Smith antibodies in PROFILE: a multi-ethnic lupus cohort. *Clin. Rheumatol.* 2015;34(7):1217-23.
- 84 Bruner BF, Guthridge JM, Lu R, et al. Comparison of autoantibody specificities between traditional and bead-based assays in a large, diverse collection of patients with systemic lupus erythematosus and family members. *Arthritis Rheum.* 2012;64(11):3677-86.
- 85 Arnett FC, Hamilton RG, Roebber MG, Harley JB, Reichlin M. Increased frequencies of Sm and nRNP autoantibodies in American blacks compared to whites with systemic lupus erythematosus. *J. Rheumatol.* 1988;15(12):1773-6.
- 86 Jurencak R, Fritzler M, Tyrrell P, Hiraki L, Benseler S, Silverman E. Autoantibodies in pediatric systemic lupus erythematosus: ethnic grouping, cluster analysis, and clinical correlations. *J. Rheumatol.* 2009;36(2):416-21.
- 87 Urowitz MB, Gladman DD, Ibanez D, et al. Evolution of disease burden over five years in a multicenter inception systemic lupus erythematosus cohort. *Arthritis Care Res (Hoboken)* 2012;64(1):132-7.
- 88 Alarcon GS, Calvo-Alen J, McGwin G, Jr., et al. Systemic lupus erythematosus in a multiethnic cohort: LUMINA XXXV. Predictive factors of high disease activity over time. *Ann. Rheum. Dis.* 2006;65(9):1168-74.
- 89 Contreras G, Lenz O, Pardo V, et al. Outcomes in African Americans and Hispanics with lupus nephritis. *Kidney Int.* 2006;69(10):1846-51.
- 90 Plantinga L, Lim SS, Patzer R, et al. Incidence of End-Stage Renal Disease Among Newly Diagnosed Systemic Lupus Erythematosus Patients: The Georgia Lupus Registry. *Arthritis Care Res (Hoboken)* 2016;68(3):357-65.
- 91 Nee R, Martinez-Osorio J, Yuan CM, et al. Survival Disparity of African American Versus Non-African American Patients With ESRD Due to SLE. *Am. J. Kidney Dis.* 2015;66(4):630-7.
- 92 Gomez-Puerta JA, Feldman CH, Alarcon GS, Guan H, Winkelmayr WC, Costenbader KH. Racial and Ethnic Differences in Mortality and Cardiovascular Events Among Patients With End-Stage Renal Disease Due to Lupus Nephritis. *Arthritis Care Res (Hoboken)* 2015;67(10):1453-62.

- 93 Contreras G, Li H, Gonzalez-Suarez M, et al. Kidney allograft survival of African American and Caucasian American recipients with lupus. *Lupus* 2014;23(2):151-8.
- 94 Bruce IN, O'Keeffe AG, Farewell V, et al. Factors associated with damage accrual in patients with systemic lupus erythematosus: results from the Systemic Lupus International Collaborating Clinics (SLICC) Inception Cohort. *Ann. Rheum. Dis.* 2015;74(9):1706-13.
- 95 Ugarte-Gil MF, Acevedo-Vasquez E, Alarcon GS, et al. The number of flares patients experience impacts on damage accrual in systemic lupus erythematosus: data from a multiethnic Latin American cohort. *Ann. Rheum. Dis.* 2015;74(6):1019-23.
- 96 Zhang J, Gonzalez LA, Roseman JM, Vila LM, Reveille JD, Alarcon GS. Predictors of the rate of change in disease activity over time in LUMINA, a multiethnic US cohort of patients with systemic lupus erythematosus: LUMINA LXX. *Lupus* 2010;19(6):727-33.
- 97 Alarcon GS, McGwin G, Jr., Bastian HM, et al. Systemic lupus erythematosus in three ethnic groups. VII [correction of VIII]. Predictors of early mortality in the LUMINA cohort. LUMINA Study Group. *Arthritis Rheum.* 2001;45(2):191-202.
- 98 Bernatsky S, Boivin JF, Joseph L, et al. Mortality in systemic lupus erythematosus. *Arthritis Rheum.* 2006;54(8):2550-7.
- 99 Chambers SA, Allen E, Rahman A, Isenberg D. Damage and mortality in a group of British patients with systemic lupus erythematosus followed up for over 10 years. *Rheumatology (Oxford).* 2009;48(6):673-5.
- 100 Lee YH, Choi SJ, Ji JD, Song GG. Overall and cause-specific mortality in systemic lupus erythematosus: an updated meta-analysis. *Lupus* 2016.
- 101 Merrill JT, Neuwelt CM, Wallace DJ, et al. Efficacy and safety of rituximab in moderately-to-severely active systemic lupus erythematosus: the randomized, double-blind, phase II/III systemic lupus erythematosus evaluation of rituximab trial. *Arthritis Rheum.* 2010;62(1):222-33.
- 102 Dooley MA, Hogan S, Jennette C, Falk R. Cyclophosphamide therapy for lupus nephritis: poor renal survival in black Americans. *Glomerular Disease Collaborative Network. Kidney Int.* 1997;51(4):1188-95.
- 103 Appel GB, Contreras G, Dooley MA, et al. Mycophenolate mofetil versus cyclophosphamide for induction treatment of lupus nephritis. *J. Am. Soc. Nephrol.* 2009;20(5):1103-12.
- 104 Fernandez M, Alarcon GS, Calvo-Alen J, et al. A multiethnic, multicenter cohort of patients with systemic lupus erythematosus (SLE) as a model for the study of ethnic disparities in SLE. *Arthritis Rheum.* 2007;57(4):576-84.
- 105 Sutcliffe N, Clarke AE, Gordon C, Farewell V, Isenberg DA. The association of socio-economic status, race, psychosocial factors and outcome in patients with systemic lupus erythematosus. *Rheumatology (Oxford).* 1999;38(11):1130-7.
- 106 Tobon GJ, Youinou P, Saraux A. The environment, geo-epidemiology, and autoimmune disease: Rheumatoid arthritis. *Autoimmunity reviews* 2010;9(5):A288-92.

- 107 Watts RA, Gonzalez-Gay MA, Lane SE, Garcia-Porrúa C, Bentham G, Scott DG. Geoepidemiology of systemic vasculitis: comparison of the incidence in two regions of Europe. *Ann. Rheum. Dis.* 2001;60(2):170-2.
- 108 Khairallah M, Accorinti M, Muccioli C, Kahloun R, Kempen JH. Epidemiology of Behcet disease. *Ocul. Immunol. Inflamm.* 2012;20(5):324-35.
- 109 Albor C, du Sautoy T, Kali Vanan N, Turner BP, Boomla K, Schmierer K. Ethnicity and prevalence of multiple sclerosis in east London. *Mult. Scler.* 2016.
- 110 Milo R, Kahana E. Multiple sclerosis: geoepidemiology, genetics and the environment. *Autoimmunity reviews* 2010;9(5):A387-94.

Figure legends

Fig. 1. Epidemiological studies of SLE in different countries

Incidence (A) and prevalence (B) of SLE in epidemiological studies across the world. Where multiple studies are available for a country or region, the largest and/or most recent data are represented. Error bars show 95% confidence intervals for incidence and prevalence. Where confidence intervals were not published, they were calculated based on case number and incidence/prevalence. *Countries/administrative regions with nationwide data.

Fig. 2. Choropleth map showing prevalence of SLE around the world

Countries lacking prevalence data are shown in grey. Coloured circles highlight prevalence data from Puerto Rico, Martinique and Hong Kong.

Fig. 3. Bubble chart summarising studies of incidence and prevalence of SLE over time

Linear regression models (white line) were calculated in R, with region within 95% confidence interval shaded grey.

Figure 1





Figure 2




Figure 3

