

This is a post-peer-review, pre-copyedit version of an article published in Supportive Care in Cancer. The final authenticated version is available online at:
<http://dx.doi.org/10.1007/s00520-018-4146-9>

Post-Traumatic Stress in Head and Neck Cancer Survivors and their Partners

Running Title: Post-Traumatic Stress in HNC Survivors and their Partners

Authors: Elisavet Moschopoulou ¹, Iain Hutchison ², Kamaldeep Bhui ¹, Ania Korszun ¹

Affiliations

1. Centre for Psychiatry, Wolfson Institute of Preventive Medicine, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, EC1A 7BE
2. Oral and Maxillofacial Surgery, The Royal London Hospital, Barts Health NHS Trust, Whitechapel, London, E1 1BB

Corresponding author:

Elisavet Moschopoulou

elisavet.moschopoulou@qmul.ac.uk

+44 (0)20 7882 2023

ORCID ID:

Elisavet Moschopoulou: 0000-0003-3568-3748

Kamaldeep Bhui: 0000-0002-9205-2144

Ania Korszun: 0000-0001-9253-6491

Acknowledgments: This work was supported by a Doctoral Scholarship from Saving Faces – The Facial Surgery Research Foundation.

ABSTRACT

Purpose: Head and neck cancer (HNC) diagnosis and treatment are distressing and have immediate detrimental impacts on functioning and quality of life (QoL). Nevertheless, little is known about long-term psychosocial effects. The aim of this study was to determine the prevalence and correlates of clinical post-traumatic stress disorder (PTSD) and subclinical post-traumatic stress symptoms (PTSS) in HNC patients surviving more than 2 years since treatment and in their partners.

Methods: HNC survivors identified from the cancer registry of a London hospital and their partners completed measures of PTSS, depression and anxiety, fear of cancer recurrence, social support, appearance concerns and health-related QoL. Data regarding their clinical and demographic characteristics were also collected. Correlations, as well as linear and logistic regression coefficients, were calculated to estimate associations with PTSS scores.

Results: In this analysis of 93 HNC survivors, at a mean of 6 years ($SD=4$) after treatment, 33.4% reported PTSS and 11.8% met criteria for post-traumatic stress disorder (PTSD). Fear of cancer recurrence was independently associated with PTSS ($p<.01$). In subgroup analyses of patient-partner dyads 15.4% of patients and 12.8% of partners reported PTSD, with a further 33.3% of patients and 25.7% of partners demonstrating PTSS. Patients' and partners' scores did not differ significantly ($p >.05$).

Conclusions: This is the first examination of post-traumatic stress in survivors of HNC and shows that high levels of cancer-related PTSS exist for many years after diagnosis in both patients and their partners.

Keywords: Cancer, Head and Neck Neoplasms, Survivors, Caregivers, Post-Traumatic Stress Disorder

BACKGROUND

Head and Neck Cancer (HNC) is currently the sixth most common cancer worldwide^[1,2] with a steadily increasing incidence^[3]. Five-year survival rates have risen to around 50% with an increase of cases related to human papillomavirus (HPV) infection. HPV-positive HNC patients are much younger and have a better prognosis^[4,5]. Thus, there are more HNC survivors facing long-term sequelae of HNC treatment. HNC is now better viewed as a chronic disease making quality of life (QoL) and overall wellbeing as treatment outcomes highly relevant. Depending on the site, HNC can significantly impact eating, drinking, swallowing, appearance and social interaction, with marked adverse effects on QoL and psychosocial function^[3,6,7].

The focus of HNC studies has been more on the potential demographic, functional and clinical characteristics associated with survivorship and QoL^[8], whereas psychosocial factors, particularly depression and anxiety, are the factors most strongly associated with poorer QoL^[9]. In survivors (i.e. those who have completed their primary treatment for HNC), prevalence rates for clinically significant symptoms of depression and anxiety are reported as 19-31% and 16% respectively^[10,9]. Caregivers of HNC patients are also at risk for psychological distress and previous studies have suggested that caregivers may suffer equal or even higher levels of distress than patients^[11,12]. A diagnosis of HNC places a significant burden on the patients' carers^[11], but less is known about the specific issues they face.

HNC treatment can be intrusive resulting in permanent functional and disfiguring changes, thereby increasing the likelihood of a post-traumatic stress reaction. Despite the potentially highly traumatic nature of HNC^[13], the prevalence of post-traumatic stress symptoms (PTSS) and post-traumatic stress disorder (PTSD) in long-term HNC survivors has not been investigated. PTSD is a psychiatric condition that affects some people when they have witnessed or experienced a traumatic event involving the risk of serious injury or death^[14]. Symptoms such as flashbacks, avoiding cancer-related experiences, and increased anxiety, have been reported in other cancer survivor groups with PTSD prevalence rates of 5-17%^[15-18]. In newly diagnosed HNC patients, 12% of patients and 29% of partners met criteria for PTSD caseness^[19]. Richardson and colleagues (2016) found that 19% of HNC patients^[20] and their caregivers^[21] met criteria for PTSD six months after treatment. To the best of the authors' knowledge there have been no studies investigating whether cancer-related PTSS exist in long-term HNC survivors and in their partners. Therefore, the purpose of this study

was to (i) estimate the prevalence of PTSS in a cohort of British HNC survivors, (ii) identify potential demographic, clinical and psychosocial correlates of PTSS and (iii) compare PTSS levels in a sub-group of HNC survivors and their partners.

METHODS

Participants

Prospective participants were identified through the cancer registry at Barts Health NHS Trust. Eligible patients were those who had received a diagnosis of HNC, had completed active treatment (i.e. were in the follow-up phase), and were ≥ 2 years after diagnosis. Potential participants were sent an invitation letter asking whether they would be willing for the research team to invite their spouse to the study.

The study questionnaire along with the information sheet and consent forms were mailed to those interested in participating. Patients and their partners were provided with separate postage-paid return envelopes. No financial incentives or any other forms of compensation were used. Ethics approval was obtained from the Office for Research Ethics Committees Northern Ireland (ORECNI) (Reference: 15/NI/0009).

Measures

Demographic and clinical variables

Sociodemographic information sought included gender, ethnicity, education, marital status, and work status. Clinical variables included cancer site and stage, treatment type, time since treatment and whether there had been a recurrence. Medical comorbidity was measured using the Self-Administered Comorbidity Questionnaire (SCQ)[22]. Participants' self-reported history of anxiety or depression preceding their HNC diagnosis were also evaluated.

Psychosocial Variables and QoL

The Hospital Anxiety and Depression Scale (HADS)^[23], which has been validated for use in cancer patients^[24], was selected to assess participants' levels of psychological distress. This instrument consists of two subscales: 7 items assessing levels of depressive symptoms and 7 items for levels of anxiety. Higher scores (range = 0-21) indicate increased distress. Social

support (SS) was measured using the ENRICH Social Support Inventory (ESSI)^[25], a reliable 7-item scale previously used in cancer patients^[26]. A total score can be produced by summation, with greater scores representing greater social support. Fear of cancer recurrence (FoR) was assessed using a 7-item questionnaire measuring worries associated with the risk of cancer coming back and their impact on daily life^[27]. This instrument has been previously used in HNC patients^[28]. Scores range from 6-40 with higher scores representing greater FoR. The Assessment of Life Threat and Treatment Intensity (ALTTIQ)^[29] is a 7-item questionnaire used to examine how threatening and intense patients perceive cancer and its treatment to be^[17,30]. Responses are summed and scores range from 7-35. The Global QoL scale of the EORTC QLQ-C30 Version 3^[31] was used as the main QoL outcome. Responses are transformed to fit a 0-100 distribution. The QLQ-C30 has been extensively used in studies of HNC patients^[32]. Finally, the Derriford Appearance Scale (DAS24)^[33] was used to measure levels of appearance concerns. The total score ranges from 11 to 96, with lower scores reflecting low levels of social anxiety and avoidant behaviour.

Post-traumatic Stress

The PTSD Checklist-Civilian Version (PCL-C)^[34] is a validated 17-item self-report scale assessing the severity of traumatic stress symptoms^[35]. Responses range from 1 “Not at all” to 5 “Extremely” and a total score (range =17-85) can be obtained by summation. Scores ≥ 44 are classified as indicative of PTSD for adults who have experienced acute trauma^[36]. The PCL reflects the DSM-IV (Diagnostic and Statistical Manual of Mental Disorders – Fourth Edition) symptoms of PTSD. Using the cluster scoring method, a predefined level of symptoms in each symptom cluster (i.e. intrusive thoughts, avoidant behaviour, hyperarousal) needs to be endorsed in order for the scores to indicate PTSD^[19]. Meeting the threshold in two symptom clusters indicates partial PTSD (i.e. PTSS)^[17]. Questions were modified to reflect cancer and its treatment as the traumatic stressor of interest.

Statistical Methods

In bivariate analyses, t-tests and analysis of variance (ANOVA) were used in order to compare mean levels of PCL-C scores. Pearson’s correlation coefficient was used to examine associations between continuous variables and the PCL-C. Potential correlates of PTSS were then assessed using multiple linear regression analyses with patients’ PCL-C score as the outcome variable. The association of the same variables with full or partial PTSD was assessed using binary logistic regression. A stepwise modelling approach was followed

whereby only variables that were significantly associated ($p < .05$) with PCL-C in bivariate analyses were included in multiple regression models.

Regarding the dyadic data analyses, differences between patients' and partners' scores on the PCL-C, as well as on their depression and anxiety scores, were examined using paired t-tests and McNemar's χ^2 test, as appropriate. Mann Whitney U and Kruskal-Wallis H tests were used for comparisons within subgroups due to their small sample size. Correlation coefficients were calculated to examine the association between all continuous variables and participants' PCL-C scores. Agreement between patients' and partners' levels of post-traumatic stress was assessed using weighted Cohen's Kappa. In addition, overall PCL-C scores in dyads were examined for absolute agreement using the two-way mixed-effects intraclass correlation coefficient (ICC). All data were coded and analysed using IBM SPSS Statistics Version 23.

RESULTS

As shown in the consort diagram (Figure 1), the final study sample consisted of 93 HNC survivors while the dyadic analysis was conducted on data from a subsample of 39 patient-partner dyads. There was no time lag between patients' and partners' return of the completed questionnaire. Table 1 presents the sample characteristics. Patients' mean (\pm SD) age at enrolment was 66 years (11) and the overall male to female ratio was 1:0.72. The majority of patients were White British (84.8%) and their education was at least of high school level. Sixty eight percent were partnered and 66% were not in paid employment. Patients were at a mean (\pm SD) of 6.3 years (4) after treatment and 56% of them had an oral cavity tumour. The majority had early stage cancer (i.e. stage one or two) (66%) and did not have a recurrence (83%). Eighty seven percent were treated with surgery either alone or with adjuvant chemo-radiation. Fifteen percent reported a history of anxiety while 25.8% reported a history of depression.

Prevalence of Post-traumatic Stress in HNC survivors and their partners

In this sample of HNC survivors (N=93), the mean (\pm SD) PCL-C score was 28.2 (11.5). As shown in Table 2, 11.8% of patients scored ≥ 44 indicating the presence of PTSD. Based on

the cluster scoring method, 12.9% of patients met the criteria for estimated PTSD caseness. A further 10.8% rated two out of three PTSD symptom clusters as moderately to extremely reoccurring (i.e. 'partial PTSD'). Overall, 46.3% met criteria for at least one PTSD symptom cluster. Across the symptom clusters, 33.3% met criteria for intrusion, 29% for hyperarousal, and 20.4% for avoidance.

With regards to the prevalence of post-traumatic stress in the subsample of 39 patient-partner dyads, the mean (\pm SD) PCL-C score was 28.6 (10.4) for patients and 27.8 (12.5) for partners. Patients' and partners' PTSS scores were not significantly different ($t(38) = .35, p = .731$) and did not correlate with each other ($r = .23, p = .15$). Thirteen percent of patients and 15.4% of partners scored ≥ 44 indicating PTSD caseness.

Using the cluster scoring method, 6/39 (15.4%) of patients and 5/39 (12.8%) of partners endorsed the threshold symptom level reflecting PTSD caseness. A further 15.4% of patients and 10.3% of partners rated two out of three PTSD symptom clusters as moderately to extremely reoccurring (partial PTSD). Overall, 48.7% of patients and 38.5% of partners met criteria for at least one PTSD symptom cluster. Both patients and carers rated more highly symptoms of intrusion, with 35.9% and 33.3% respectively meeting the threshold. Patients endorsed more symptoms of avoidance (25.6%) and hyperarousal (33.3%) in comparison to their partners (12.8% and 28.2% respectively) however the difference was not statistically significant (Table 3).

Concordance analysis for levels of post-traumatic stress (i.e. symptomatic in 3 symptom cluster, symptomatic in 2 symptom clusters, symptomatic in 1 symptom cluster and asymptomatic) between patients and their partners showed that the strength of agreement was poor ($\kappa_w = .187, p = .127$). Similarly, the ICC test showed little agreement between patients' and partners' overall PCL-C scores (ICC = .378, 95% CI (-.202 to .676), $F(38,38) = 1.59, p = .078$). Dyads with male caregivers ($n = 13$) appeared more congruent than dyads with female caregivers ($n = 26$) however this did not reach statistical significance.

Bivariate Associations with PTSS

Table 1 presents the bivariate associations between patients' total PCL-C scores and the variable categories of interest. There were no significant associations between clinical variables and levels of post-traumatic stress.

Among demographic variables, a negative PCL-C association was found for age at the time of completing the questionnaire. There were no other significant associations between demographic variables and PCL-C scores.

Among psychosocial variables, there were significant PCL-C associations for FoR, symptoms of anxiety and depression, as well as for appearance concerns. Levels of perceived threat were also highly positively correlated with patients' PCL-C scores. SS levels were not significantly associated with PCL-C scores. Finally, there was a large negative correlation between PCL-C scores and QoL.

Multivariable Analyses

Table 4 presents the results of the multiple linear regression analysis for patients' PCL-C scores and the variables that were statistically significant in bivariate analyses ($p < .05$). Results of a binary logistic regression for full or partial PTSD and the same explanatory factors are also presented in Table 4. Although the full linear regression model accounted for 73.4% of the variance in PCL-C scores ($p < .001$), FoR was the only independent correlate of PTSS. There was a favourable statistical trend ($p < .1$) for symptoms of anxiety, QoL and appearance concerns however these were not statistically significant ($p > .05$). In the logistic regression model, FoR was the only significant correlate of PTSS.

Dyads Subsample: Bivariate Analyses

Although we were underpowered to detect differences in PTSS levels between patients and their partners or to investigate interactions within subgroups of partners, we carried out bivariate analyses to identify any trends in the data.

With regards to their background characteristics, patients' and partners' symptoms of depression and anxiety, as well as their SS levels did not differ significantly (anxiety, $t(38) = 1.17$, $p = .25$; depression, $t(38) = .04$, $p = .97$; SS $t(38) = 1.79$, $p = .082$) and were not significantly associated with each other (anxiety, $r = -.006$, $p = .97$; depression, $r = .23$, $p = .17$; SS, $r = .29$, $p = .07$). Furthermore, 10% of patients and 12.8% of partners reported previous anxiety while 23% of patients and 23% of partners reported previous depression.

Regarding potential correlates of PTSS levels in partners, between the clinical variables, prior history of depression was associated with higher PCL-C scores ($U = 64$, $p = < .05$). Among demographics the only significant relationship was for gender ($U = 52$, $p = < .01$) indicating

that female partners (n=26) experienced more severe PTSS than their male counterparts (n=13).

Among psychosocial variables, significant PCL-C relationships were found for symptoms of depression ($r = .570$, $p = <.01$), anxiety ($r = .766$, $p = <.01$) and levels of social support ($r = -.560$, $p = <.01$).

Regarding possible interactions between partners' PCL-C scores and patients' characteristics or vice versa, there was a statistically significant relationship between partners' PCL-C scores and patients' QoL ($r = -.329$, $p = <.05$). Patients' PCL-C score was associated only with partners' SS levels ($r = -.445$, $p = <.01$).

CONCLUSIONS

This is the first study to examine post-traumatic stress in long-term HNC survivors and their partners. In this sample of HNC survivors, 13% of patients met diagnostic criteria for PTSD, with an additional 33% experiencing some PTSS. This is a much higher prevalence in comparison to that of the general adult population in the UK (4.4%)^[37] but similar to other cancer survivor populations such as haematological (8%) or breast (12%) cancer^[17,16]. Our findings are also comparable to those of Posluszny et al (2015)[19] who reported a 12% prevalence of PTSD in HNC patients approximately two months after diagnosis, though Richardson et al (2016)[20] in their study of HNC patients 6 months after treatment reported a higher rate of 19%. In our sample, mean time since treatment was 6 years suggesting that PTSS may persist for many years after diagnosis. In the context of the growing literature on PTSD/PTSS amongst HNC patients, longitudinal research is needed in order to understand better the duration and course of these symptoms.

Results from our subgroup analyses showed that 15.4% of patients and 12.8% of partners reported PTSD. Scores from a further 33.3% of patients and 25.7% of partners were indicative of PTSS. These results are not in line with the findings of Posluszny et al (2015) who reported significantly higher levels of PTSD in partners than in patients. Nevertheless, Richardson's findings suggested a similar prevalence rate between HNC patients and their caregivers [21,20]. Further research employing greater sample sizes is needed.

Clinical correlates of PTSS and PTSD

In bivariate analysis, younger age was significantly associated with higher PTSS; consistent with findings from the PTSD literature^[35]. Younger survivors may face more financial and social challenges^[17] and the emotional response of older HNC patients is more positive^[38]. It is striking that there was no significant effect of cancer stage, site or treatment. This is in agreement with previous findings^[35,21] and emphasises that PTSS experienced by cancer survivors may not consequently be connected to the severity of cancer or to socio-demographic factors. Contrary to previous findings, time since treatment was not associated with reduced PTSS^[17,39]. Although patients with previous depression and anxiety were at higher risk for psychological distress and PTSD, this did not have a significant impact on PTSS scores. Reports of PTSS are more likely to reflect difficulties in adjustment to cancer^[40]. Medical comorbidity has been shown to be a risk factor for PTSS^[17] however this was not replicated in our sample.

FoR was the strongest correlate of post-traumatic stress. FoR is a common emotional response to cancer but differs from a psychopathological response that is characterised by features of avoidance and hyperarousal such as PTSD. Nevertheless, FoR, depending on its intensity, can be difficult to distinguish from anxiety and PTSS.

QoL, anxiety and appearance concerns showed significant but weak trends consistent with previous research^[40]. In our sample, mean patient QoL was very close to the 2014 European general population mean (\pm SD) of 75.7 (21.2)[41]. Other studies have reported a negative correlation between appearance concerns and depression^[42], however, our study is the first to present findings on the association with PTSS. As the face is difficult to avoid, facial disfigurement may be acting as a constant reminder and trigger of trauma.

Survivors and their partners experienced similar levels of PTSS but they showed little agreement in their appraisals of post-traumatic stress symptoms. Trends in the data indicated that female partners, those with a history of depression and those with less SS showed higher PTSS. Higher PTSS in partners was also related to more depressive symptoms, anxiety and lower patient QoL. Although further investigation is needed in order to confirm these patterns, our findings are in line with previous research that suggests a proportion of HNC partners experience poorer psychological health in comparison to population norms and that caregivers are affected by patients' well-being (and vice versa)^[11,12]. Further study of the factors affecting patient-caregiver PTSS congruence is warranted as this may influence coping with cancer overall.

Study Limitations

This study has some limitations, including the cross-sectional design, the small sample size and the inclusion of patients from one cancer centre. The cross-sectional design limits our ability to infer causal relationships or to draw conclusions about the duration of PTSS since diagnosis. Although the inclusion of partners is a strength, given how little we know about partners' emotional response to HNC, our sample size was too small for sub-group analyses amongst patients or dyads. Our participants were recruited from a single centre, limiting the generalisability of our results. Furthermore, our sample consisted of predominantly white patients and the recurrence rate was low which may further affect the generalisability of our findings. Nevertheless, our results show levels slightly higher than those reported in other cancer survivor groups ^[17,35]. Finally, the assessment of PTSS/PTSD in this study was based on the diagnostic criteria of DSM-IV. This is a limitation as there are notable changes to the PTSD criteria in DSM-V. Therefore relevancy of these finding to DSM-V is not known. However, our results add to the existing literature on the prevalence of cancer-related PTSD and allow a reliable comparison of PTSS/PTSD rates between HNC survivors and different patient groups, as well as between HNC survivors and newly diagnosed patients.

In conclusion, we found that in some HNC patients, symptoms of cancer-related PTSD exist for many years and also affect caregivers. Clinicians and researchers need to be aware of this type of psychological response to HNC in survivors and take this into account when addressing the needs of patients and their partners. PTSD is a treatable condition and referral to psychological services should be considered. Psychological interventions to reduce PTSD symptoms in HNC patients have shown some effectiveness [43,44] but further research is required for the development of targeted interventions that can also be used at an earlier stage to prevent PTSS in cancer survivors. Our study has identified potentially modifiable factors that may play a role in the development of cancer-related PTSD – namely, fear of recurrence. Understanding the issues HNC survivors and their caregivers face will enable identification of those at highest risk of suffering from post-traumatic stress.

Compliance with Ethical Standards

Ethical Approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee

and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent: Informed consent was obtained from all individual participants included in the study.

Conflict of Interest Statement: The authors have no conflict of interest to declare.

REFERENCES

1. Pfster DG, Spencer S, Brizel DM, Burtness B, Busse PM, Caudell JJ, Cmelak AJ, Colevas AD, Dunphy F, Eisele DW, Foote RL, Gilbert J, Gillison ML, Haddad RI, Haughey BH, Hicks WL, Hitchcock YJ, Jimeno A, Kies MS, Lydiatt WM, Maghami E, McCaffrey T, Mell LK, Mittal BB, Pinto HA, Ridge JA, Rodriguez CP, Samant S, Shah JP, Weber RS, Wolf GT, Worden F, Yom SS, McMillian N, Hughes M (2015) Head and Neck Cancers, Version 1.2015: Featured Updates to the NCCN Guidelines. *J Natl Compr Canc Netw* 13 (7):847-856
2. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *International journal of cancer Journal international du cancer* 136 (5):E359-386. doi:10.1002/ijc.29210
3. National Head and Neck Cancer Audit (DAHNO) (2015) National Head and Neck Cancer Audit - 10th Annual Report. Health and Social Care Information Centre, Leeds
4. Boscolo-Rizzo P, Pawlita M, Holzinger D (2016) From HPV-positive towards HPV-driven oropharyngeal squamous cell carcinomas. *Cancer Treat Rev* 42:24-29. doi:10.1016/j.ctrv.2015.10.009
5. Friedman JM, Stavas MJ, Cmelak AJ (2014) Clinical and scientific impact of human papillomavirus on head and neck cancer. *World J Clin Oncol* 5 (4):781-791. doi:10.5306/wjco.v5.i4.781
6. Howren MB, Christensen AJ, Karnell LH, Funk GF (2013) Psychological factors associated with head and neck cancer treatment and survivorship: Evidence and opportunities for behavioral medicine. *Journal of Consulting and Clinical Psychology* 81 (2):299-317. doi:10.1037/a0029940
7. Semple C, Parahoo K, Bessell A, McCaughan E, Humphris G, Mills M (2013) Cochrane review: Effectiveness of psychosocial interventions for patients with head and neck cancer. *European Journal of Cancer* 49:754-755
8. Rettig EM, D'Souza G, Thompson CB, Koch WM, Eisele DW, Fakhry C (2016) Health-related quality of life before and after head and neck squamous cell carcinoma: Analysis of the Surveillance, Epidemiology, and End Results-Medicare Health Outcomes Survey linkage. *Cancer* 122 (12):1861-1870. doi:10.1002/cncr.30005
9. Dunne S, Mooney O, Coffey L, Sharp L, Desmond D, Timon C, O'Sullivan E, Gallagher P (2016) Psychological variables associated with quality of life following primary treatment for head and neck cancer: a systematic review of the literature from 2004 to 2015. *Psychooncology*. doi:10.1002/pon.4109

10. Neilson KA, Pollard AC, Boonzaier AM, Corry J, Castle DJ, Mead KR, Gray MCL, Smith DI, Trauer T, Couper JW (2010) Psychological distress (depression and anxiety) in people with head and neck cancers. *Medical Journal of Australia* 193 (5):48-51
11. Longacre ML, Ridge JA, Burtness BA, Galloway TJ, Fang CY (2012) Psychological functioning of caregivers for head and neck cancer patients. *Oral Oncology* 48 (1):18-25. doi:10.1016/j.oraloncology.2011.11.012
12. Sterba KR, Zapka J, Cranos C, Laursen A, Day TA (2015) Quality of Life in Head and Neck Cancer Patient-Caregiver Dyads: A Systematic Review. *Cancer Nurs.* doi:10.1097/ncc.0000000000000281
13. Björklund M, Sarvimäki A, Berg A (2010) Living with head and neck cancer: a profile of captivity. *Journal of Nursing and Healthcare of Chronic Illness* 2 (1):22-31. doi:10.1111/j.1752-9824.2010.01042.x
14. American Psychiatric Association (2013) *Diagnostic and Statistical Manual of Mental Disorders: Fifth Edition.* Arlington
15. Black EK, White CA (2005) Fear of recurrence, sense of coherence and posttraumatic stress disorder in haematological cancer survivors. *Psychooncology* 14 (6):510-515. doi:10.1002/pon.894
16. O'Connor M, Christensen S, Jensen AB, Moller S, Zachariae R (2011) How traumatic is breast cancer? Post-traumatic stress symptoms (PTSS) and risk factors for severe PTSS at 3 and 15 months after surgery in a nationwide cohort of Danish women treated for primary breast cancer. *Br J Cancer* 104 (3):419-426. doi:10.1038/sj.bjc.6606073
17. Smith SK, Zimmerman S, Williams CS, Preisser JS, Clipp EC (2008) Post-traumatic stress outcomes in non-Hodgkin's lymphoma survivors. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 26 (6):934-941. doi:10.1200/jco.2007.12.3414
18. Kornblith AB, Herndon JE, 2nd, Weiss RB, Zhang C, Zuckerman EL, Rosenberg S, Mertz M, Payne D, Jane Massie M, Holland JF, Wingate P, Norton L, Holland JC (2003) Long-term adjustment of survivors of early-stage breast carcinoma, 20 years after adjuvant chemotherapy. *Cancer* 98 (4):679-689. doi:10.1002/cncr.11531
19. Posluszny DM, Dougall AL, Johnson JT, Argiris A, Ferris RL, Baum A, Bovbjerg DH, Dew MA (2015) Posttraumatic stress disorder symptoms in newly diagnosed patients with head and neck cancer and their partners. *Head Neck* 37 (9):1282-1289. doi:10.1002/hed.23760
20. Richardson AE, Morton RP, Broadbent E (2016) Coping strategies predict post-traumatic stress in patients with head and neck cancer. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery.* doi:10.1007/s00405-016-3960-2
21. Richardson AE, Morton RP, Broadbent EA (2016) Illness perceptions and coping predict post-traumatic stress in caregivers of patients with head and neck cancer. *Support Care Cancer.* doi:10.1007/s00520-016-3285-0
22. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN (2003) The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum* 49 (2):156-163. doi:10.1002/art.10993
23. Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. *Acta psychiatrica Scandinavica* 67 (6):361-370
24. Mitchell AJ, Meader N, Symonds P (2010) Diagnostic validity of the Hospital Anxiety and Depression Scale (HADS) in cancer and palliative settings: a meta-analysis. *Journal of affective disorders* 126 (3):335-348. doi:10.1016/j.jad.2010.01.067
25. Mitchell PH, Powell L, Blumenthal J, Norton J, Ironson G, Pitula CR, Froelicher ES, Czajkowski S, Youngblood M, Huber M, Berkman LF (2003) A short social support measure for patients recovering from myocardial infarction: the ENRICH Social Support Inventory. *Journal of cardiopulmonary rehabilitation* 23 (6):398-403
26. Zhou ES, Penedo FJ, Lewis JE, Rasheed M, Traeger L, Lechner S, Soloway M, Kava BR, Antoni MH (2010) Perceived stress mediates the effects of social support on health-related quality of life among

- men treated for localized prostate cancer. *J Psychosom Res* 69 (6):587-590.
doi:10.1016/j.jpsychores.2010.04.019
27. Ghazali N, Cadwallader E, Lowe D, Humphris G, Ozakinci G, Rogers SN (2013) Fear of recurrence among head and neck cancer survivors: longitudinal trends. *Psycho-Oncology* 22 (4):807-813.
doi:10.1002/pon.3069
28. Rogers SN, Scott B, Lowe D, Ozakinci G, Humphris GM (2010) Fear of recurrence following head and neck cancer in the outpatient clinic. *European Archives of Oto-Rhino-Laryngology* 267 (12):1943-1949. doi:10.1007/s00405-010-1307-y
29. Stuber ML, Kazak AE, Meeske K, Barakat L, Guthrie D, Garnier H, Pynoos R, Meadows A (1997) Predictors of posttraumatic stress symptoms in childhood cancer survivors. *Pediatrics* 100 (6):958-964
30. Smith SK, Zimmerman S, Williams CS, Benecha H, Abernethy AP, Mayer DK, Edwards LJ, Ganz PA (2011) Post-traumatic stress symptoms in long-term non-Hodgkin's lymphoma survivors: does time heal? *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 29 (34):4526-4533. doi:10.1200/jco.2011.37.2631
31. Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, Filiberti A, Flechtner H, Fleishman SB, de Haes JC, et al. (1993) The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *Journal of the National Cancer Institute* 85 (5):365-376
32. Bjordal K, de Graeff A, Fayers PM, Hammerlid E, van Pottelsberghe C, Curran D, Ahlner-Elmqvist M, Maher EJ, Meyza JW, Bredart A, Soderholm AL, Arraras JJ, Feine JS, Abendstein H, Morton RP, Pignon T, Huguenin P, Bottomly A, Kaasa S (2000) A 12 country field study of the EORTC QLQ-C30 (version 3.0) and the head and neck cancer specific module (EORTC QLQ-H&N35) in head and neck patients. EORTC Quality of Life Group. *European journal of cancer (Oxford, England : 1990)* 36 (14):1796-1807
33. Carr T, Moss T, Harris D (2005) The DAS24: A short form of the Derriford Appearance Scale DAS59 to measure individual responses to living with problems of appearance. *British Journal of Health Psychology* 10 (2):285-298. doi:10.1348/135910705X27613
34. Weathers F, Litz B, Herman D, Huska J, Keane T (1994) The PTSD checklist-civilian version (PCL-C). Boston, MA: National Center for PTSD
35. Hahn EE, USA UoCUFSopHLAC, USA KPSCDoRaEPC, Hays RD, USA UoCUFSopHLAC, University of California Department of Medicine DoGIMaHSR, David Geffen School of Medicine Los Angeles CA USA, USA RCSMC, Kahn KL, University of California Department of Medicine DoGIMaHSR, David Geffen School of Medicine Los Angeles CA USA, USA RCSMC, Litwin MS, USA UoCUFSopHLAC, University of California Department of Urology DGSoMLACU, Ganz PA, USA UoCUFSopHLAC, USA UoCDoCPaCRJCCCLAC (2016) Post-traumatic stress symptoms in cancer survivors: relationship to the impact of cancer scale and other associated risk factors. *Psycho-Oncology* 24 (6):643-652.
doi:10.1002/pon.3623
36. Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA (1996) Psychometric properties of the PTSD Checklist (PCL). *Behaviour research and therapy* 34 (8):669-673
37. McManus S, Bebbington P, Jenkins R, Brugha T (2016) Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. Leeds: NHS Digital
38. Rogers SN, Audisio RA, Lowe D (2015) Do the elderly raise different issues when using the Patient Concerns Inventory in routine head and neck cancer follow-up clinics? *European Journal of Cancer Care* 24 (2):189-197. doi:10.1111/ecc.12289
39. Abbey G, Thompson SB, Hickish T, Heathcote D (2015) A meta-analysis of prevalence rates and moderating factors for cancer-related post-traumatic stress disorder. *Psychooncology* 24 (4):371-381. doi:10.1002/pon.3654
40. Cordova MJ, Riba MB, Spiegel D (2017) Post-traumatic stress disorder and cancer. *The Lancet Psychiatry*. doi:[http://dx.doi.org/10.1016/S2215-0366\(17\)30014-7](http://dx.doi.org/10.1016/S2215-0366(17)30014-7)

41. Hinz A, Singer S, Brahler E (2014) European reference values for the quality of life questionnaire EORTC QLQ-C30: Results of a German investigation and a summarizing analysis of six European general population normative studies. *Acta oncologica (Stockholm, Sweden)* 53 (7):958-965. doi:10.3109/0284186x.2013.879998
42. Clarke S-A, Newell R, Thompson A, Harcourt D, Lindenmeyer A (2014) Appearance concerns and psychosocial adjustment following head and neck cancer: A cross-sectional study and nine-month follow-up. *Psychology Health & Medicine* 19 (5):505-518. doi:10.1080/13548506.2013.855319
43. Kangas M, Milross C, Taylor A, Bryant RA (2013) A pilot randomized controlled trial of a brief early intervention for reducing posttraumatic stress disorder, anxiety and depressive symptoms in newly diagnosed head and neck cancer patients. *Psycho-Oncology* 22 (7):1665-1673. doi:10.1002/pon.3208
44. Kangas M, Milross C, Bryant RA (2014) A brief, early cognitive-behavioral program for cancer-related ptsd, anxiety, and comorbid depression. *Cognitive and Behavioral Practice* 21 (4):416-431

Table 1 Characteristics of the patient sample and bivariate associations with PTSS (PCL-C scores)

Variable	Mean (SD), Range	Frequency (Valid %)	PCL-C Mean (SD)	Correlation (r) †	p-value‡
Age at enrolment, years (n = 93)	66 (11), 30-92			-.249	.016
Gender (n = 93)					.205
Male		54 (58.1)	29.5 (12.1)		
Female		39 (41.9)	26.4 (10.4)		
Ethnicity (n = 92, missing = 1)					.694
White British		78 (84.8)	28.2 (11.3)		
White Other		7 (7.6)	25.7 (9.1)		
Asian		5 (5.4)	32.6 (15.5)		
Black		2 (2.2)	33.5 (23.3)		
Education (n = 77, missing = 16)					.527
>High School		40 (51.9)	29.4 (11.3)		
≤High School		37 (48.1)	27.6 (12.5)		
Relationship Status (n = 92, missing = 1)					.186
Partnered		63 (67.7)	29.3 (12.3)		
Non-partnered		29 (31.2)	26.2 (9.3)		
Work Status (n = 91, missing = 2)					.359
Employed		31 (34.1)	29.9 (12.2)		
Unemployed		60 (65.9)	27.6 (11.2)		
Site (n = 93)					.158
Oral Cavity		52 (55.9)	28.4 (12)		
Oropharynx		22 (23.7)	26.8 (8.2)		
Nasopharynx		4 (4.3)	34 (15.6)		
Nasal Cavity		5 (5.4)	21.8 (4.6)		
Salivary gland		2 (2.2)	27 (11.3)		
Hypopharynx		2 (2.2)	29 (2.8)		
Larynx		2 (2.2)	17.5 (.1)		
Unknown primary		4 (4.3)	42 (17.9)		
Stage (n = 87, missing = 6)					.640
Tx (tumour cannot be evaluated)		7 (8)	34.3 (17.9)		
T1		23 (26.4)	28.6 (11)		
T2		34 (39.1)	26.7 (10)		
T3		10 (11.5)	28.1 (9.2)		
T4		13 (14.9)	28.4 (14)		

Variable	Mean (SD), Range	Frequency (Valid %)	PCL-C Mean (SD)	Correlation (r) †	p-value‡
Treatment (n = 93)					.957
Surgery		32 (34.4)	28.9 (13.9)		
Surgery & Radiotherapy		23 (24.7)	27 (10)		
Chemotherapy & Radiotherapy		9 (9.7)	26.7 (3.4)		
Chemotherapy & Radiotherapy & Surgery		26 (28)	28.4 (11.6)		
Radiotherapy		2 (2.2)	31 (18.4)		
Other		1 (1.1)	36 (-)		
Time since Treatment, months (n = 84, missing = 9)	75 (48), 24-165			-.039	.727
Recurrence (n = 90, missing = 3)					.961
No		75 (83.3)	28.5 (11.7)		
Yes		15 (16.7)	28.7 (11)		
Self-administered Comorbidity (possible range = 0-45) (n = 82, missing = 11)	5 (3.6), 0-18			.193	.083
History of depression (n = 93)					.083
Yes		24 (25.8)	31.7 (10.9)		
No		69 (74.2)	27 (11.5)		
History of anxiety (n = 92, missing = 1)					.140
Yes		14 (15.2)	31 (14)		
No		78 (84.8)	27.7 (11.1)		
HADS Depression Score (n = 92, missing = 1)	3.6 (3.3), 0-14			.625	.000
HADS Anxiety Score (n = 92, missing = 1)	5.4 (3.9), 0-18			.676	.000
Social Support Score (n = 92, missing = 1)	28.1 (6.2), 9-34			-.144	.172
Appearance Concerns Score (n = 88, missing = 5)	26.4 (11.1), 11-66			.584	.000

Variable	Mean (SD), Range	Frequency (Valid %)	PCL-C Mean (SD)	Correl ation (r) †	p- value‡
Quality of Life Score (n = 91, missing = 2)	72.2 (22.5), 17-100			-.516	.000
Fear of Recurrence Score (n = 91, missing = 2)	17.3 (8.1), 6-39			.743	.000
Appraisal of Life Threat and Treatment Intensity Score (n = 85, missing = 8)	20.7 (7.5), 7-35			.555	.000

Abbreviations: PTSS, Post-Traumatic Stress Symptoms; PCL-C, Post-Traumatic Stress Disorder Checklist – Civilian Version; N, number of patients; SD, standard deviation

† Pearson product-moment correlation coefficient

‡ Independent sample t-test or Analysis of Variance

Table 2 Prevalence of cancer-related post-traumatic stress in head and neck cancer survivors (N=93)

PTSD Measure†	Frequency (%)	95% Confidence Intervals
PCL-C Score \geq 44	11 (11.8)	6.7 to 20
Symptomatic in 3 Symptom Clusters	12 (12.9)	7.5 to 21.2
Symptomatic in 2 Symptom Clusters	10 (10.8)	6 to 18.7
Symptomatic in 1 Symptom Cluster	21 (22.6)	15.3 to 32.1
Non-symptomatic	50 (53.8)	43.7 to 63.6
Met Cluster B (intrusion) criteria	31 (33.3)	24.6 to 43.4
Met Cluster C (avoidance) criteria	19 (20.4)	13.5 to 29.7
Met Cluster D (hyperarousal) criteria	27 (29)	20.8 to 38.9

Abbreviations: HNC, Head and Neck Cancer; PTSD, Post-Traumatic Stress Disorder

† As measured with the PTSD Checklist – Civilian Version (PCL-C). Symptom clusters include at least one intrusion symptom, three avoidance symptoms and two hyperarousal symptoms. Symptomatic in all three clusters constitutes full PTSD.

Table 4 Multiple Linear Regression for survivors' PCL-C Score and Logistic Regression for Meeting Symptom Criteria for Full or Partial PTSD (N=78)

Variable	PCL-C Score			Full or Partial PTSD†		
	B	95% CI	<i>p</i>	Odds Ratio	95% CI	<i>p</i>
Age at questionnaire completion	-.05	-.18 to .08	.467	.98	.91 to 1.05	.533
HADS Anxiety Score	.53	-.04 to 1.10	.068	1.14	.83 to 1.56	.427
HADS Depression Score	.36	-.33 to 1.05	.304	1.12	.83 to 1.51	.468
Appraisal of life threat and treatment intensity	.15	-.13 to .43	.300	1.02	.87 to 1.21	.790
Fear of Recurrence	.60	.31 to .88	.000	1.18	1.01 to 1.37	.038
Appearance Concerns	.16	-.02 to .34	.074	1.04	.95 to 1.14	.403
Quality of Life	-.08	-.16 to .01	.086	.97	.93 to 1.01	.128

Abbreviations: B, Beta Coefficient; PCL-C, Post-Traumatic Stress Disorder Checklist – Civilian Version; PTSD, Post-Traumatic Stress Disorder; CI, Confidence Intervals

† No. of events/no. of patients = 20/78

Table 3 Prevalence of cancer-related post-traumatic stress in dyads of HNC survivors and their partners (N=39)

Measure	Patient Frequency (%)	Partners Frequency (%)	Test Statistic	p value
PCL Mean Score (SD)	28.6 (10.4)	27.8 (12.5)	t(38) = .35	.731
Symptomatic in 3 Symptom Clusters	6 (15.4)	5 (12.8)	---*	1
Symptomatic in 2 Symptom Clusters	6 (15.4)	4 (10.3)	---*	.754
Symptomatic in 1 Symptom Cluster	7 (17.9)	6 (15.4)	---*	1
Non-symptomatic	20 (51.3)	24 (61.5)		
Met Cluster B (intrusion) criteria	14 (35.9)	13 (33.3)	---*	1
Met Cluster C (avoidance) criteria	10 (25.6)	5 (12.8)	---*	.180
Met Cluster D (hyperarousal) criteria	13 (33.3)	11 (28.2)	---*	.791

* McNemar's test

Fig 1 Consort diagram for study participants