INSTRUCTIONS ON THE ANNOTATION OF PDF FILES

To view, print and annotate your article you will need Adobe Reader version 9 (or higher). This program is freely available for a whole series of platforms that include PC, Mac, and UNIX and can be downloaded from <http://get.adobe.com/reader/>. The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/reader/tech-specs.html>.

Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file.

PDF ANNOTATIONS	
Adobe Reader version 9	Adobe Reader version X and XI
<p>When you open the PDF file using Adobe Reader, the Commenting tool bar should be displayed automatically; if not, click on 'Tools', select 'Comment & Markup', then click on 'Show Comment & Markup tool bar' (or 'Show Commenting bar' on the Mac). If these options are not available in your Adobe Reader menus then it is possible that your Adobe Acrobat version is lower than 9 or the PDF has not been prepared properly.</p> <p>(Mac)</p> <p>PDF ANNOTATIONS (Adobe Reader version 9)</p> <p>The default for the Commenting tool bar is set to 'off' in version 9. To change this setting select 'Edit Preferences', then 'Documents' (at left under 'Categories'), then select the option 'Never' for 'PDF/A View Mode'.</p> <p>(Changing the default setting, Adobe version 9)</p>	<p>To make annotations in the PDF file, open the PDF file using Adobe Reader XI, click on 'Comment'.</p> <p>If this option is not available in your Adobe Reader menus then it is possible that your Adobe Acrobat version is lower than XI or the PDF has not been prepared properly.</p> <p>This opens a task pane and, below that, a list of all Comments in the text. These comments initially show all the changes made by our copyeditor to your file.</p>

HOW TO...		
Action	Adobe Reader version 9	Adobe Reader version X and XI
Insert text	<p>Click the 'Text Edits' button on the Commenting tool bar. Click to set the cursor location in the text and simply start typing. The text will appear in a commenting box. You may also cut-and-paste text from another file into the commenting box. Close the box by clicking on 'x' in the top right-hand corner.</p>	<p>Click the 'Insert Text' icon on the Comment tool bar. Click to set the cursor location in the text and simply start typing. The text will appear in a commenting box. You may also cut-and-paste text from another file into the commenting box. Close the box by clicking on '_' in the top right-hand corner.</p>
Replace text	<p>Click the 'Text Edits' button on the Commenting tool bar. To highlight the text to be replaced, click and drag the cursor over the text. Then simply type in the replacement text. The replacement text will appear in a commenting box. You may also cut-and-paste text from another file into this box. To replace formatted text (an equation for example) please Attach a file (see below).</p>	<p>Click the 'Replace (Ins)' icon on the Comment tool bar. To highlight the text to be replaced, click and drag the cursor over the text. Then simply type in the replacement text. The replacement text will appear in a commenting box. You may also cut-and-paste text from another file into this box. To replace formatted text (an equation for example) please Attach a file (see below).</p>
Remove text	<p>Click the 'Text Edits' button on the Commenting tool bar. Click and drag over the text to be deleted. Then press the delete button on your keyboard. The text to be deleted will then be struck through.</p>	<p>Click the 'Strikethrough (Del)' icon on the Comment tool bar. Click and drag over the text to be deleted. Then press the delete button on your keyboard. The text to be deleted will then be struck through.</p>
Highlight text/ make a comment	<p>Click on the 'Highlight' button on the Commenting tool bar. Click and drag over the text. To make a comment, double click on the highlighted text and simply start typing.</p>	<p>Click on the 'Highlight Text' icon on the Comment tool bar. Click and drag over the text. To make a comment, double click on the highlighted text and simply start typing.</p>
Attach a file	<p>Click on the 'Attach a File' button on the Commenting tool bar. Click on the figure, table or formatted text to be replaced. A window will automatically open allowing you to attach the file. To make a comment, go to 'General' in the 'Properties' window, and then 'Description'. A graphic will appear in the PDF file indicating the insertion of a file.</p>	<p>Click on the 'Attach File' icon on the Comment tool bar. Click on the figure, table or formatted text to be replaced. A window will automatically open allowing you to attach the file. A graphic will appear indicating the insertion of a file.</p>
Leave a note/ comment	<p>Click on the 'Note Tool' button on the Commenting tool bar. Click to set the location of the note on the document and simply start typing. <u>Do not use this feature to make text edits.</u></p>	<p>Click on the 'Add Sticky Note' icon on the Comment tool bar. Click to set the location of the note on the document and simply start typing. <u>Do not use this feature to make text edits.</u></p>

HOW TO...		
Action	Adobe Reader version 9	Adobe Reader version X and XI
Review	To review your changes, click on the 'Show' button on the Commenting tool bar. Choose 'Show Comments List'. Navigate by clicking on a correction in the list. Alternatively, double click on any mark-up to open the commenting box.	Your changes will appear automatically in a list below the Comment tool bar. Navigate by clicking on a correction in the list. Alternatively, double click on any mark-up to open the commenting box.
Undo/delete change	To undo any changes made, use the right click button on your mouse (for PCs, Ctrl-Click for the Mac). Alternatively click on 'Edit' in the main Adobe menu and then 'Undo'. You can also delete edits using the right click (Ctrl-click on the Mac) and selecting 'Delete'.	To undo any changes made, use the right click button on your mouse (for PCs, Ctrl-Click for the Mac). Alternatively click on 'Edit' in the main Adobe menu and then 'Undo'. You can also delete edits using the right click (Ctrl-click on the Mac) and selecting 'Delete'.

SEND YOUR ANNOTATED PDF FILE BACK TO ELSEVIER

Save the annotations to your file and return as instructed by Elsevier. Before returning, please ensure you have answered any questions raised on the Query Form and that you have inserted all corrections: later inclusion of any subsequent corrections cannot be guaranteed.

FURTHER POINTS

- Any (grey) halftones (photographs, micrographs, etc.) are best viewed on screen, for which they are optimized, and your local printer may not be able to output the greys correctly.
- If the PDF files contain colour images, and if you do have a local colour printer available, then it will be likely that you will not be able to correctly reproduce the colours on it, as local variations can occur.
- If you print the PDF file attached, and notice some 'non-standard' output, please check if the problem is also present on screen. If the correct printer driver for your printer is not installed on your PC, the printed output will be distorted.

AUTHOR QUERY FORM

ELSEVIER	Journal: YMOD Article Number: 4936	
--	---	--

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof.

Location in article	Query / Remark: Click on the Q link to find the query's location in text Please insert your reply or correction at the corresponding line in the proof
Q1	<p>If there are any drug dosages in your article, please verify them and indicate that you have done so by initialing this query</p> <p>Is the sentence that begins "Subjects" as you meant?</p> <p>Q2 The first paragraph of "Material and methods" says "4 specialist practices." What is correct?</p> <p>Q3 Is it OK to change the percentage of women so that the total is 100% with the percentage of men added (21.1%)?</p> <p>Q4 Reference 20—is it correct that this was published in 1998?</p> <p>Q5 Please confirm that given names and surnames have been identified correctly.</p> <div data-bbox="303 1415 893 1585"> <p>Please check this box or indicate your approval if you have no corrections to make to the PDF file</p> <input data-bbox="790 1468 877 1553" type="checkbox"/> </div>

Thank you for your assistance.

Highlights

- No one has evaluated factors that predict uptake of orthodontic treatment by adults.
- We assessed the importance of 4 factors in predicting treatment uptake.
- Age, marital status, and oral health—related quality of life predicted uptake.

What factors predict the uptake of orthodontic treatment among adults?

Ama Johal^a and Easter Joury^b
London, United Kingdom

Introduction: Our aim was to evaluate the factors that predict orthodontic treatment uptake among adults attending a specialist practice. **Methods:** A cross-sectional controlled design was adopted in a private practice setting. The test group included 62 adults seeking fixed orthodontic treatment. The controls were 52 parents of children undergoing orthodontics but who had not undergone treatment themselves. All subjects completed a set of validated questionnaires: the Rosenberg self-esteem scale, the shortened version of the oral health impact profile, and the demographic and socioeconomic position characteristics. The dental health component and the aesthetic component of the index of orthodontic treatment need were used to assess the severity of the malocclusions. **Results:** A 100% response rate was achieved. Subjects without a partner ($P < 0.001$) and with a high oral health impact ($P < 0.001$) or with a need for orthodontic treatment (as assessed by the clinician or the subject using the aesthetic component of the index of orthodontic treatment need; $P = 0.003$ and $P = 0.031$, respectively) were more likely to have orthodontic treatment than were their counterparts with a partner (odds ratio [OR] = 20.8; 95% confidence interval [CI] = 4.63-93.25) and with a low oral health impact (OR = 5.3; 95% CI = 2.36-11.88) or with no treatment need (OR = 3.6 and 4.4; 95% CI = 1.57-8.99 and 1.15-16.77, respectively). Self-esteem and demographic and socioeconomic position characteristics were not significantly associated with orthodontic treatment uptake ($P > 0.05$). **Conclusions:** The significance of age, marital status, and the shortened version of the oral health impact profile in predicting the uptake of orthodontic treatment among adults was demonstrated. (Am J Orthod Dentofacial Orthop 2015; ■:■-■)

Enhancing appearance and improving psychosocial function appear to play important roles in an adult's decision to initiate orthodontic treatment.¹⁻³ However, with limited research about adults seeking treatment, it appears that esthetic and functional improvement in occlusion are the 2 most commonly cited reasons by adults for undergoing combined surgical-orthodontic treatment, according to Cunningham et al.⁴ Research in adults has focused primarily on their motivations for seeking treatment, with Pabari et al⁵ reporting that adult motives tend to be numerous and varied, with their psychological traits closer to those among the general population than to

orthognathic patients. Both McKiernan et al⁶ and Sergl and Zentner⁷ used a questionnaire-based study and found that among adults, the primary motivating factor for orthodontic treatment was a desire to improve dental appearance, followed by facial appearance. Sergl and Zentner identified that a functional benefit was also a key motivator for seeking treatment.

Orthodontic treatment uptake has been shown to vary considerably in differing populations, from 5% to 60%, depending on the country.^{8,9} Research to date on the factors that predict orthodontic treatment uptake has focused on adolescent populations and identified a number of factors, including sex,¹⁰ ethnicity,^{11,12} availability of orthodontic services,¹³ socioeconomic status,¹³ and treatment need.¹⁴

Harris and Glassell¹² demonstrated that the greater uptake of orthodontic treatment in girls occurred because of preferential self-selection and not necessarily greater need. The available evidence in relation to socioeconomic factors and treatment uptake shows controversial findings in adolescents. Badran and Al Khateen¹³ reported socioeconomic class as a significant predictor for treatment, with a greater frequency of uptake among adolescents from high and middle socioeconomic classes compared with those from a

From the Department of Oral Growth and Development, Institute of Dentistry, Queen Mary University, London, United Kingdom.

^aReader.

^bVisiting research assistant.

All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest, and none were reported.

Ama Johal and Easter Joury are joint first authors and contributed equally to this work.

Address correspondence to: Ama Johal, Institute of Dentistry, Turner St, Whitechapel, London E1 2AD, England, United Kingdom; e-mail, a.s.johal@qmul.ac.uk.

Submitted, September 2014; revised and accepted, January 2015.

0889-5406/\$36.00

Copyright © 2015 by the American Association of Orthodontists.

<http://dx.doi.org/10.1016/j.ajodo.2015.01.025>

lower class. Breistein and Burden,¹⁵ however, found no evidence to support this association. The uptake of orthodontic treatment may also be influenced by the availability and in turn the ease of access to such services, and it relates strongly to the availability of government-subsidized treatment.^{10,13,16} This serves to highlight the complex interactions and the reasons that the uptake of orthodontic treatment varies across cultural and socioeconomic backgrounds; these have yet to be identified, particularly in adults.

Recently, the focus has been more on patients' own perceptions of oral health status and oral health care systems to understand their needs, satisfaction with treatment, and ultimately the perceived overall quality of health systems in adults.¹⁷ Thus, oral health-related quality of life (OHRQOL), perceived orthodontic treatment need, and self-esteem may play important roles in determining treatment uptake. Feu et al¹⁸ compared OHRQOL in adolescents seeking orthodontic treatment with age-matched peers not seeking treatment and found that those who sought treatment were 3.1 times more likely to have a worse OHRQOL than those in the comparison group. Helm et al¹⁹ attempted to evaluate the influence of the aesthetic component (AC) of the index of treatment need (IOTN) as a motivator for seeking treatment, reporting unfavorable perceptions of teeth and high dental awareness in adults with a malocclusion. Only Mandall et al¹⁴ reported the AC of the IOTN as a predictor of orthodontic treatment, but that study was limited to adolescents. In relation to self-esteem, this has only been evaluated in relation to any observed change as a result of treatment but not as a potential predictor of orthodontic treatment uptake.

Currently, there is no evidence to evaluate the factors that may predict the uptake of orthodontic treatment, especially in light of the greater numbers of adults undergoing treatment. The aim of this study was therefore to evaluate the role of sociodemographic background, orthodontic treatment need, OHRQOL, and self-esteem in terms of their ability to predict orthodontic treatment uptake among adults attending a specialist practice.

MATERIAL AND METHODS

In this study, we used a cross-sectional controlled design, for which ethical approval was obtained from the research ethics committee of Queen Mary University (reference number, QMREC2009/33) in London, United Kingdom. All subjects aged 18 years and above who fulfilled the selection criteria were recruited from 4 specialist practices in southeast England into 2 groups, with written informed consent obtained. The test group

was recruited from adult patients who were due to receive active fixed orthodontic treatment to correct their malocclusion in the specialist practice. The control group was recruited from parents of children undergoing orthodontic treatment in the same settings with no history of orthodontic treatment themselves. Subjects were excluded if they were not literate and fluent in English, or had caries, periodontal disease, recent dental treatment or orthognathic surgery, or a craniofacial deformity. A sample of 114 patients distributed into 2 groups was estimated to be sufficient to demonstrate a 3-fold or greater odds ratio in explanatory variables between the test and control groups with respect to orthodontic treatment uptake, with a power of 80% at a significance level of 5%. The calculation assumed no more than a 70% frequency of exposure to the explanatory variables in the test or the control group.

Explanatory variables included sociodemographic background, orthodontic treatment need, OHRQOL, and self-esteem. Sociodemographic variables included age, sex, ethnicity, marital status, and socioeconomic position indicators: occupation, education, and employment status.²⁰ Occupation is considered an indicator of social class. The Registrar General's Classification of Occupations was used to allocate social class (groups I-V) based on each participant's occupation. These 5 groups are broadly dichotomized into nonmanual (high social class, groups I-III(N) and manual (low social class, groups IIIM-V). In the case of unemployment or retirement, the Registrar General's Classification of Occupations provides no classification. Thus, this information was considered missing. Education was measured by the highest qualification obtained, with high levels indicating university or postgraduate qualifications.²¹ Employment status information included being an employee or self-employed. Marital status included not having a partner (never married, separated, divorced, or widowed) or having a partner (married, remarried, or cohabiting).

Orthodontic treatment need was assessed using the dental health component (DHC) and the AC of the IOTN.²² The DHC and the AC of the IOTN were assessed by trained and calibrated examiners. In addition, each patient's self-perceived dental esthetics was assessed with the AC of the IOTN. The scores of the DHC of the IOTN were dichotomized into 2 categories of need for orthodontic treatment: moderate or lower need, and great or very great need. The scores of the AC of the IOTN were also dichotomized into 2 categories of need for orthodontic treatment: no need (scores 1-4), and borderline or definite need (scores 5-10).²³

Self-esteem was measured using the Rosenberg self-esteem scale.²⁴ It has proven validity and

reliability for general population and orthodontic patients.²⁵ The Rosenberg self-esteem scale consists of 10 questions—5 positive and 5 negative—and uses a Likert scale in which the responses for the positive and negative questions are weighted differently with a 4-point scale, including strongly agree, agree, disagree, and strongly disagree. The scale's total score ranges from 0 to 30. Total scores were dichotomized into high and low levels of self-esteem based on the median values.

The oral health impact profile (OHIP-14) was used to measure the impact of oral health on quality of life.²⁶ The OHIP-14 has 7 conceptualized domains (2 items per domain): functional limitation, physical pain, psychological discomfort, physical disability, psychological disability, social disability, and handicap. The respondents are asked to rate how frequently they experienced an oral health impact (as described by each item). In turn, the response to each item is scored on a 5-point Likert scale: 0, never; 1, hardly ever; 2, occasionally; 3, fairly often; and 4, very often or every day. Thus, the summary of OHIP-14 scores can range from 0 to 56. A higher total sum of the OHIP-14 indicates a greater negative impact on OHRQOL.²⁷ Total scores were dichotomized into high and low levels based on the median values.

A piloted self-completed questionnaire was used to collect the sociodemographic, OHRQOL, and self-esteem data, which demonstrated no problems with the interpretation of the questions; the patients required less than 10 minutes to complete the questionnaire. All patients were advised by the clinicians that they would be available to address any uncertainties about the questions. The need for orthodontic treatment data was collected by a clinical examination.

Statistical analysis

Data analysis was performed with the Statistical Package for the Social Sciences software (version 17.0; SPSS, Chicago, Ill). First, the effect of the explanatory variables on orthodontic treatment uptake was assessed using simple logistic regression analysis. Second, the explanatory variables that were significant at the 0.2 level were selected to enter a regression model.²⁸ This step was intended to confirm the significance and show the collective effect of predictors of orthodontic treatment uptake.

RESULTS

A response rate of 100% was obtained. In total, 114 adults, from 7 specialist practices, were recruited for the study. These included 62 test and 52 control subjects. Men comprised 21.1% of the sample.

Table 1. Frequency of sociodemographic variables, odds ratios, and 95% confidence intervals to predict orthodontic treatment uptake with simple logistic regression analyses (n = 114)

Variable	Base	Frequency of orthodontic treatment uptake	OR (95% CI)	P value
Age (y)				
≥40	76	33 (43.4%)	1	
<40	34	29 (85.3%)	7.6 (2.64–21.63)	<0.001
Missing values	4			
Sex				
Male	24	14 (58.3%)	1	
Female	90	48 (53.3%)	0.8 (0.33–2.03)	0.662
Missing values	0			
Ethnicity				
White	101	54 (53.5%)	1	
Other	13	8 (61.5%)	1.4 (0.43–4.55)	0.583
Missing values	0			
Marital status				
With a partner	82	33 (40.2)	1	
Without a partner	30	28 (93.3)	20.8 (4.63–93.25)	<0.001
Missing values	2			
Social class				
High	94	47 (50%)	1	
Low	14	11 (71.4%)	2.5 (0.73–8.54)	0.144
Missing values	6			
Education				
High	39	23 (59%)	1	
Low	74	38 (51.4%)	0.7 (0.34–1.61)	0.440
Missing values	1			
Employment				
Self-employed	20	8 (40%)	1	
Employee	88	49 (55.7%)	1.9 (0.70–5.07)	0.209
Missing values	6			

The difference in orthodontic treatment uptake between subjects aged less than 40 years (85.3%) and those aged 40 years and above (43.4%) was large and statistically significant ($P < 0.001$; Table 1). Subjects less than 40 years old were more likely to have orthodontic treatment than were their counterparts over 40 years of age (odds ratio [OR] = 7.6; 95% confidence interval [CI] = 2.64–21.63; Table 1).

With respect to marital status, the difference in orthodontic treatment uptake between subjects with a partner (93.3%) and those without a partner (40.2%) was also large and statistically significant ($P < 0.001$; Table 1). Subjects without a partner were more likely to have orthodontic treatment than were their counterparts with a partner (OR = 20.8; 95% CI = 4.63–93.25; Table 1). Among the other sociodemographic variables—sex, ethnicity, social class, employment, and education—only social class was significant at the 0.2 level (Table 1).

Table II. Frequency of need for orthodontic treatment, OHRQOL, and self-esteem variables, odds ratios, and 95% confidence intervals to predict orthodontic treatment uptake with simple logistic regression analyses (n = 114)

Variable	Base	Frequency of orthodontic treatment uptake	OR (95% CI)	P value
DHC of the IOTN				
Moderate and below	64	26 (40.6%)	1	
Great and very great	34	20 (58.8%)	2.1 (0.90-4.86)	0.088
Missing values	16			
C-AC of the IOTN*				
No need	59	21 (35.6%)	1	
Need	42	28 (66.7%)	3.6 (1.57-8.33)	0.003
Missing values	13			
S-AC of the IOTN†				
No need	90	41 (45.6%)	1	
Need	14	11 (78.6%)	4.4 (1.15-16.77)	0.031
Missing values	10			
OHRQOL				
Low impact	59	21 (35.6%)	1	
High impact	55	41 (74.5%)	5.3 (2.36-11.88)	<0.001
Missing values	0			
Self-esteem				
Low	60	33 (55%)	1	
High	54	29 (53.7%)	0.9 (0.45-1.99)	0.890
Missing values	0			

*AC of the IOTN scored by the clinician; †AC of the IOTN scored by the subject.

With respect to orthodontic treatment need, the difference in orthodontic treatment uptake between subjects with a need (as assessed by the clinician or the subject using the AC of the IOTN) and those without a need was statistically significant ($P = 0.003$ and $P = 0.031$, respectively; Table II). Subjects with a need for orthodontic treatment were more likely to have it than were their counterparts without a need (OR = 3.6 and 4.4; 95% CI = 1.57-8.99 and 1.15-16.77, respectively; Table II). The difference in orthodontic treatment need as assessed by the DHC of the IOTN was significant at the 0.2 level ($P = 0.088$; Table II).

With regard to the OHRQOL, the difference in orthodontic treatment uptake between subjects with a high oral health impact (74.5%) and those with a low oral health impact (35.6%) was large and statistically significant ($P < 0.001$; Table II). Subjects who had a high oral health impact were more likely to have orthodontic treatment than were their counterparts who had a low oral health impact (OR = 5.3; 95% CI = 2.36-11.88; Table II).

The difference between subjects with high and low levels of self-esteem was small and not statistically significant at the 0.2 level (Table II).

A regression model was used to confirm the significance of the identified predictors of orthodontic treatment uptake and show their collective effects. The correlation matrix of the significant variables at the 0.2 level showed significant associations between the 3 measures of orthodontic treatment need as well as a significant association between the AC of the IOTN as scored by the clinician and the OHIP-14. To prevent colinearity, only the most significant orthodontic treatment need variable that was not significantly correlated with the other significant variables was selected to enter the regression model: the AC of the IOTN as scored by the subject. The regression model confirmed the significance of age, marital status, and OHRQOL in predicting the uptake of orthodontic treatment (Table III).

DISCUSSION

This cross-sectional controlled design study is the first to date to evaluate the factors that predict orthodontic treatment uptake among adults attending specialist practices. The relevance of this form of research is justified by the observed increases in adult patients in clinical practices.^{29,30} McKiernan et al⁶ reported that this increased acceptance of treatment by adults reflects an increased overall awareness of their appearance, the social acceptability of treatment, the improved appearance of appliances, and the increasing dissatisfaction with the previous treatment interventions.

Thus, an understanding of the factors that predict uptake and the possibility of identifying them would add to the process of informed consent and ultimately patient satisfaction with treatment. To date, no study has explored the role of OHRQOL on the uptake of orthodontic treatment.

The ratios of men to women in the 2 groups were similar, but most participants were women (78.9%). Previous research has shown that female subjects are more dissatisfied with the appearance of their dentition and perceive a need for braces more often than do male subjects.^{4,12} A clear statistical difference was detected in terms of the influence of the ages of the adults in the uptake of treatment; it was greatest in subjects less than 40 years old.⁴ This is consistent with the findings of previous studies, with reported mean ages of 34 and 25 years.^{5,6} Furthermore, with respect to marital status, the difference in orthodontic treatment uptake between subjects with a partner and those without a partner was also large and statistically significant. These factors may

Table III. Frequency distribution, unadjusted odds ratios, adjusted odds ratios, and 95% confidence intervals to predict orthodontic treatment uptake with a multiple logistic regression analysis (n = 114)

Variable	Base	Frequency of orthodontic treatment uptake	Unadjusted OR (95% CI)	P value	Adjusted OR (95% CI)	P value
Age (y)						
<40	76	33 (43.4%)	1		1	
≥40	34	29 (85.3%)	7.6 (2.64–21.63)	<0.001	4.2 (1.21–14.88)	0.024
Marital status						
With a partner	82	33 (40.2)	1		1	
Without a partner	30	28 (93.3)	20.8 (4.63–93.25)	<0.001	9.9 (1.93–50.60)	0.006
Social class						
High	94	47 (50%)	1		1	
Low	14	11 (71.4%)	2.5 (0.73–8.54)	0.144	1.9 (0.40–8.72)	0.430
S-AC of the IOTN*						
No need	90	41 (45.6%)	1		1	
Need	14	11 (78.6%)	4.4 (1.15–16.77)	0.031	3.2 (0.48–21.4)	0.229
OHRQOL						
Low impact	59	21 (35.6%)	1		1	
High impact	55	41 (74.5%)	5.3 (2.36–11.88)	<0.001	3.1 (1.11–8.43)	0.030

*AC of the IOTN scored by the subject.

reflect the fact that physical attractiveness plays a major role in social interactions; particularly for those who are young and single, this has increased importance. From infancy to adulthood, beautiful people are treated preferentially and viewed more positively; they are judged to be more socially skilled, to be more desirable as friends, to have higher intellectual abilities, to be more successful occupationally, and to be less guilty of crime.^{31,32} Pabari et al⁵ demonstrated, using a qualitative methodology, that a desire to improve the smile and straighten the teeth was the primary motivating factor for adults seeking treatment.

With respect to orthodontic treatment need, the difference in treatment uptake between subjects with a need, as assessed by both the clinician and the subject using the AC of the IOTN, and those without a need was statistically significant. Most participants seeking orthodontic treatment were aware of their malocclusion, as indicated by their scores on the AC of the IOTN, which indicated some esthetic impairment.¹⁸ This agrees with the results of Helm et al,¹⁹ who found unfavorable perceptions of teeth and high dental awareness in adults with malocclusion. Again, this adds weight to previous studies that have focused on evaluating the motives of adults seeking treatment.^{5,6} Studies evaluating treatment uptake in adolescents showed that those who sought treatment had more severe malocclusions and greater esthetic impacts, as judged by both the clinicians and the patients.^{14,18}

Self-esteem did not appear to influence the uptake of orthodontic treatment of adults. No previous study has evaluated the role of self-esteem in orthodontic

treatment uptake in any age group. This lack of significance observed in our study might be interpreted in view of other studies that evaluated the impact of orthodontic treatment and malocclusion on self-esteem. For example, several studies have reported that a dentofacial deformity did not necessarily imply lower self-esteem.^{33,34} Thus, self-esteem might not be related to the perceived need for orthodontic treatment; therefore, it might not be related to the uptake of orthodontic treatment.

With regard to the OHRQOL, the difference in orthodontic treatment uptake between subjects with a high oral health impact and those with a low oral health impact was large and statistically significant. Similar findings were reported in a study of adolescents seeking orthodontic treatment; those who sought treatment had greater negative OHRQOL impacts.¹⁸ The authors applied the same instrument of the OHRQOL measure, the OHIP-14, which was originally developed for use in adult populations, but they reported that it had successfully been applied to adolescents in some studies.

The multivariate analysis showed the significance of age, marital status, and OHRQOL in predicting the uptake of orthodontic treatment. Although direct comparison with the literature is not possible, research among adolescent patients shows that self-perceived esthetics, as assessed by the AC of the IOTN, appears to be a strong predictor of negative OHRQOL impacts, which in turn influence the decision to have treatment.^{14,18,35} Thus, it appears that the uptake of orthodontic treatment by adults is influenced by various factors of which clinicians should be aware when assessing these

persons and obtaining informed consent, to achieve the best outcome for their patients.

This study is not without its limitations. In terms of sample selection, the control group was recruited from parents with no previous history of orthodontic treatment; they attended with their children who were being treated, since the data were collected in the specialist practices, and these parents provided a convenient sample. The use of a cross-sectional design limited data collection to 1 time point; therefore, no temporal relationship could be established between the independent variables such as the OHRQOL and orthodontic treatment uptake. Whereas a cohort study design allows exploration of such a temporal relationship, it requires preliminary evidence of significant associations from cross-sectional studies before adopting such an expensive and time-consuming study design. Our sample size was another potential limitation. The large confidence intervals of the significant independent variables suggest that the sample size was small. Yet, no previous study could help in calculating the sample size for our study. A further potential limitation relates to missing data in some sociodemographic and orthodontic treatment need variables. However, the amounts of missing data were too small to introduce any potential bias into our findings. To further validate such an assumption, a separate group was constituted of subjects with missing data (in each of the sociodemographic and orthodontic treatment need variables with missing data). The statistically significant differences were checked between each group of subjects with missing data and the group of subjects with complete data using logistic regression analyses. No significant differences were identified, suggesting that the small amount of missing data was unlikely to introduce any bias into these findings.

CONCLUSIONS

The significance of age, marital status, and OHRQOL in predicting the uptake of orthodontic treatment among adults was demonstrated.

REFERENCES

- Byrne D, London O, Reeves K. The effects of physical attractiveness, sex, and attitude similarity on interpersonal attraction. *J Pers* 1968;36:259-71.
- Greenwald A, Banaji M. Implicit social cognition: attitudes, self-esteem and stereotypes. *Psychol Rev* 1995;102:4-27.
- Katz R. Relationships between eight orthodontic indices and an oral self-image satisfaction scale. *Am J Orthod* 1978;73:328-33.
- Cunningham S, Hunt N, Feinmann C. Psychological aspects of orthognathic surgery: a review of literature. *Int J Adult Orthod Orthognath Surg* 1995;10:159-64.
- Pabari S, Moles D, Cunningham S. Assessment of motivation and psychological characteristics of adult orthodontic patients. *Am J Orthod Dentofacial Orthop* 2011;140:e263-72.
- McKiernan E, McKiernan F, Jones M. Psychological profiles and motives of adults seeking orthodontic treatment. *Int J Adult Orthod Orthognath Surg* 1992;7:187-98.
- Serfl HG, Zentner A. A study of psychological aspects of adult orthodontic treatment. *Int J Adult Orthognath Surg* 1997;12:17-22.
- Foster Page LA, Thomson WM. Malocclusion and uptake of orthodontic treatment in Taranaki 12-13-year-olds. *N Z Dent J* 2005;101:98-105.
- Nobile C, Pavia M, Fortunato L, Angelillo I. Prevalence and factors related to malocclusion and orthodontic treatment need in children and adolescents in Italy. *Eur J Public Health* 2007;17:637-41.
- Kerosuo H, Abdulkarim E, Kerosuo E. Subjective need and orthodontic treatment experience in a Middle East country providing free orthodontic services: a questionnaire survey. *Angle Orthod* 2002;72:565-70.
- Proffit WR, Fields HW Jr, Moray LJ. Prevalence of malocclusion and orthodontic treatment need in the United States: estimates from the NHANES III survey. *Int J Adult Orthod Orthognath Surg* 1998;13:97-106.
- Harris EF, Glassell BE. Sex differences in the uptake of orthodontic services among adolescents in the United States. *Am J Orthod Dentofacial Orthop* 2011;140:543-9.
- Badran SA, Al Khateen S. Factors influencing the uptake of orthodontic treatment. *J Public Health Dent* 2013;73:339-44.
- Mandall NA, Wright J, Conboy F, Kay E, Harvey L, O'Brien KD. Index of orthodontic treatment need as a predictor of orthodontic treatment uptake. *Am J Orthod Dentofacial Orthop* 2005;128:703-7.
- Breistein B, Burden DJ. Equity and orthodontic treatment: a study among adolescents in Northern Ireland. *Am J Orthod Dentofacial Orthop* 1998;113:408-13.
- Drugan CS, Hamilton S, Naqvi H, Boyles JR. Inequality in uptake of orthodontic services. *Br Dent J* 2007;202:15-22.
- de Oliveira C, Sheiham A. Orthodontic treatment and its impact on oral health-related quality of life in Brazilian adolescents. *J Orthod* 2004;31:20-7.
- Feu D, de Oliveira BH, de Oliveira Almeida MA, Kiyak HA, Miguel JA. Oral health related quality of life and orthodontic treatment seeking. *Am J Orthod Dentofacial Orthop* 2010;138:152-9.
- Helm S, Kreiborg S, Solow B. Psychosocial implications of malocclusion: a 15-year follow-up study in 30-year-old Danes. *Am J Orthod* 1985;87:110-8.
- Adult Dental Health Survey. Oral Health in the United Kingdom 1998, Office for National Statistics. London: United Kingdom: Stationary Office; 1998.
- Liberatos P, Link B, Kelsey J. The measurement of social class in epidemiology. *Epidemiol Rev* 1988;10:87-121.
- Brooke P, Shaw W. Development of an index of orthodontic treatment priority. *Eur J Orthod* 1989;11:309-20.
- Richmond S, Shaw W, O'Brien K, Buchanan I, Stephens C, Andrews M, et al. The relationship between the index of orthodontic treatment need and consensus opinion of a panel of 74 dentists. *Br Dent J* 1995;178:370-4.
- Rosenberg M. Society and the adolescent self-image. Middletown, Conn: Wesleyan University Press; 1965.
- Shaw W, Richmond S, Kenealy P, Kingdon A, Worthington H. A 20-year cohort study of health gain from orthodontic treatment: psychological outcome. *Am J Orthod Dentofacial Orthop* 2007;132:146-57.

- 721 26. Slade G. Derivation and validation of a short-form oral health
722 impact profile. *Community Dent Oral Epidemiol* 1997;25:284-90. 781
- 723 27. Slade G, Spencer A. Development and evaluation of the oral health
724 impact profile. *Community Dent Health* 1994;11:3-11. 782
- 725 28. Altman D. *Practical statistics for medical research*. London, United
726 Kingdom: Chapman and Hall; 1991. 783
- 727 29. Khan R, Horrocks E. A study of adult orthodontic patients and their
728 treatment. *Br J Orthod* 1991;18:183-94. 784
- 729 30. Nattrass C, Sandy J. Adult orthodontics—a review. *Br J Orthod*
730 1995;22:331-7. 785
- 731 31. Baldwin D. Appearance and aesthetics in oral health. *Community*
732 *Dent Oral Epidemiol* 1980;8:244-56. 786
- 733 787
- 734 788
- 735 789
- 736 790
- 737 791
- 738 792
- 739 793
- 740 794
- 741 795
- 742 796
- 743 797
- 744 798
- 745 799
- 746 800
- 747 801
- 748 802
- 749 803
- 750 804
- 751 805
- 752 806
- 753 807
- 754 808
- 755 809
- 756 810
- 757 811
- 758 812
- 759 813
- 760 814
- 761 815
- 762 816
- 763 817
- 764 818
- 765 819
- 766 820
- 767 821
- 768 822
- 769 823
- 770 824
- 771 825
- 772 826
- 773 827
- 774 828
- 775 829
- 776 830
- 777 831
- 778 832
- 779 833
- 780 834
- 835
- 836
- 837
- 838
- 839
- 840
32. Jacobsen A. Psychological aspects of dentofacial esthetics and
orthognathic surgery. *Angle Orthod* 1984;54:18-35.
33. O'Regan J, Dewey M, Slade P, Lovious B. Self-esteem and aes-
thetics. *Br J Orthod* 1991;18:111-8.
34. Varela M, Garcia-Camba J. Impact of orthodontics on the psycho-
logic profile of adult patients: a prospective study. *Am J Orthod*
Dentofacial Orthod 1995;108:142-8.
35. Kok YV, Mageson P, Harradine NW, Sprod AJ. Comparing a
quality of life measure and the aesthetic component of the
index of orthodontic treatment need (IOTN) in assessing
orthodontic treatment need and concern. *J Orthod* 2004;31:
312-8.