Heart Failure Drug Modifies the Intrinsic Dynamics of the Pre-Power Stroke State of Cardiac Myosin

Shaima Hashem,^{a†} William George Davies^a and Arianna Fornili^{a,b*}

^a School of Biological and Chemical Sciences, Queen Mary University of London, London, E1 4NS, United Kingdom

^b The Thomas Young Centre for Theory and Simulation of Materials, London, WC1E 6BN, United Kingdom

Abstract

Omecamtiv mecarbil (OM), currently investigated for the treatment of heart failure, is the first example of a new class of drugs (cardiac myotropes) that can modify muscle contractility by directly targeting sarcomeric proteins. Using atomistic molecular dynamics simulations, we show that the binding of OM to the pre-power stroke state of cardiac myosin inhibits the functional motions of the protein and potentially affects Pi release from the nucleotide binding site. We also show that the changes in myosin ATPase activity induced by a set of OM analogues can be predicted from their relative affinity to the pre-power stroke state compared to the near rigor one, indicating that conformational selectivity plays an important role in determining the activity of these compounds.

Introduction

Omecamtiv Mecarbil (OM) is a first-in-class cardiac myotrope in clinical trial for the treatment of heart failure.¹⁻⁴ Differently from the currently available heart failure drugs, OM increases heart contractility by directly and selectively binding cardiac myosin, the motor protein responsible for the mechanical work at the basis of muscle contraction. Myotropes are considered as a new promising class of drugs for skeletal and cardiac disease since they can affect muscle contractility without altering Ca²⁺-dependent pathways, thus reducing the probability and extent of Ca²⁺-related side effects.^{5,6}

OM has been shown to increase the fraction of time spent by myosin heads in the state where they are strongly bound to actin (duty ratio), which is related to the observed increase in the rate of actin-activated Pi-release and in force production as measured by fractional shortening of cardiac myocytes.² While these findings support the view that OM is a myosin activator, other observations point to a more complex type of mechanism. Indeed, different in vitro studies have shown that OM decreases the rate of Pi release in the absence of actin², reduces the sliding velocity of actin filaments in motility assays⁷ and slows down⁸ or even suppresses⁹ the power stroke, the force-generating motion of the myosin heads when bound to actin. These features seem more typical of an inhibitor rather than an activator.

Previous Molecular Dynamics (MD) studies by our group highlighted that OM binding induces a reduction in the motions of the motor domain of cardiac myosin¹⁰, which involved in particular the converter and lever arm subdomains of the protein (here collectively named as CLD). Because of the limited availability of experimental structures of OM-bound myosin at that time, the study was performed on the near rigor (NR) state¹¹, where the CLD is in a 'down' conformation (Figure 1B). However, evidence indicates that a more relevant state for the OM mechanism of action is the pre-power stroke (PPS) one, where the CLD is rotated upwards (Figure 1A) and myosin is ready to bind actin¹². Weak actin binding of myosin in the PPS state is followed by the steps that have been shown

to be affected by OM, namely the transition to strongly bound states accompanied by Pi release and then the power stroke, where the CLD rotates back to the down position.

In the following, we present the first computational study of the effects of OM on the dynamics of the pre-power stroke state of myosin and of the interactions that are responsible for them, aimed at providing a detailed atomistic model to be used in the future for the design of new compounds with modified activity profiles. Describing protein dynamics has been indeed repeatedly shown to be essential when investigating protein function and its modulation by mutations and small molecules, in particular for sarcomeric proteins (see Ref. ^{13–18} for recent examples). We also extend our investigation to a set of OM analogues, in order to characterise the relationship between their binding interactions with myosin and their effect on the protein function.

Methods

System preparation

Molecular Dynamics (MD) simulations were carried out on the human β -cardiac myosin motor domain (residues 1 to 783), starting from X-ray structures of bovine β -cardiac myosin in the PPS state with (PDB ID: 5N69) and without (PDB ID: 5N6A) OM¹². The structure for the human sequence (96% sequence identity with bovine for residues 1-783) was generated with MODELLER using the bovine X-ray structures and, for missing residues, NR X-ray (PDB ID: 4PA0)¹¹ and MD¹⁰ structures. Both the Apo and OM-bound systems were simulated with Mg^{2+} , ADP and Pi in the nucleotide binding site, using coordinates from the PPS X-ray structures (vanadate in the OM-bound structure was first converted into phosphate). Pi was treated as doubly protonated ($H_2PO_4^-$), following previous studies of myosin bound to $MgADP.Pi$ ¹⁹. Distance restraints were added between N6 of ADP and O of Y134 to prevent large rearrangements of ADP in its binding site (piecewise linear/harmonic restraint, with an upper bound of 4.5 Å and a force constant of 12 kcal/mol/Å²). In preliminary MD simulations of the Apo system, the N-terminal segment (residues ~1-20) of the N-terminal domain was too flexible and it tended to unfold. Distance restraints (force constant = 12 kcal/mol/Å²) were thus applied between C α atoms of residues S4 and E22 (upper bound = 17 Å) and V8 and E137 (upper bound = 9 Å) and between C δ of E22 and O of Y142 (upper bound = 4 Å).

MD simulation protocol

The protein was described with the Amber99SB*-ILDN force field²⁰ and OM with GAFF²¹ parameters as described in Ref. 10. ADP and Pi were described using parameters from Ref. 22. AmberTools²³ and Acpye²⁴ were used to convert topology and parameter files to the GROMACS format. The system was solvated using a truncated octahedral box of TIP3P water molecules, using a minimal distance of 12 Å between the protein and the walls of the box. The charge of the ionisable residues was set to that of their standard protonation state at pH 7. The systems were neutralised by adding counter-ions, for a total of ~ 208,000 atoms. The simulations were run adopting the same protocol and parameters used for the NR state¹⁰. The Apo system required longer equilibrations, so that the length of each of the equilibration steps was doubled. Production simulations were then run for 300 ns, saving the coordinates every 1 ps. Four replicas were performed for each state to check for reproducibility of the results, for a total simulation time (production only) of 2.4 μ s (2 X 4 X 300ns).

Analysis of MD trajectories

Dynamic cross correlation matrices (DCCM) and Mutual Information (MI) networks from the M32K25 Structural Alphabet (SA) were calculated as described in Ref. 10. In particular, regions preferentially connected to a source residue a were determined by calculating the shortest distances in the MI network between each of the 4-residue fragments j_a containing a and all the protein fragments. The distances were then standardised by calculating their Z-scores ζ_{ja} and an overall ζ profile was derived by calculating the minimum over the ζ_{ja} profiles. Fragments with negative ζ values are closer to the source site than the average, so that they can be considered as preferentially connected to it. A ζ profile was calculated from each of the two consensus MI matrices calculated over the 4 replicas for each system (PPS Apo and OM-bound) and the $\Delta\zeta$ differences were then calculated by subtracting the Apo profile from the OM-bound one.

A Principal Component Analysis (PCA) was performed on the pseudo-trajectory obtained by concatenating together all the PPS and NR simulations (16 X 300-ns production trajectories). The PCA was performed with GROMACS tools using Ca atoms coordinates extracted every 100 ps.

In order to reduce the noise in the analyses, high-flexibility regions including or close to parts of the structure missing in the original PDB files (residues 1-23, 198-218, 296-300, 367-371, 404-411, 568-573, 621-647, 723-742 and 781-783) were excluded from DCCM, MI, PCA and RMSD calculations.

Hydrogen bond (HB) networks were analysed to identify OM-induced changes in inter-residue interactions around the OM binding site. The frequency of occurrence (or occupancy) of HB interactions was calculated for all residue pairs within 10 Å of the OM binding site. HB interaction matrices were calculated for each of the PPS and NR trajectories with the `g_hbond` tool from GROMACS, using a cutoff of 30° on the H-Donor-Acceptor angle and of 3.5 Å on the Donor-Acceptor distance. Average matrices (avgHB) were then calculated over the four replicas for each conformation (PPS and NR) and binding state (Apo and OM-bound). Difference matrices (Δ HBFreq) were finally calculated as:

$$\Delta\text{HBFreq(PPS)} = \text{AvgHB(OMbound_PPS)} - \text{AvgHB(Apo_PPS)}$$

$$\Delta\text{HBFreq(NR)} = \text{AvgHB(OMbound_NR)} - \text{AvgHB(Apo_NR)}$$

Only pairs with $|\Delta\text{HBFreq}| \geq 0.4$ were considered. A positive ΔHBFreq value indicates that the corresponding hydrogen bond is strengthened by OM binding.

MDpocket²⁵ (stand-alone version) was used to detect the changes in the protein cavities during MD simulations. This method is based on the repeated application of fpocket²⁶ on snapshots extracted from a MD trajectory. In fpocket, cavities are detected as sets of α -spheres, i.e. spheres in contact with four atoms and with no atom inside. MDpocket then determines the frequency of occurrence of cavities during the simulation for each point of a 1-Å spaced grid, which can be visualised as a frequency map. MDpocket analyses were performed on each PPS Apo and OM-bound trajectory containing protein structures sampled every 5 ps, using the default fpocket parameters for α -sphere detection. A common frame (initial structure) was added as the first frame of each trajectory in order to have the same grid definition for all the MDpocket analyses. All frames in each trajectory were aligned to the first one by best fit superimposition.

Representative structures of OM-bound trajectories were used for the analysis of the OM-myosin interactions (Figure 1 and Table S1) and for the docking of OM analogues (see below). They were determined by cluster analysis performed with the 'gromos'²⁷ clustering method implemented in

GROMACS, using a cutoff of 2 Å on structures sampled every 100 ps. The four OM-bound trajectories were first concatenated. Each structure in the concatenated trajectory was then fitted to the first one using all the C α atoms in the protein. The RMSD values used for the clustering were then calculated on all OM non-hydrogen atoms. A dominant cluster was found, with similar population in the four replicas (Table S2). The centroid of this cluster was used as representative structure for the PPS state and compared with the NR representative derived in the same way¹⁰.

Contact analysis in the nucleotide binding site was performed using R and the bio3D library²⁸. All the residues in the outer and inner backdoor I tube as indicated in Ref. 19 were included. Two residues were considered to be in contact if the minimum distance calculated over all the pairs of non-hydrogen side chain atoms was < 4 Å. The analysis was performed on each PPS Apo and OM-bound trajectory with structures sampled every 100 ps.

Molecular Docking

OM analogues were docked to PPS and NR representative structures extracted from the OM-bound MD simulations by clustering (see above). The analogues were selected among those found in Refs. 29 and 30 as having a lower (group 1), similar (group 2) or higher (group 3) % increase in myosin ATPase activity compared to OM, measured for each compound with a sarcomeric assay at 10 mM. Initial ligand 3D-structures were downloaded from PubChem³¹ or, when not available, generated with the Online SMILES Translator and Structure File Generator Translator web interface (<https://cactus.nci.nih.gov/translate/>) of the CACTVS system.

AutoDock Tools³² were used to prepare the systems and generate the pdbqt files for subsequent docking. In particular, non-polar hydrogen atoms were merged with the corresponding heavy atoms and ligand amide bonds were set as rotatable. A 20X20X20 Å³ grid box was defined with 1-Å spacing and centred on the OM binding site. Docking was performed with AutoDock Vina³³, using an exhaustiveness of 3,000 and a maximum number of 9 binding poses. For each ligand, five docking calculations were run using different random seeds and the run leading to the lowest binding affinity was selected. The steric, hydrophobic and hydrogen-bond components of the binding affinity were calculated by re-running vina with the --score_only option for each binding pose. Control calculations were performed where OM was re-docked to both myosin conformations using the same parameters described above. Vina found the correct binding poses (RMSD from target < 2 Å) in both cases, which were ranked as best solution (lowest predicted binding affinity) for PPS (RMSD=0.78 Å) and as second best solution for NR (RMSD=1.40 Å). The corresponding binding affinities (-7.6 kcal/mol for NR and -9.7 kcal/mol for PPS) confirm that OM binds more strongly to the PPS state compared to the NR one by ~2 kcal/mol, as expected on the basis of the available experimental data (K_d values measured on cardiac myosin bound to different nucleotides)¹².

Docking results were analysed using PyMOL³⁴ and in-house R scripts. Linear regression models were built using the lm() R function. Absolute binding affinities and NR - PPS binding affinity differences were first independently rescaled as Z-scores. Different linear models (Models 1 to 5) were then built by fitting different linear combinations of scores to the % increase in myosin ATPase activity. The individual scores were calculated as PPS binding affinity Z-scores + NR - PPS binding affinity difference Z-scores (Models 1, 2 and 5), PPS binding affinity Z-scores (Model 3) and NR binding affinity Z-scores (Model 4). Averages calculated over all the Vina binding poses were used for Models 1-4, while only the best binding pose (lowest binding affinity) was used for Model 5. The best

combination was then identified on the basis of the adjusted R-squared value and the statistical significance of the linear combination coefficients.

Results

OM interacts more strongly with the PPS state

OM-myosin interactions were analysed on representative structures extracted from MD simulations using a cluster analysis. For each state (PPS and NR), the centroid of the most populated cluster was considered (Table S2). The strength of the interactions was also evaluated by calculating their frequency of occurrence over the whole trajectories (Methods). Results were compared with previous simulations of the OM-bound NR state¹⁰.

OM binds the PPS conformation in a deep pocket between the N-terminal and CLD sub-domains of the myosin motor domain

(Figure 1A and C). The ligand is stably bound to the binding site (Figure S1A) and its structure remains close to the starting one throughout the simulations (Figure S1B), with the RMSD calculated over non-hydrogen atoms below 2.0 Å for almost 80% of all the frames in the four replicas.

Comparison with the NR complex (Figure 1B and D) shows that, apart from a small subset of residues (bold in Table S1), the residue composition of the NR and PPS pocket is different, reflecting the large conformational difference between the two states in the CLD region. The orientation of the molecule is also different: in the NR state the carboxymethyl piperazine end (a in Figure 1D) is buried deep in the pocket and close to the relay helix (cyan), while in the PPS state (Figure 1C) the molecule is flipped, with the d end (methylpyridin-3-yl ring) closest to the relay helix.

A common element in the OM-myosin interactions formed in the two states is the involvement of the urea group (c) in hydrogen bonds with nearby residues. These bonds are particularly strong in the PPS state, where this moiety forms multiple high-occupancy interactions with D168, N711 and R712 (average occupancy > 0.5, Table S3). This could explain why derivatives of OM need to retain this functional group to preserve activity^{29,30}. No hydrogen bonds of comparable strength were found among the OM-myosin interactions in the NR state (Table S3). The PPS state contains also a larger number of stable OM-myosin contacts, with 11 residues found in contact with OM through their side chains for more than 70% of the simulations (average in Table S4) compared to the 7 residues observed for the NR state. In particular, stable contacts are formed in the PPS state with the aromatic or hydrophobic side chains of Y164, L770 and I713, while they are absent or less stable in the NR simulations.

Figure 1. OM binding site in PPS and NR structures from MD. A-B. Cartoon representation of PPS (A) and NR (B) structures of the motor domain bound to OM (purple sticks) extracted from MD simulations. Different colours are used for the N-terminal (green), U50K (red), L50K (grey), relay helix (cyan) and CLD (blue) sub-domains. C-D. Close-up view of the OM binding site in the PPS (C) and NR (D) structures. Atoms within 4 Å from OM are represented as white surface, the corresponding residues as sticks and OM-myosin hydrogen bonds as yellow dashed lines. Hydrogen-bond forming residues are labelled.

The analysis of the simulations indicates that the presence of OM increases the coupling between the subdomains in the region around the binding site also through direct hydrogen bonding interactions between the residues and that this effect is more pronounced for the PPS state (Figures S2 and S3). Indeed, OM binding to the PPS state is found to cause larger variations in the frequency of occurrence of hydrogen bonds (red in Figure S2) than the NR one (blue). The OM-induced stabilisation of the hydrogen bonds ($\Delta\text{HBfreq} > 0$) is more extensive in the PPS state and leads on average to stronger hydrogen bonds, in particular those within the relay helix (intra RH in Table S5) and those coupling the CLD with the N-terminal subdomain (N-term-CLD) or the relay helix (RH-CLD).

Figure 2. Linear regression model of the change in the ATPase activity of cardiac myosin induced by OM analogues. The experimental change measured for group 1 (cyan), group 2 (blue) and group 3 (orange) compounds is plotted against the change predicted using PPS vs NR binding affinity differences (Model 1 in Table S7).

Conformational selectivity modulates the activity of OM analogues

Ligand-protein interactions in the OM binding site were further investigated by modelling the complexes between OM analogues with known activity and the two conformational states of myosin. Analogues were selected to sample different levels of activity while keeping their chemical structure as similar as possible to OM (Figure S4). The selected compounds, which cause a lower (group 1), similar (group 2) or higher (group 3) increase in cardiac myosin ATPase activity compared to OM^{29,30}, were docked to the binding site and the resulting interactions analysed (Methods). Control calculations were also performed where OM was re-docked to both myosin conformations. The docking procedure identified the correct binding poses (RMSD from target < 2 Å) in both cases (Supplementary Methods). The corresponding binding affinities (-7.6 kcal/mol for NR and -9.7 kcal/mol for PPS) confirm that OM binds more strongly to the PPS state compared to the NR, as expected on the basis of the available experimental data¹² and our MD simulations.

Consistently with the behaviour observed with OM, all the analogues were found to have higher affinity to the PPS conformational state than the NR one (Table S6). However, the affinity difference was found to be more pronounced for groups 2 and 3 (1.3 kcal/mol on average) compared to group 1 (0.8 kcal/mol on average). If the single contributions to the affinity are considered, groups 2 and 3 tend to have larger steric and hydrogen bonded terms (both absolute values and differences) compared to group 1. The opposite trend was observed for the hydrophobic terms, which tend to be more important in group 1 compared to the other two. Correspondingly, a comparison of the myosin-ligand complexes showed a different distribution in the ligand-protein polar contacts for the different groups in particular in the PPS state (Figure S5), where group 1 ligands form interactions mostly with the N-terminal domain, while groups 2 and 3 form a more extensive network of interactions that cross-link the N-terminal domain, the CLD and the relay helix.

In order to further investigate the relationship between the binding affinity of the compounds and their effect on myosin ATPase activity, different linear regression models were built using different

combinations of energy terms (Table S7). A comparison of the adjusted R-squared values shows that the largest proportion of ATPase activity variance (~66%, Model 1) is explained by the model that uses steric and hydrogen-bond differences between the PPS and NR state (Figure 2). Adding the hydrophobic term (Model 2) does not lead to any improvement. Interestingly, using only the PPS (Model 3) or NR (Model 4) terms rather than their difference produces models that explain only ~35% of the variance or less. At last, considering only the terms corresponding to a single binding pose (Model 5) rather than averaging them over the different Vina solutions, leads to a poor model, suggesting that multiple binding poses might be relevant at least for some of these compounds.

OM inhibits the dynamics of the PPS state

Further analysis of the PPS MD simulations shows that the increased (direct or OM-mediated) interactions between the different subdomains observed in the presence of OM are associated with a reduction in the global dynamics compared to the Apo simulations. This is particularly evident for the CLD, relay helix and N-terminal sub-domains, which show a strong reduction in flexibility (Figure 3 and S6), accompanied by an increase in dynamic cross correlation between the CLD and the nearby subdomains (red lines in Figure 4A and S7). A comparison of the RMSF profiles averaged over the replicas shows that the flexibility reduction is in general more pronounced for the PPS state (positive red values in Figure S8B) than the NR one (positive blue), especially in the N-terminal SH3 region and in the relay helix. This is consistent with the behaviour of the RMSD time evolution, which shows a much larger difference between the plateau values of Apo and OM-bound PPS simulations (Figure S9A) compared to NR ones (Figure S9B).

A principal component analysis of the pseudo-trajectory obtained by concatenating all the simulations performed so far (Apo and OM-bound motor domain starting from either the PPS or NR conformation) shows a dominant pseudo-motion (~70% of the overall variance) involving mostly the rotation of the CLD (PC1 in Figures 5 and S10). Data in Figure 5 are plotted so that an increase in

Figure 3. Effect of OM on the flexibility of the motor domain in the PPS state. Root mean square fluctuations (RMSF) colour are mapped onto the motor domain average structure from blue (0 Å) to red (3.4 Å) for one of the Apo (A) and OM-bound (B) simulations. The thickness of the tube representation is proportional to the RMSF value.

Figure 4. Effect of OM on global and local correlations in the PPS state. A. Difference dynamic cross correlation matrices (Δ DCCM) calculated between a pair of OM-bound and Apo simulations mapped onto the initial structure. The red edges connect residue pairs with Δ DCCM > 0.16. B. Mapping of $\Delta\zeta$ scores for the fragment starting from D168 (orange sphere). Regions with negative $\Delta\zeta$ scores are highlighted by using a thicker tube representation (with the thickness proportional to the magnitude of $\Delta\zeta$ values) and coloured according to their sub-domain. A negative $\Delta\zeta$ score indicates a stronger preferential connection to the D168 fragment in OM-bound simulations compared to Apo ones.

PC1 corresponds to a recovery stroke-like rotation of the CLD, while a decrease corresponds to a power stroke-like motion. The presence of OM (blue in Figure 5) greatly constrains the motor domain in both directions, this effect however is much more evident in the PPS simulations. Indeed, when in the PPS Apo state (solid green lines), the protein samples conformations that are either further along the recovery stroke compared to the initial structure ($PC1 > 100 \text{ \AA}$) or moving back towards the NR state in the power-stroke direction ($PC1 < 70 \text{ \AA}$). OM (solid blue lines) inhibits motion in both directions and freezes the domain to conformations close to the starting one ('P' letter). The higher peaks observed for the PPS OM-bound PC1 projections compared to the NR ones (dashed blue lines) indicate a more pronounced inhibitory effect of OM when bound to the PPS state. Motion is inhibited by OM also along PC2 and PC3 (Figures S10 and S11 B-C), which involve smaller amplitude ($\sim 5\%$ variance) rotations of the N-terminal subdomain (PC2) and of CLD (PC3). It is also to be noted that for each state, the different replicas explore similar regions of the 1D or 2D projections on the PC space (Figure 5 and S10), indicating a good level of sampling convergence, especially for the OM-bound simulations.

Figure 5. Principal Component Analysis of the PPS and NR trajectories. A. Porcupine representation of the first principal component (PC1) calculated on the pseudo-trajectory obtained by concatenating all the 16 trajectories together (Apo/OM-bound PPS and NR). The orange spikes show the direction and relative amplitude of motion of each residue along the PC. Spikes shorter than 20% of the maximum length are not shown. Highly flexible unstructured regions were excluded from the calculation. B. Distribution of projections of PPS (solid lines) and NR (dashed lines) trajectories of the Apo (green shades) and OM-bound (blue shades) simulations. The position of initial structures is indicated with letters (n = Apo NR, N = OM-bound NR, p = Apo PPS, P = OM-bound PPS).

Consistently with the overall dynamics of the protein, the C-terminal end of the key structural element relay helix shows on average a strong reduction in flexibility when OM is bound (Figure S12). OM-bound structures (blue) are stabilised in a kinked conformation close to the initial one (transparent cartoon in Figure S12B), while Apo simulations (green) can sample more bent conformations. A similar behaviour was observed in the NR simulations, where however OM stabilised a straight conformation of the helix¹⁰.

OM closes possible Pi release pathways in actin-unbound myosin

In order to identify possible correlations coupling the OM binding site with distant regions in PPS structures, a network of local correlated motions was reconstructed using the M32K25 Structural Alphabet (Supplementary

Figure 6. Pocket analysis in the nucleotide binding site. Pocket frequency maps for PPS Apo (A) and OM-bound (B) simulations. The maps are represented as isosurfaces connecting points with pocket frequency = 0.3. The position of the residues lining the backdoor I tube is indicated in magenta in the cartoon representation, while the approximate position of the tube is indicated as a dashed magenta arrow in A. The discontinuity between the Pi-containing pocket and the backdoor I tube observed in the OM-bound simulation is indicated with an orange arrow in B. Inset panels on the right show backdoor I residues in contact (distance $< 4 \text{ \AA}$) for at least 50% of the trajectory (edges). The narrower backdoor I tubes observed in OM-bound simulations seem to be correlated with the presence of S180-E466, R243-A463 and R237-E466 contacts (orange edges).

Methods)³⁵ and the 4-residue fragments with enhanced preferential connection scores ($\Delta\zeta < 0$) to the OM binding site were identified (Figure 4B). The $\Delta\zeta$ profiles of the different fragments covering the OM-binding region present similar features (Figure S13 and S14). Remarkably, Switch 2, part of the G-helix and the $\beta 5$ and $\beta 7$ strands from the transducer show large negative $\Delta\zeta$ values in most of the cases. Except for $\beta 7$, these same elements were preferentially connected to OM-binding fragments also in the NR state, indicating that these OM-enhanced long-range couplings are a general feature that does not depend on the specific position of the CLD sub-domain.

The local correlation analysis thus suggests that OM-binding to the PPS state might affect the nucleotide binding site, since it enhances the coupling with one of its most important regions (Switch 2). Changes in the shape of the binding pocket and other cavities in this region during the simulations were analysed using MDpocket (Supplementary Methods). 3D maps of pocket frequency values were generated (surfaces in Figure 6), providing the fraction of time a given point in space is found to be part of a pocket. A channel connecting the ATP binding site with the external surface of the protein was observed in three out of four Apo simulations (green surfaces in Figure 6A and S15). The channel is lined by residues from the N-terminal and U50K sub-domains (magenta) previously reported to be part of one of the main Pi release pathways ('backdoor I' inner and outer tube)¹⁹. Interestingly, the channel is closed in all the OM-bound simulations (Figure 6B and S15), indicating that the presence of OM might make Pi release through this pathway more difficult. The narrower channels observed in the presence of OM seem to be correlated with the presence of persistent contacts connecting Switch-2 residues (A463 and E466) with either Switch 1 (R237 and R243) or the P-loop (S180) (orange edges in Figure 6B and S16).

Discussion

The MD simulations presented in this work both confirm the results from previous simulations of the NR state and provide new crucial PPS-specific information on the molecular mechanism of action of OM.

The NR simulations indicated that the presence of OM can hinder the recovery stroke (CLD down to up transition), but they did not give any information on the effect of OM on the power stroke. The present findings indicate that OM has an even stronger inhibitory effect on the CLD rotation when the CLD is up. This would suggest that the presence of OM would slow down or even prevent rotation of the CLD from the up to the down conformation required for the power-stroke motion. This agrees very well with experimental data from different biophysical techniques (including FRET and single molecule optical trapping) that show that OM inhibits the power stroke by decreasing its velocity⁸ and size⁹, to the point of completely suppressing it at high concentrations. It is however to be noted that, while the present simulations show a reduction in the intrinsic dynamics of the PPS state and in particular in the rotational motions of the CLD, their length does not allow a complete power stroke transition to be observed. Further simulations, possibly combined with enhanced sampling techniques, will need to be performed in the future to directly observe the effect of OM on the power stroke.

The modelling of myosin-ligand complexes for a series of close OM analogues indicated that a) their effect on the ATPase activity of myosin is modulated by their ability to selectively bind the PPS state, i.e, by their relative PPS vs NR binding affinities rather than absolute ones; b) steric and hydrogen-bonding contributions tend to be more important than the hydrophobic ones in determining the activity of these compounds and c) more active compounds tend to form interactions that cross-couple multiple sub-domains in the protein. The regression model used to fit the experimental data, while

accounting for a surprisingly large proportion of data variance considering its simplicity, does not completely explain the changes in ATPase activity. This indicates that other factors might be at play, for example possible ligand-induced long-range effects on myosin structure and dynamics, which are not included in this model. It is also to be noted that the considerations above are valid for the selected subset of OM analogues. Building a general predictive model for myosin activators would require a larger and more diverse dataset than then one considered here.

In addition to inhibiting the CLD motions, simulations show that OM can indeed affect functional regions that are distant from its binding region. In the PPS state, the presence of OM seems to particularly affect the nucleotide binding site, since a) it induces large increments in the preferential connection between the OM binding site and Switch 2 and b) it closes possible pathways for Pi release. The second result in particular could explain why OM decreases the rate of Pi release in the absence of actin^{2,7}. The escape of Pi from its binding site is never observed in our simulations, as expected considering that their timescale (300 ns) is much shorter than the time required for Pi release (ms to s). However, a channel connecting the Pi side of the nucleotide binding site to the molecular surface is present in the Apo simulations, forming a possible escape route for the molecule. The residues lining the channel are part of the main pathway for Pi release from the PPS state of actin-unbound myosin (backdoor I), as previously identified with MD simulations¹⁹. Our results indicate that the narrower channels observed when OM is bound to the protein are due to OM-induced contacts between specific residues in these regions, which increase the interactions between Switch 2 and nearby regions and thus disfavour the release of Pi from this route.

It is important to note that these results are not in contradiction with the OM-induced increase of Pi release observed in the presence of actin (actin-activated Pi release), since the escape of Pi from the actomyosin complex has been shown to occur mostly through a different route (backdoor II)¹⁹, which is formed by the opening of the P-loop during the transition from the weakly to strongly actin-bound myosin state. Since the release of Pi through backdoor II is faster than when using other escape routes, a possibility is that OM increases the actin-activated Pi release rate by favouring the weakly-to-strongly actin bound state transition and thus the opening of backdoor II. Future simulations of actin-bound myosin complexes will need to be performed to clarify this point.

Overall, the present findings are compatible with the current view that OM is not a traditional myosin activator but presents a mixture of activating and inhibitory effects. In particular, they support the current mechanistic model that OM increases heart contractility by preferentially binding to myosin states with the CLD in the up conformation compared to the down one and by promoting the formation of strongly interacting actomyosin complexes where the power stroke is inhibited. This prolongs the time during which myosin is strongly bound to actin and enhances muscle contraction by promoting thin filament activation and cooperative binding of fully functional myosin motor domains that are not bound to OM⁹. This up-regulation could be further enhanced by the OM-induced stabilisation of the ON state of the myosin heads in the thick filament over the OFF one observed using fluorescent probes³⁶ or hypothesised on the basis of MD simulations¹⁴.

ASSOCIATED CONTENT

The Supporting Information is available free of charge at <https://pubs.acs.org/>

Table S1. Residues within 4 Å of OM functional groups in the PPS and NR states. Table S2. Cluster analysis of OM-bound simulations. Tables S3 and S4. Frequency of occurrence (occupancy) of OM-myosin hydrogen bonds and contacts during OM-bound PPS and NR simulations. Table S5 - Average

frequency of occurrence (occupancy) of hydrogen bonds near the OM binding site (AvgHB). Table S6 – Activity and binding affinity data for OM and analogues. Table S7 – Linear models of % increase in ATPase activity for OM and analogues.

Figure S1. Stability of OM binding to myosin. Figure S2. Difference in HB frequency of occurrence (ΔHBfreq) of hydrogen bonds near the OM binding site. Figure S3. OM-stabilised hydrogen bonds near the OM binding site. Figure S4. Chemical structures of OM analogues. Figure S5. Models of OM analogues bound to the PPS and NR conformations. Figure S6. RMSF of the motor domain (PPS state) in Apo and OM-bound simulations. Figure S7. Comparison of dynamic cross-correlation networks of OM-bound and Apo simulations of the PPS state. Figure S8. Comparison of RMSF profiles. Figure S9. Time evolution of RMSD in PPS and NR simulations. Figure S10. Principal component analysis of PPS and NR trajectories (Apo and OM-bound states). Figure S11. First three Principal Components of the concatenated pseudo-trajectory (PPS and NR, Apo and OM-bound states). Figure S12. Bending of the relay helix during the MD simulations of the PPS state. Figure S13. Profiles of $\Delta\zeta$ scores calculated for fragments in the OM-binding site. Figure S14. Mapping of $\Delta\zeta$ scores calculated for fragments in the OM-binding site. Figure S15. Pocket analysis in the nucleotide binding site. Figure S16. Inter-residue contacts in the backdoor I region.

AUTHOR INFORMATION

Corresponding Author

*Arianna Fornili, email: a.fornili@qmul.ac.uk

Present Addresses

†Department of Chemistry, University of Pisa, Via Giuseppe Moruzzi 13, 56124 Pisa, Italy

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources

Royal Society of Chemistry Undergraduate Research Bursary (grant number U19-1596) to WGD.

Acknowledgements

This work made use of time on ARCHER granted via the UK High-End Computing Consortium for Biomolecular Simulation, HECBioSim (<http://hecbiosim.ac.uk>), supported by EPSRC (grant no. EP/L000253/1). The work was also partly supported by a Royal Society of Chemistry Undergraduate Research Bursary (grant number U19-1596) to WGD.

REFERENCES

- (1) Mitchell A Psotka; Stephen S Gottlieb; Gary S Francis; Allen, L. A.; Teerlink, J. R.; Jr, K. F. A.; Rosano, G. M. C.; Lancellotti, P. Cardiac Calcitropes, Myotropes, and Mitotropes. *J. Am. Coll. Cardiol.* **2019**, *73*, 2345–2353. <https://doi.org/10.1016/j.jacc.2019.02.051>.
- (2) Malik, F. I.; Hartman, J. J.; Elias, K. a; Morgan, B. P.; Rodriguez, H.; Brejc, K.; Anderson, R. L.; Sueoka, S. H.; Lee, K. H.; Finer, J. T.; Sakowicz, R.; Baliga, R.; Cox, D. R.; Garard, M.; Godinez, G.; Kawas, R.; Kraynack, E.; Lenzi, D.; Lu, P. P.; Muci, a; Niu, C.; Qian, X.; Pierce, D. W.; Pokrovskii, M.; Suehiro, I.; Sylvester, S.; Tochimoto, T.; Valdez, C.; Wang, W.; Katori, T.; Kass, D. a; Shen, Y.-T.; Vatner, S. F.; Morgans, D. J. Cardiac Myosin Activation: A Potential Therapeutic Approach for Systolic Heart Failure. *Science* **2011**, *331*, 1439–1443. <https://doi.org/10.1126/science.1200113>.
- (3) Kaplinsky, E.; Mallarkey, G. Cardiac Myosin Activators for Heart Failure Therapy: Focus on Omecamtiv Mecarbil. *Drugs Context* **2018**, *7*, 1–10. <https://doi.org/10.7573/dic.212518>.
- (4) Muir, W. W.; Hamlin, R. L. Myocardial Contractility: Historical and Contemporary Considerations. *Front. Physiol.* **2020**, *11*. <https://doi.org/10.3389/fphys.2020.00222>.
- (5) Ochala, J.; Sun, Y.-B. Novel Myosin-Based Therapies for Congenital Cardiac and Skeletal Myopathies. *J. Med. Genet.* **2016**, *53*, 651–654. <https://doi.org/10.1136/jmedgenet-2016-103881>.
- (6) Hwang, P. M.; Sykes, B. D. Targeting the Sarcomere to Correct Muscle Function. *Nat. Rev. Drug Discov.* **2015**, *14*, 313–328. <https://doi.org/10.1038/nrd4554>.
- (7) Liu, Y.; White, H. D.; Belknap, B.; Winkelmann, D. A.; Forgacs, E. Omecamtiv Mecarbil Modulates the Kinetic and Motile Properties of Porcine β -Cardiac Myosin. *Biochemistry* **2015**, *54*, 1963–1975. <https://doi.org/10.1021/bi5015166>.
- (8) Rohde, J. A.; Thomas, D. D.; Muretta, J. M.; Failure, H.; Mecarbil, O.; Myosin, F. A Heart Failure Drug Changes the Mechano-Enzymology of the Cardiac Myosin Powerstroke. *Proc. Natl. Acad. Sci.* **2017**, *114*, E1796–E1804. <https://doi.org/10.1073/pnas.1611698114>.
- (9) Woody, M. S.; Greenberg, M. J.; Barua, B.; Winkelmann, D. A.; Goldman, Y. E.; Ostap, E. M. Positive Cardiac Inotrope Omecamtiv Mecarbil Activates Muscle despite Suppressing the Myosin Working Stroke. *Nat. Commun.* **2018**, *9*, 3838. <https://doi.org/10.1038/s41467-018-06193-2>.
- (10) Hashem, S.; Tiberti, M.; Fornili, A. Allosteric Modulation of Cardiac Myosin Dynamics by Omecamtiv Mecarbil. *PLoS Comput. Biol.* **2017**, *13*, e1005826. <https://doi.org/10.1371/journal.pcbi.1005826>.
- (11) Winkelmann, D.; Forgacs, E.; Miller, M. T.; Stock, A. M. Structural Basis for Drug-Induced Allosteric Changes to Human β -Cardiac Myosin Motor Activity. *Nat. Commun.* **2015**, *6*, 7974. <https://doi.org/10.1038/ncomms8974>.
- (12) Planelles-Herrero, V. J.; Hartman, J. J.; Robert-Paganin, J.; Malik, F. I.; Houdusse, A. Mechanistic and Structural Basis for Activation of Cardiac Myosin Force Production by Omecamtiv Mecarbil. *Nat. Commun.* **2017**, *8*, 190. <https://doi.org/10.1038/s41467-017-00176-5>.
- (13) Viswanathan, M. C.; Schmidt, W.; Franz, P.; Rynkiewicz, M. J.; Newhard, C. S.; Madan, A.; Lehman, W.; Swank, D. M.; Preller, M.; Cammarato, A. A Role for Actin Flexibility in Thin Filament-Mediated Contractile Regulation and Myopathy. *Nat. Commun.* **2020**, *11*, 2417. <https://doi.org/10.1038/s41467-020-15922-5>.
- (14) Robert-Paganin, J.; Auguin, D.; Houdusse, A. Hypertrophic Cardiomyopathy Disease Results from Disparate Impairments of Cardiac Myosin Function and Auto-Inhibition. *Nat. Commun.* **2018**, *1*–13. <https://doi.org/10.1038/s41467-018-06191-4>.
- (15) Aprahamian, M. L.; Tikunova, S. B.; Price, M. V.; Cuesta, A. F.; Davis, J. P.; Lindert, S. Successful Identification of Cardiac Troponin Calcium Sensitizers Using a Combination of Virtual Screening and ROC Analysis of Known Troponin C Binders. *J. Chem. Inf. Model.* **2017**, *57*, 3056–3069. <https://doi.org/10.1021/acs.jcim.7b00536>.
- (16) Bowman, J. D.; Lindert, S. Computational Studies of Cardiac and Skeletal Troponin. *Front. Mol. Biosci.* **2019**, *6*. <https://doi.org/10.3389/fmolb.2019.00068>.
- (17) Tiberti, M.; Lechner, B.-D.; Fornili, A. Binding Pockets in Proteins Induced by Mechanical Stress. *J. Chem. Theory Comput.* **2019**, *15*, 1–6. <https://doi.org/10.1021/acs.jctc.8b00755>.
- (18) Porter, J. R.; Meller, A.; Zimmerman, M. I.; Greenberg, M. J.; Bowman, G. R. Conformational Distributions of Isolated Myosin Motor Domains Encode Their Mechanochemical Properties. *eLife* **2020**, *9*, e55132. <https://doi.org/10.7554/eLife.55132>.
- (19) Cecchini, M.; Alexeev, Y.; Karplus, M. Pi Release from Myosin: A Simulation Analysis of Possible Pathways. *Structure* **2010**, *18*, 458–470. <https://doi.org/10.1016/j.str.2010.01.014>.
- (20) Lindorff-Larsen, K.; Maragakis, P.; Piana, S.; Eastwood, M. P.; Dror, R. O.; Shaw, D. E. Systematic Validation of Protein Force Fields against Experimental Data. *PLoS ONE* **2012**, *7*, e32131. <https://doi.org/10.1371/journal.pone.0032131>.
- (21) Wang, J.; Wolf, R. M.; Caldwell, J. W.; Kollman, P. A.; Case, D. A. Development and Testing of a General Amber Force Field. *J. Comput. Chem.* **2004**, *25*, 1157–1174. <https://doi.org/10.1002/jcc.20035>.
- (22) Meagher, K. L.; Redman, L. T.; Carlson, H. A. Development of Polyphosphate Parameters for Use with the AMBER Force Field. *J. Comput. Chem.* **2003**, *24*, 1016–1025. <https://doi.org/10.1002/jcc.10262>.

- (23) Case, D. A.; Berryman, J. T.; Betz, R. M.; Cerruti, D. S.; Cheatham, T. E.; Darden, T. A.; Duke, R. E.; Giese, T. J.; Gohlke, H.; Goetz, A. W.; Homeyer, N.; Izadi, S.; Janowski, P.; Kaus, J.; Kovalenko, A.; Lee, T. S.; LeGrand, S.; Li, P.; Luchko, T.; Luo, R.; Madeij, B.; Merz, K.; Monard, G.; Needham, P.; Nguyen, H.; Nguyen, H. T.; Omelyan, I.; Onufriev, A.; Roe, D. R.; Roitberg, A.; Salomon-Ferrer, R.; Simmerling, C. L.; Smith, W.; Swails, J.; Walker, R. C.; Wang, J.; Wolf, R. M.; Wu, X.; York, D. M.; Kollman, P. A. (2015), Amber 2015, University of California, San Francisco.
- (24) Sousa da Silva, A. W.; Vranken, W. F. ACPYPE - AnteChamber PYthon Parser InterfacE. *BMC Res Notes* **2012**, *5*, 367. <https://doi.org/10.1186/1756-0500-5-367>.
- (25) Schmidtke, P.; Bidon-Chanal, A.; Luque, F. J.; Barril, X. MDpocket: Open-Source Cavity Detection and Characterization on Molecular Dynamics Trajectories. *Bioinformatics* **2011**, *27*, 3276–3285. <https://doi.org/10.1093/bioinformatics/btr550>.
- (26) Le Guilloux, V.; Schmidtke, P.; Tufféry, P. Fpocket: An Open Source Platform for Ligand Pocket Detection. *BMC Bioinformatics* **2009**, *10*, 168–11. <https://doi.org/10.1186/1471-2105-10-168>.
- (27) Daura, X.; van Gunsteren, W. F.; Mark, A. E. Peptide Folding: When Simulation Meets Experiment. *Angew. Chem. Int. Ed.* **1999**, *38*, 236–240. <https://doi.org/10.1002/9781118183373.fmatter>.
- (28) Skjærven, L.; Yao, X. Q.; Scarabelli, G.; Grant, B. J. Integrating Protein Structural Dynamics and Evolutionary Analysis with Bio3D. *BMC Bioinformatics* **2014**, *15*, 399. <https://doi.org/10.1186/s12859-014-0399-6>.
- (29) Manickam, M.; Jalani, H. B.; Pillaiyar, T.; Sharma, N.; Boggu, P. R.; Venkateswararao, E.; Lee, Y.-J.; Jeon, E.-S.; Jung, S.-H. Exploration of Flexible Phenylpropylurea Scaffold as Novel Cardiac Myosin Activators for the Treatment of Systolic Heart Failure. *Eur. J. Med. Chem.* **2017**, *134*, 379–391. <https://doi.org/10.1016/j.ejmech.2017.04.005>.
- (30) Manickam, M.; Jalani, H. B.; Pillaiyar, T.; Boggu, P. R.; Sharma, N.; Venkateswararao, E.; Lee, Y.-J.; Jeon, E.-S.; Son, M.-J.; Woo, S.-H.; Jung, S.-H. Design and Synthesis of Sulfonamidophenylethylureas as Novel Cardiac Myosin Activator. *Eur. J. Med. Chem.* **2018**, *143*, 1869–1887. <https://doi.org/10.1016/j.ejmech.2017.10.077>.
- (31) Kim, S.; Chen, J.; Cheng, T.; Gindulyte, A.; He, J.; He, S.; Li, Q.; Shoemaker, B. A.; Thiessen, P. A.; Yu, B.; Zaslavsky, L.; Zhang, J.; Bolton, E. E. PubChem 2019 Update: Improved Access to Chemical Data. *Nucleic Acids Res.* **2019**, *47*, D1102–D1109. <https://doi.org/10.1093/nar/gky1033>.
- (32) Morris, G. M.; Huey, R.; Lindstrom, W.; Sanner, M. F.; Belew, R. K.; Goodsell, D. S.; Olson, A. J. AutoDock4 and AutoDockTools4: Automated Docking with Selective Receptor Flexibility. *J. Comput. Chem.* **2009**, *30*, 2785–2791. <https://doi.org/10.1002/jcc.21256>.
- (33) Trott, O.; Olson, A. J. AutoDock Vina: Improving the Speed and Accuracy of Docking with a New Scoring Function, Efficient Optimization, and Multithreading. *J. Comput. Chem.* **2009**, *31*, 455–461. <https://doi.org/10.1002/jcc.21334>.
- (34) Schrodinger, LLC. The PyMOL Molecular Graphics System, Version 2.0.
- (35) Pandini, A.; Fornili, A.; Fraternali, F.; Kleinjung, J. GSATools: Analysis of Allosteric Communication and Functional Local Motions Using a Structural Alphabet. *Bioinformatics* **2013**, *29*, 2053–2055. <https://doi.org/10.1093/bioinformatics/btt326>.
- (36) Kampourakis, T.; Zhang, X.; Sun, Y.-B.; Irving, M. Omecamtiv Mercabil and Blebbistatin Modulate Cardiac Contractility by Perturbing the Regulatory State of the Myosin Filament. *J. Physiol.* **2018**, *596*, 31–46. <https://doi.org/10.1113/JP275050>.

For Table of Contents Only

Heart Failure Drug Modifies the Intrinsic Dynamics of the Pre-Power Stroke State of Cardiac Myosin

Shaima Hashem,^{a†} William George Davies^a and Arianna Fornili^{a,b*}

^a School of Biological and Chemical Sciences, Queen Mary University of London, London, E1 4NS, United Kingdom.

^b The Thomas Young Centre for Theory and Simulation of Materials, London, WC1E 6BN, United Kingdom

[†] Present address: Department of Chemistry, University of Pisa, Via Giuseppe Moruzzi 13, 56124 Pisa, Italy

*corresponding author, email: a.fornili@qmul.ac.uk

Supplementary Information

Table S1. Residues within 4 Å of OM functional groups in the PPS and NR states.

OM functional group	PPS ^a	NR ^a
Carboxymethyl-piperazine (a)	K146, R147, S148, N160, Q163, Y164 N711, R721, A767, L770 , Y722, G771, E774	M90, A91, L96, V101, S118, G119, L120 M493 I702, C705 P710
Fluoro-benzyl (b)	T167, Y164, D168 I713, N711, R712, L770	A91, G119, L120 N711, P710, R712
Amino-carbamoyl (c)	Y164, D168 N711, R712	A91, M92 R712
Methylpyridin-3-yl (d)	D168, Y164 H492 H666 P710, N711, R712, K762	A91, M92, F84 N711, R712, I713, L770

^aThe residues were determined on representative structures extracted from MD simulations of the motor domain in the OM-bound PPS and NR state. Representative structures were determined through cluster analysis of the concatenated trajectories of each state (see Methods). Residues are coloured according to their sub-domain (see Figure 1 legend). Residues common to the PPS and NR pockets are highlighted in bold.

Table S2. Cluster analysis of OM-bound simulations.

# cluster	population ^a (%)	RMSD ^b (Å)	MD1 ^c (%)	MD2 ^c (%)	MD3 ^c (%)	MD4 ^c (%)
1	87.8	1.5	24.4	19.0	23.3	21.1
2	7.9	2.9	0.6	5.3	0.0	2.1
3	2.2	1.7	0.0	0.0	1.6	0.5
4	0.7	2.7	0.0	0.0	0.0	0.7

^aFraction of the concatenated OM-bound trajectory belonging to each cluster.

^bRMSD from the OM X-ray structure non-hydrogen atoms. The fitting of the structures was performed using all the C^α atoms in the protein.

^cContribution to the cluster population from each OM-bound simulation.

Table S3. Frequency of occurrence (occupancy) of OM-myosin hydrogen bonds during OM-bound PPS and NR simulations

PPS ^a						NR ^{a,b}					
Resid ^c	MD1	MD2	MD3	MD4	Average	Resid ^c	OMA1	OMA2	OMB1	OMB2	Average
N711	0.83	0.58	0.81	0.68	0.72	R712	0.18	0.47	0.61	0.31	0.39
D168	0.71	0.58	0.82	0.30	0.60	C705	0.04	0.50	0.02	0.47	0.25
R712	0.54	0.60	0.50	0.45	0.52	A91	0.11	0.22	0.25	0.15	0.18
R147	0.07	0.13	0.00	0.19	0.10	N711	0.09	0.22	0.22	0.00	0.14
Q163	0.01	0.18	0.01	0.05	0.06	S118	0.06	0.12	0.00	0.10	0.07
N160	0.00	0.03	0.18	0.00	0.05	K762	0.21	0.00	0.01	0.01	0.06

^aHydrogen bonds calculated with VMD on snapshots sampled every 100 ps using a threshold of 3.5 Å on the D(onor)-A(cceptor) distance and 30° on the D-H-A angle. Only residues with frequency > 0.1 in at least one replica are reported.

^bNR values are taken from Ref. 10 in the main text and reported here for ease of comparison.

^cResidues are coloured according to their subdomain (see Figure 1 legend).

Table S4. Frequency of occurrence (occupancy) of OM-myosin contacts during OM-bound PPS and NR simulations.

Resid ^c	PPS ^a					Average	Resid	NR ^{a,b}					Average
	OM1	OM2	OM3	OM4	Average			OMA1	OMA2	OMB1	OMB2		
Y164	1.00 (1.00)	1.00 (1.00)	1.00 (1.00)	1.00 (1.00)	1.00 (1.00)	1.00 (1.00)	A91	0.96 (0.84)	1.00 (0.97)	1.00 (0.96)	1.00 (0.93)	0.99 (0.93)	
R712	1.00 (0.99)	1.00 (0.96)	1.00 (0.99)	0.97 (0.86)	0.99 (0.95)	0.99 (0.95)	R712	0.94 (0.93)	1.00 (0.99)	1.00 (1.00)	1.00 (1.00)	0.99 (0.98)	
L770	0.97 (0.97)	0.91 (0.87)	0.96 (0.96)	0.97 (0.97)	0.95 (0.94)	0.95 (0.94)	P710	0.90 (0.10)	1.00 (0.15)	1.00 (0.36)	1.00 (0.15)	0.97 (0.19)	
N711	1.00 (1.00)	0.90 (0.78)	1.00 (1.00)	0.90 (0.89)	0.95 (0.92)	0.95 (0.92)	S118	0.99 (0.57)	0.97 (0.85)	0.88 (0.72)	0.96 (0.85)	0.95 (0.75)	
T167	0.98 (0.98)	0.94 (0.94)	0.97 (0.97)	0.88 (0.88)	0.94 (0.94)	0.94 (0.94)	M92	0.83 (0.60)	1.00 (0.93)	0.98 (0.85)	1.00 (0.94)	0.95 (0.83)	
H666	0.97 (0.97)	0.88 (0.88)	0.95 (0.95)	0.95 (0.95)	0.94 (0.94)	0.94 (0.94)	C705	0.94 (0.94)	0.96 (0.96)	0.87 (0.77)	1.00 (1.00)	0.94 (0.92)	
P710	0.97 (0.00)	0.88 (0.00)	0.93 (0.01)	0.82 (0.00)	0.90 (0.00)	0.90 (0.00)	I702	0.71 (0.63)	0.91 (0.91)	0.96 (0.82)	0.97 (0.97)	0.89 (0.83)	
N160	0.94 (0.94)	0.69 (0.68)	0.94 (0.94)	0.88 (0.88)	0.86 (0.86)	0.86 (0.86)	L96	0.70 (0.70)	0.86 (0.86)	0.80 (0.80)	0.83 (0.83)	0.80 (0.80)	
Q163	0.94 (0.91)	0.84 (0.82)	0.78 (0.74)	0.76 (0.71)	0.83 (0.79)	0.83 (0.79)	N711	0.28 (0.22)	0.96 (0.85)	0.99 (0.48)	0.95 (0.50)	0.80 (0.51)	
I713	0.71 (0.71)	0.84 (0.84)	0.74 (0.74)	0.82 (0.82)	0.77 (0.77)	0.77 (0.77)	M90	0.50 (0.50)	0.97 (0.68)	0.55 (0.50)	0.99 (0.92)	0.75 (0.65)	
E774	0.89 (0.89)	0.47 (0.47)	0.87 (0.87)	0.83 (0.83)	0.76 (0.76)	0.76 (0.76)	G701	0.52 (0.00)	0.61 (0.00)	0.97 (0.00)	0.47 (0.00)	0.64 (0.00)	
Y722	0.73 (0.73)	0.56 (0.56)	0.81 (0.81)	0.81 (0.81)	0.72 (0.72)	0.72 (0.72)	M493	0.54 (0.54)	0.62 (0.62)	0.59 (0.59)	0.60 (0.60)	0.59 (0.59)	
R147	0.94 (0.93)	0.84 (0.82)	0.05 (0.05)	0.97 (0.97)	0.70 (0.69)	0.70 (0.69)	G119	0.89 (0.00)	0.60 (0.00)	0.35 (0.00)	0.33 (0.00)	0.54 (0.00)	
D168	0.74 (0.74)	0.60 (0.60)	0.88 (0.88)	0.35 (0.35)	0.64 (0.64)	0.64 (0.64)	L93	0.16 (0.00)	0.52 (0.00)	0.60 (0.00)	0.81 (0.01)	0.52 (0.00)	
K146	0.89 (0.89)	0.26 (0.24)	0.86 (0.86)	0.21 (0.21)	0.56 (0.55)	0.56 (0.55)	T94	0.15 (0.11)	0.56 (0.51)	0.60 (0.43)	0.75 (0.67)	0.51 (0.43)	
R721	0.57 (0.57)	0.44 (0.44)	0.49 (0.49)	0.50 (0.50)	0.50 (0.50)	0.50 (0.50)	L770	0.15 (0.15)	0.59 (0.59)	0.47 (0.47)	0.48 (0.48)	0.42 (0.42)	
G771	0.45 (0.00)	0.55 (0.00)	0.19 (0.00)	0.61 (0.00)	0.45 (0.00)	0.45 (0.00)	E497	0.13 (0.13)	0.30 (0.30)	0.67 (0.67)	0.50 (0.50)	0.40 (0.40)	
S148	0.78 (0.76)	0.06 (0.05)	0.00 (0.00)	0.86 (0.76)	0.42 (0.40)	0.42 (0.40)	V698	0.90 (0.26)	0.01 (0.00)	0.51 (0.24)	0.05 (0.01)	0.37 (0.13)	
L120	0.46 (0.46)	0.56 (0.56)	0.06 (0.06)	0.59 (0.57)	0.42 (0.41)	0.42 (0.41)	F121	0.28 (0.28)	0.00 (0.00)	0.81 (0.81)	0.32 (0.32)	0.35 (0.35)	
E497	0.18 (0.18)	0.12 (0.12)	0.64 (0.64)	0.08 (0.08)	0.26 (0.26)	0.26 (0.26)	F84	0.17 (0.17)	0.08 (0.08)	0.25 (0.25)	0.60 (0.60)	0.27 (0.27)	
H492	0.04 (0.04)	0.08 (0.08)	0.45 (0.45)	0.05 (0.05)	0.16 (0.16)	0.16 (0.16)	I713	0.10 (0.09)	0.77 (0.77)	0.09 (0.09)	0.13 (0.13)	0.27 (0.27)	
A767	0.47 (0.05)	0.00 (0.00)	0.12 (0.01)	0.00 (0.00)	0.15 (0.01)	0.15 (0.01)	V101	0.00 (0.00)	0.16 (0.16)	0.08 (0.08)	0.39 (0.39)	0.16 (0.16)	
E500	0.11 (0.11)	0.13 (0.13)	0.10 (0.10)	0.09 (0.09)	0.11 (0.11)	0.11 (0.11)	L120	0.39 (0.30)	0.04 (0.03)	0.07 (0.07)	0.05 (0.04)	0.14 (0.11)	
							K762	0.36 (0.36)	0.03 (0.03)	0.02 (0.02)	0.03 (0.03)	0.11 (0.11)	
							E500	0.08 (0.08)	0.01 (0.01)	0.21 (0.21)	0.04 (0.04)	0.08 (0.08)	
							F489	0.13 (0.13)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.03 (0.03)	

^aA residue is considered in contact with OM if the OM-residue minimum distance calculated over non-hydrogen atoms is < 4 Å. Values reported in parentheses are calculated considering only the atoms in the side chains. Only residues with frequency >= 0.1 in at least one replica are reported.

^bNR values are taken from Ref. 10 and reported here for ease of comparison.

^cResidues are coloured according to their sub-domain (see Figure 1 legend).

Table S5 - Average frequency of occurrence (occupancy) of hydrogen bonds near the OM binding site (AvgHB).

HB ^a	PPS Apo	PPS OM-bound	NR Apo	NR OM-bound
Intra N-term				
R17-S148	0.00	0.46	0.00	0.00
Intra RH				
E497-Y501	0.49	0.91	0.76	0.84
E499_K503	0.33	0.81	0.00	0.00
Intra SH1				
R671-N696	0.00	0.00	0.34	0.84
I704-G708	0.45	0.89	0.34	0.10
I704-K707	0.00	0.00	0.14	0.54
Intra CLD				
L714-F718	0.69	0.95	0.46	0.95
Y715-T761	0.33	0.78	0.00	0.00
N-term/CLD				
T94-N711	0.00	0.00	0.10	0.57
K146-D778	0.01	0.69	0.00	0.00
R169-D717	0.21	0.83	0.00	0.00
RH/CLD				
E500-K762	0.42	0.60	0.19	0.65
E504-K762	0.27	0.88	0.72	0.89
E504-T761	0.23	1.00	0.00	0.00

^aInter-residue hydrogen bonds within 10 Å of OM. Colour is used to highlight pairs where the HB frequency difference is largest in the PPS (blue) or the NR (red) state. Only pairs with $|\Delta\text{HBFreq}| > 0.4$ for at least one of the two states (PPS or NR) are shown.

Table S6 – Activity and binding affinity data for OM and analogues.

Compound ^a	% ATPase (exp) ^b	% ATPase (predicted) ^c	Binding affinity (NR) ^d	Binding affinity (PPS) ^d	Average Steric (NR) ^e	Average Steric (PPS) ^e	Average Hydrophobic (NR) ^e	Average Hydrophobic (PPS) ^e	Average hydrogen bond (NR) ^e	Average hydrogen bond (PPS) ^e
Group 1										
2a	10.3	18.0	-7.2	-8.0	-5.3	-5.5	-1.1	-1.7	-0.5	-0.7
5a	1.6	21.3	-6.8	-7.5	-5.0	-5.2	-1.1	-1.8	-0.3	-0.5
6a	11.9	13.5	-6.8	-7.7	-5.0	-5.4	-1.3	-1.8	-0.3	-0.3
11a	20.3	10.9	-7.0	-7.7	-5.3	-5.4	-0.9	-1.6	-0.3	-0.5
Group 2										
OM	58.0	70.0	-7.8	-9.7	-6.7	-6.6	-0.4	-0.6	-0.6	-1.2
1a	51.1	47.0	-7.3	-8.2	-4.8	-5.4	-1.1	-1.7	-0.9	-1.0
9a	53.3	58.3	-6.9	-7.9	-4.9	-5.6	-1.1	-1.6	-0.7	-0.6
15b	35.0	42.1	-6.9	-8.6	-5.2	-5.3	-0.9	-1.5	-0.9	-1.3
21a	44.0	65.0	-6.3	-7.6	-4.9	-5.4	-0.5	-0.9	-0.9	-1.1
Group 3										
10b	122.4	85.2	-7.5	-8.5	-5.4	-5.6	-0.8	-1.0	-0.9	-1.4
13b	91.6	119.4	-7.2	-8.5	-5.4	-6.1	-0.8	-1.0	-0.6	-1.0
14a	81.4	60.4	-6.6	-7.7	-4.9	-5.6	-0.8	-1.2	-0.7	-0.7
14b	95.1	65.0	-6.8	-8.5	-5.1	-5.6	-0.9	-1.5	-0.7	-1.0

^aCompounds are divided into three groups, with lower (group 1, light blue), similar (group 2, blue) or higher (group 3, orange) % increase in ATPase activity (experimental data) compared to OM.

^bChange in ATPase activity from Ref. 29 (**a** compounds) and Ref. 30 (**b** compounds) in the main text. The compound numbering of the original publication was used.

^cChange in ATPase activity predicted from the linear regression model 1 (Table S7).

^dBinding affinity (kcal/mol) predicted by AutoDock Vina (best binding pose).

^eIndividual components (steric, hydrophobic and hydrogen-binding) of the binding affinity (kcal/mol), averaged over all the Vina binding poses.

Table S7 – Linear models of % increase in ATPase activity for OM and analogues.

Models^a	Adjusted R-squared	Intercept	Intercept p-value	c1	c1 p-value	c2	c2 p-value	c3	c3 p-value
1	0.66	52.1	8.4E-04	-27.2	1.7E-03	-32.6	1.2E-03	-	-
2	0.62	53.3	2.7E-03	-28.6	1.5E-02	-3.2	8.4E-01	-35.4	4.1E-02
3	0.35	67.4	5.2E-01	-52.0	3.5E-01	-126.6	5.2E-02	-	-
4	0.32	194.2	4.4E-02	-33.9	4.1E-01	-223.0	2.4E-02	-	-
5	0.11	36.1	1.0E-01	-16.7	2.4E-01	-11.6	2.0E-01	-	-

^aThe linear models $Intercept + c_1x_1 + c_2x_2 (+ c_3x_3)$ were built-up using the following combinations of variables:

Model 1: x_1 = average steric score, x_2 = average HB score

Model 2: x_1 = average steric score, x_2 = average hydrophobic score, x_3 = average HB score

Model 3: x_1 = average steric score, x_2 = average HB score (PPS energy only)

Model 4: x_1 = average steric score, x_2 = average HB score (NR energy only)

Model 5: x_1 = steric score, x_2 = HB score (best binding pose only)

Figure S1. Stability of OM binding to myosin. A. Time evolution of the OM-protein distance for the PPS OM-bound simulations. The distance was calculated as the minimum value over all possible pairs of non-hydrogen atoms. B. Superimposition of the X-ray (blue) and cluster 1 representative (orange) OM structure. The protein is shown as white cartoon. The overall cluster population is reported in parentheses.

Figure S2. Difference in HB frequency of occurrence (ΔHBfreq) of hydrogen bonds near the OM binding site. Only hydrogen bonds between residues within 10 Å of OM are reported. Green lines indicate the -0.4 and 0.4 thresholds.

Figure S3. OM-stabilised hydrogen bonds near the OM binding site. Hydrogen bonds with $\Delta\text{HBfreq} > 0.4$ are indicated with yellow dashed lines on structures extracted from PPS (A) and NR (B) OM-bound simulations.

Figure S4. Chemical structures of OM analogues. The numbering of the compounds is the same as the one used in the original publication (a=Ref. 29, b=Ref. 30).

Figure S5. Models of OM analogues bound to the PPS and NR conformations. The ligands are represented as cyan (group1), blue (group 2) and orange (group 3) lines. The first 9 binding poses from Vina to the PPS (top row) and NR (bottom row) conformations are considered for each compound. Residues in the binding site are represented as sticks (carbon atoms in white), while polar contacts are shown as yellow dashed lines. Residues involved in recurring contacts are labelled.

Figure S6. RMSF of the motor domain (PPS state) in Apo and OM-bound simulations. The RMSF is colour mapped onto the motor domain structure from blue (0 Å) to red (3.4 Å) for each Apo (A) and OM-bound (B) replica. The average structure is used for each simulation. The thickness of the tube representation is proportional to the RMSF value. High flexibility regions and the OM binding site are also labelled.

Figure S7. Comparison of dynamic cross-correlation networks of OM-bound and Apo simulations of the PPS state. Difference DCCM matrices (ΔDCCM) calculated between pairs of OM-bound and Apo simulations are mapped onto the initial OM-bound structure. Edges connect residue pairs that have a positive (red) and a negative (green) ΔDCCM value, using a threshold of 0.16. The CLD domain is highlighted in blue. Highly flexible unstructured regions were excluded from the calculation (see Supplementary Methods)

Figure S8. Comparison of RMSF profiles. (A) Profiles of RMSF values for PPS Apo (green) and OM-bound (blue) simulations. All replicas are reported. RMSF values were calculated on C α atoms considering only the production part of the trajectories. (B) Difference between average Apo and OM-bound RMSF profiles calculated for PPS (red) and NR (blue) simulations. Positive values indicate residues with higher average flexibility in Apo simulations compared to OM-bound ones.

Figure S9. Time evolution of RMSD in PPS and NR simulations. The time evolution of RMSD values is reported for each replica of OM-bound (blue) and Apo (green) simulations of the PPS (A) and NR (state). Highly flexible unstructured regions were excluded from the calculation (see Methods).

Figure S10. Principal component analysis of PPS and NR trajectories (Apo and OM-bound states). A/C. Distribution of projections of PPS (solid lines) and NR (dashed lines) trajectories on PC2 (A) and PC3 (C). B/D. PPS (dots) and NR (crosses) trajectories of the Apo (green shades) and OM-bound (blue shades) simulations are projected onto PC1 (x axis) and PC2 (B) or PC 3 (D) (y axis). The position of initial structures is indicated with letters (n = Apo NR, N = OM-bound NR, p = Apo PPS, P = OM-bound PPS). PCs are calculated on the pseudo-trajectory obtained by concatenating all the 16 trajectories together. Highly flexible unstructured regions were excluded from the calculation (see Methods). The contribution of each PC to the total variance is reported in the axis label.

Figure S11. First three Principal Components of the concatenated pseudo-trajectory (PPS and NR, Apo and OM-bound states). In the porcupine representation, the orange spikes show the direction and relative amplitude of motion of each residue along the PC. Spikes shorter than 20% of the maximum length are not shown. Highly flexible unstructured regions were excluded from the calculation (see Methods).

Figure S12. Bending of the relay helix during the MD simulations of the PPS state. The relay helix bending was measured by calculating the angle formed by residues I478, H492 and E500 (C α atoms only). A. Probability distribution of the angle values observed during Apo (green) and OM-bound (blue) simulations. Arrows indicate the value of the angle measured for representative experimental structures of the pre-power stroke state (PDB IDs shown). B. Representative structures from PPS Apo (green) and OM-bound (blue) simulations. The energy-minimised experimental structure (PDB ID: 5N69) is shown as transparent white cartoon.

Figure S13. Profiles of $\Delta\zeta$ scores calculated for fragments in the OM-binding site. Plots of the difference between OM and Apo preferential connection scores ($\Delta\zeta$), calculated using fragments in the OM binding site as source sites. Similar profiles were observed when using source fragments close to each other, so that plots are shown only for selected fragments representative of the different profile types, namely K146, D168, H666 and R712 (orange dots). A negative value indicates a stronger preferential connection to the source fragment in OM-bound simulations compared to Apo ones. Negative bars are coloured according to their sub-domain. Grey areas highlight the position of functional regions, namely the seven strands of the central β -sheet (b1 to b7), the P-loop (P), the F helix (FH), Loop 1 (L1), the G helix (GH), Switch 1 (SW1), the β -bulge (bb), the OH loop (OH), Switch 2 (SW2), the Strut (St) and the SH1 helix (SH1).

Figure S14. Mapping of $\Delta\zeta$ scores calculated for fragments in the OM-binding site. Regions with negative values are coloured according to their sub-domain. The thickness of the tube is proportional to the magnitude of $\Delta\zeta$ values. Mapping is shown for K146 (A), D168 (B), H666 (C) and R712 (D). The position of the source fragment is indicated with an orange sphere. Highly flexible unstructured regions were excluded from the calculation (see Methods)

Figure S15. Pocket analysis in the nucleotide binding site. Pocket frequency maps calculated with MDpocket are represented as green (PPS Apo) and blue (PPS OM-bound) isosurfaces connecting points with frequency = 0.3 for each replica. The discontinuity between the Pi-containing pocket and the backdoor I channel observed in the OM-bound simulations and in Apo MD4 is indicated with an orange arrow. The position of the residues lining the inner and outer backdoor I tube is indicated in magenta, while the approximate position of the tube is indicated as a dashed magenta arrow in Apo MD1-3.

Figure S16. Inter-residue contacts in the backdoor I region. Backdoor I residues that are in contact (distance < 4 Å) for at least 50% of the trajectory are connected by an edge. Pocket frequency maps calculated with MDpocket are represented as green (PPS Apo) and blue (PPS OM-bound) isosurfaces connecting points with frequency = 0.3. The position of the residues lining the inner and outer backdoor I tube is indicated in magenta in the cartoon representation. The narrower backdoor I tubes observed in OM-bound simulations seem to be correlated with the presence of S180-E466, R243-A463 and R237-E466 contacts (orange edges).