

1 **Evaluating aziridinyl nitrobenzamide compounds as**
2 **leishmanicidal prodrugs**

3
4 **Andrew A. Voak¹, Karin Seifert², Nuala A. Helsby³ and Shane R. Wilkinson¹.**

5
6 ¹Queen Mary Pre-Clinical Drug Discovery Group, School of Biological and Chemical
7 Sciences, Queen Mary University of London, London, UK.

8 ²Immunology and Infection Department, London School of Hygiene and Tropical Medicine,
9 London, UK

10 ³Department of Molecular Medicine and Pathology, University of Auckland, Private Bag
11 92019, Auckland, New Zealand.

12
13
14 **corresponding author:** s.r.wilkinson@qmul.ac.uk

15
16
17
18
19
20
21 **Running title:** *Identifying anti-parasitic nitroaromatic compounds*

22 **Present address:** Andrew Voak, Immunology and Infection Department, London School of
23 Hygiene and Tropical Medicine, London, United Kingdom

25 **Abstract**

26 Many of the nitroaromatic agents used in medicine function as prodrugs and must undergo
27 activation before exerting their toxic effects. In most cases this is catalysed by FMN-
28 dependent type I nitroreductases (NTRs), a class of enzyme absent from higher eukaryotes
29 but expressed by bacteria and several eukaryotic microbes including trypanosomes and
30 *Leishmania*. Here, we utilize this difference to evaluate whether a library of aziridinyl
31 nitrobenzamides have activity against *Leishmania major*. Biochemical screens using purified
32 *L. major* NTR (LmNTR) revealed that compounds containing an aziridinyl-2,4-dinitrobenzyl
33 core were effective substrates for the enzyme and showed that the 4-nitro group was
34 important for this activity. To facilitate drug screening against intracellular amastigote
35 parasites, we generated leishmanial cells that expressed the luciferase reporter gene and
36 optimized a mammalian infection model in a 96-well plate format. A subset of aziridinyl-2,4-
37 dinitrobenzyl compounds possessing a 5-amide substituent displayed significant growth
38 inhibitory properties against the parasite, with the most potent agents generating 50 %
39 inhibitory concentrations of <100 nM towards the intracellular form. This antimicrobial
40 activity was shown to be LmNTR specific since *L. major* NTR^{+/-} heterozygote parasites were
41 slightly resistance to the most aziridinyl dinitrobenzyl agents tested. When the most potent
42 leishmanicidal agents were screened against the mammalian cells in which the amastigote
43 parasites were propagated, no growth inhibitory effect was observed at concentration up to
44 100 μM. We conclude that the aziridinyl nitrobenzamides represent a new lead structure that
45 may have the potential to treat leishmanial infections.

46


47 **Introduction**

48 Leishmaniasis represents a series of insect transmitted, blood-borne diseases caused by
49 more than 20 different protozoan parasite species belonging to the genus *Leishmania*. These
50 infections are endemic throughout many tropical and sub-tropical countries where 350
51 million people are at risk of infection (1). Estimates indicate that up to 12 million individuals
52 are currently infected by these protozoan parasites, with up to 2 million new cases and 50,000
53 deaths occurring each year (1). Recently, due to military activity, population migration,
54 modern medical practices, intravenous drug usage and global warming the number of new
55 cases in non-endemic areas has increased stimulating interest from pharmaceutical companies
56 in these previously neglected infections (2-4). Drugs currently represent the only treatments
57 available to combat leishmaniasis. For more than 60 years, front-line therapies have been
58 based on pentavalent antimonial compounds but their use is problematic as they are toxic and
59 require medical supervision to administer with clinical resistance now commonplace (5,6). In
60 light of this worrying situation, a range of alternative treatments such as amphotericin B,
61 paromomycin and miltefosine are now available but these too are far from ideal as they can
62 be expensive and require medical administration, with some having teratogenic and other
63 unwanted toxicity problems (7). Therefore, there is an urgent requirement for new, safer, cost
64 effective anti-leishmanial treatments.

65 Nitroaromatic compounds are used predominantly as broad spectrum antibiotics to treat
66 various urinary and gastrointestinal tract infections. They are characterised by possessing at
67 least one nitro group attached to an aromatic ring, that usually has a heterocyclic structure
68 (e.g. imidazole, furan or thiazole) (8). However, following concerns over their safety, the use
69 of many nitroaromatics has been discontinued in Europe and USA although they are
70 commonly prescribed elsewhere ((9-11); reviewed in (12)). It is now apparent that several
71 nitro-based compounds are not as toxic as initially thought (13-15). Such observations have



72 stimulated renewed interested in this group of agents with calls made for the reinstatement of
73 nitrofurantoin as a treatment for uncomplicated urinary tract infections, whilst several others
74 have emerged as lead structures to treat various microbial infections and different forms of
75 cancer, with fexinidazole and PA-824 undergoing evaluation to treat visceral leishmaniasis
76 (16-24).

77 Most nitroaromatic compounds used in medicine function as prodrugs and must undergo
78 activation before mediating their therapeutic effects, reactions catalysed by nitroreductases
79 (NTRs). Based on oxygen sensitivity, flavin cofactor and the reduction products, NTRs can
80 be broadly divided into two classes, type I or type II NTRs (25). In many bacteria and
81 eukaryotic microorganisms, FMN-containing type I NTRs use NAD(P)H to drive the
82 sequential two electron reduction of the conserved nitro (-NO₂) group to a hydroxylamine
83 derivative (-NHOH) via an unstable nitroso intermediate (-NO) (see reactions 1 and 2). The
84 hydroxylamine can be processed further to produce DNA damaging/crosslinking adducts and
85 other toxic molecules such as open chain nitriles and reactive dialdehydes (18,25-32).
86 Crucially, this reaction can occur in aerobic and anerobic environments indicating that the
87 above reactions do not involve O₂ and as such, FMN-type I catalysed activity is said to be
88 “O₂ insensitive”. For bacteria and trypanosomes, this mechanism of activation appears to be
89 key for the selective toxicity of many nitroaromatic prodrugs as cells selected for resistance
90 frequently show mutations in, or decreased expression of, their FMN-type I NTR gene
91 complement (33-35).



94 In contrast, the ubiquitous NAD(P)H dependent type II NTRs use FAD or FMN as a cofactor
95 to catalyse the one electron reduction of the conserved nitro group to form an unstable nitro
96 anion radical (-NO₂^{•-}). In the presence of O₂, this radical can undergo futile cycling to

97 produce superoxide anions (O_2^-) and regenerate the parental compound (36,37) (see reactions
98 3 and 4). Mammalian enzymes, such as NAD(P)H quinone oxidoreductase 1, can catalyse a
99 two electron reduction reaction of many nitroaromatic prodrugs under aerobic conditions,
100 however, mammalian nitroreduction does not utilise FMN containing -type I enzymes (38).



103 Here, we exploit the activity of a FMN-type I NTR expressed by the protozoan parasite
104 *Leishmania major* to conduct biochemical and phenotypic screens on a small library of
105 azirindyl nitrobenzamide (ANB) compounds, agents we have previously shown to display
106 significant anti-trypanosomal activities (39). Two of these compounds were highly active
107 against intracellular form *L major* and displayed high selectivity toward the parasite. We
108 postulate that compounds based around a 5-(aziridin-1-yl)-2,4-dinitrobenzamide core
109 represent a promising new class of leishmanicidal agent.

110

111 **Materials and methods**

112 **Chemicals**

113 Aziridinyl nitrobenzamides (ANB) structures are shown in Table 1. The synthesis of NH1-
114 8 is described elsewhere (40). CB1954 was purchased from Sigma Aldrich and NH9-12 were
115 supplied by the Department of Therapeutics, NCI (USA).

116

117 **Cell culturing.**

118 *L. major* (MHOM/IL/80/Friedlin) promastigote parasites were grown at 27 °C in modified
119 M199 medium (Life Technologies) (41). Transformed parasites were grown in this medium
120 supplemented with G418 (20 µg mL⁻¹ on agar plates, 40 µg mL⁻¹ in broth) or blasticidin (10
121 µg mL⁻¹). *L. major* metacyclic form parasites were harvested from promastigote cultures as
122 described (42). These were used to infect differentiated human acute monocytic leukemia
123 (THP-1) cells at a ratio of 20 parasites per mammalian cell. The *L. major*-infected
124 monolayers were incubated overnight at 37 °C under a 5 % (v/v) CO₂ atmosphere in
125 mammalian growth medium then washed with RPMI-1640 to remove residual parasites. *L.*
126 *major* amastigote parasites were maintained in differentiated THP-1 cells at 37 °C under a 5
127 % (v/v) CO₂ atmosphere in RPMI-1640 medium.

128 The human acute monocytic leukemia cell line (THP-1) was grown at 37 °C under a 5 %
129 (v/v) CO₂ atmosphere in RPMI-1640 medium (PAA Laboratories Ltd) supplemented with 2
130 mM pyruvate, 2 mM sodium glutamate, 2.5 U mL⁻¹ penicillin and 2.5 µg mL⁻¹ streptomycin,
131 20 mM HEPES pH 7.4 and 10% (v/v) foetal calf serum. Differentiation of THP-1 towards
132 macrophage-like cells was carried out using phorbol 12-myristate 13-acetate (20 ng mL⁻¹)
133 (PMA) (Sigma-Aldrich) (43,44).

134

135 ***In vivo* studies.**

136 All animal experiments were conducted under licence in accordance with UK Home
137 Office regulations. *L. major* parasites were passaged through female BALB/c mice by
138 subcutaneous injection of 2×10^7 purified metacyclic parasites in 100 μ L RPMI medium
139 without serum into the shaved rump. Amastigotes were harvested from skin lesions and
140 allowed to transform back to promastigotes in 5 ml M199 medium.

141

142 **Anti-proliferative assays.**

143 All assays were performed in a 96-well plate format. *L. major* promastigote parasites ($5 \times$
144 10^5 mL^{-1}) or differentiated THP-1 cells (2.5×10^4 mL^{-1}) were seeded in 200 μ l growth
145 medium containing different concentrations of nitroaromatic agent. After incubation at 27 °C
146 for 5 days (*L. major*) or at 37 °C for 3 days (THP-1), 2.5 μ g resazurin (20 μ L of 0.125 μ g mL^{-1}
147 stock in phosphate buffered saline) was added to each well and the plates incubated for a
148 further 8-16 hours. Cell densities were determined by monitoring the fluorescence of each
149 culture using a Gemini Fluorescent Plate Reader (Molecular Devices (UK) Ltd, Wokingham,
150 UK) at an excitation wavelength of 530 nm, emission wavelength of 585 nm and a filter cut
151 off at 550 nm, and the drug concentration that inhibits cell growth by 50 % (IC_{50}) established.

152 Growth inhibition of luciferase expressing *L. major* amastigotes was monitored as follows.
153 THP-1 cells seeded at 2.5×10^4 mL^{-1} in 200 μ l in growth medium containing PMA (20 ng
154 mL^{-1}) were incubated at 37 °C in a 5 % (v/v) CO_2 atmosphere for 3 days. Macrophage
155 monolayers were washed with mammalian growth medium then infected with purified
156 luciferase expressing *L. major* metacyclic cells (5×10^5 cells mL^{-1}) resuspended in 200 μ L
157 mammalian growth medium. Following incubation overnight at 37 °C in a 5 % (v/v) CO_2
158 atmosphere, the cultures were washed twice in growth medium to remove non-internalised

159 parasites and the supernatant replaced with fresh growth medium containing the compound
160 under investigation. Compound treated infections were incubated for a further 3 days at 37 °C
161 under a 5 % (v/v) CO₂. The growth medium was then removed and the cells lysed in 50 µL
162 cell culture lysis reagent (Promega). Activity was then measured using the luciferase assay
163 system (Promega) and light emission measured on a β-plate counter (Perkin Elmer). The
164 luminescence is proportional to the number of live cells. The IC₅₀ value for each compound
165 was then established.

166

167 **Plasmids and parasite genetic manipulation.**

168 DNA fragments corresponding to the *L. major* 5' rRNA spacer/promoter and 3' spacer
169 rRNA regions were amplified from *L. major* Friedlin genomic DNA and sequentially cloned
170 into the *Trypanosoma cruzi* vector pTRIX-Luc (39) replacing the equivalent *T. cruzi* 5' rRNA
171 spacer/promoter and 3' spacer rRNA sequences. A polypyrimidine tract/spliced leader
172 addition site sequence located upstream the *T. cruzi* *MPX* gene was then isolated and inserted
173 between the *L. major* 5' rRNA spacer/promoter region and luciferase reporter to form the
174 integrative vector pLmRIX-Luc. Following linearization, this DNA was introduced into *L.*
175 *major* promastigotes in the logarithmic phase of growth using the Human T-cell
176 Nucleofector[®] kit and an Amaxa[®] Nucleofector[™] (Lonza AG) set to program U-033.

177 A DNA fragment encoding for the catalytic domain of *L. major* *NTR* (*LmNTR*) was
178 amplified from genomic DNA with the primers *ggatcc*CTCGACGCCGTCGAGGCCGTCG
179 and *gaattc*CTAGAACTTGTTCCACCGCAC; lower-case italics correspond to restriction
180 sites incorporated into the primers to facilitate cloning. The fragment was digested with
181 *Bam*HI/*Hind*III then cloned into the corresponding sites of the vector pTrcHis-C (Invitrogen)
182 to form the plasmid pTrcHisC-LmNTR.

183

184 **Enzyme assay.**

185 Recombinant HIS-tagged LmNTR was purified as described (45). Type I NTR activity was
186 measured spectrophotometrically by monitoring the formation of the 2- and 4-hydroxylamine
187 derivatives from the parent ANB ($\lambda = 420 \text{ nm}$, $\epsilon = 1,220 \text{ M}^{-1} \text{ cm}^{-1}$) (46). A standard reaction
188 (1 mL) containing 50 mM Tris-Cl pH7.5, 100 μM NADH and 100 μM electron acceptor was
189 incubated at room temperature for 5 min. The background reaction rate was determined and
190 the assay initiated by addition of the LmNTR (35 μg). For nifurtimox, activity was measured
191 spectrophotometrically by monitoring reduction of this compound ($\lambda = 435 \text{ nm}$, $\epsilon = 19,000$
192 $\text{M}^{-1} \text{ cm}^{-1}$) (47). Enzyme activities were expressed in nmoles of NADH oxidised per minute
193 per mg LmNTR and assumes that four NADH molecules are oxidised per molecule of ANB
194 or nifurtimox reduced (46, 47).

195

196 **Results**

197 **Construction and evaluation of luciferase expressing *Leishmania major***

198 Several drug screening systems are now in place for use with *Leishmania*, each having
199 their own advantages and disadvantages. To facilitate screening of nitroaromatic compounds,
200 which are often coloured, we developed a *L. major* line that constitutively expresses
201 luciferase. The integrative vector pLmRIX-Luc was generated by sequentially cloning DNA
202 fragments containing the 5' and 3' *L. major* Friedlin rRNA promoter/spacer sequences either
203 side of an expression cassette derived from pTEX containing the luciferase and neomycin
204 phosphotransferase genes (39). To assist luciferase mRNA processing, an untranslated
205 sequence corresponding to the *T. cruzi* mitochondrial peroxiredoxin gene polypyrimidine
206 tract/spliced leader addition site was then inserted between the 5' *L. major* rRNA region and
207 reporter gene. The DNA fragment containing *L. major* rRNA/luciferase/neomycin
208 phosphotransferase sequences was then purified following restriction digestion and
209 electroporated into promastigote form parasites. After selection with G418, the luciferase
210 activity from several clones was determined and shown to be up to 1,000-fold higher than
211 that of the parental line (Fig. 1A). The effect of luciferase expression on various *L. major* life
212 cycle stages was then evaluated. This showed that the reporter did not influence: (i) growth of
213 promastigote parasites, (ii) the ability of promastigote cells to differentiate into infective
214 metacyclic forms, (iii) invasion of tissue culture derived macrophages by metacyclics, or (iv)
215 growth of intracellular amastigote parasites (Fig. 1B and E) . Additionally the luciferase
216 expressing parasites were passaged through BALB/c mice and amastigote recovered from
217 infected animals could readily differentiate to the promastigote form and be grown in culture.
218 Therefore, it is implicit that luciferase expression has no effect on *L. major* growth,
219 differentiation, and infectivity.

220 A prerequisite for any drug screening assay requires that reporter levels provide an
221 accurate representation of cell number. Using extracts derived from serially diluted
222 promastigotes, a linear relationship between luciferase activity and parasite load was
223 observed over the range of 625 to 160,000 cells (Fig. 1C). In the case of amastigotes such a
224 correlation was shown by evaluating reporter levels in lysates from a fixed number of
225 macrophages infected for 16-20 h with varying *L. major* loads (Fig. 1D): luciferase activity
226 was linear when using a parasite:mammalian cell ratio of 2.5-80:1. With this established the
227 luciferase expressing cells were then appraised as to whether they could be used for *in vitro*
228 drug screens in a 96-well plate format using nifurtimox as the reference compound. For
229 promastigote assays, cells were grown in the presence of different concentrations of the
230 nitrofurans for 6 days and the extracts tested for luminescence activity (Fig. 2A). From the
231 resultant dose response curves nifurtimox exhibited an IC_{50} value of $7.50 \pm 0.40 \mu\text{M}$, in line
232 with the IC_{50} value ($6.30 \pm 0.10 \mu\text{M}$) obtained when using the fluorescent vital dye resazurin.
233 For the intracellular amastigote assays, differentiated THP-1 cells were infected with
234 recombinant *L. major* at a ratio of 20 parasites per mammalian cell. Following 72 hours
235 growth post-infection in the presence of drug, the luciferase activity for each culture
236 determined from which dose response curves were generated (Fig. 2B). From these plots, the
237 IC_{50} value for nifurtimox was calculated to be $0.58 \pm 0.06 \mu\text{M}$. When the susceptibility of
238 differentiated THP-1 cells to nifurtimox was determined using resazurin, the IC_{50} value was
239 calculated to be $>100 \mu\text{M}$ (Table 2) demonstrating that this nitrofuran has a selective toxicity
240 of >170 toward the intracellular parasite. Therefore, based on our data, nifurtimox shows
241 selective killing of the *L. major* amastigotes and the observed leishmanicidal activity is not
242 due to death of the mammalian cell.

243

244 **Metabolism of nitrobenzamides by the leishmanial NTR.**

245 Many nitroaromatic prodrugs undergo activation in reactions catalysed by FMN-type I
246 NTRs. Here, we evaluated whether CB1954 and twelve related ANBs could function as
247 substrates for purified HIS-tagged *L. major* NTR using NADH as an electron donor (Fig. 3).
248 Normally, assays involving this parasite enzyme can be readily followed by monitoring the
249 change in absorbance at 340 nm, corresponding to the oxidation of NADH. However, many
250 ANBs undergo a considerable change in absorbance at this particular wavelength. Instead,
251 enzyme activity was assayed by monitoring the change in absorbance at 420 nm which
252 corresponds to the appearance of the hydroxylamine metabolite (46).

253 Of the thirteen compounds screened, most (ten) were metabolised by LmNTR at a rate
254 faster than that noted for nifurtimox. These “active” compounds were related in that they all
255 contain two nitro groups at positions 2- and 4- relative to the 1-aziridinyl ring. In contrast, the
256 three ANBs that contain a single 2-nitro group were deemed “poor” substrates for the parasite
257 enzyme, having negligible activity values.

258

259 **Leishmanicidal activity and mammalian cytotoxicity of aziridinyl nitrobenzamides.**

260 To determine whether there was a correlation between LmNTR activity and anti-
261 leishmanial activity, all compounds were screened against *L. major* promastigotes and
262 luciferase expressing amastigotes. Initial tests using a fixed concentration of compound (10
263 μ M) were set up to evaluate the growth inhibitory properties of the ANB series. Out of the
264 thirteen compounds tested, five had no effect on either parasite stage at this concentration
265 (Table 2) while two others had no effect on promastigote cells but did have activity against
266 intracellular form *L. major*. Three of the five compounds that have no effect against
267 promastigote and amastigote parasites correspond to the ANBs previously identified as
268 “poor” LmNTR substrates. To evaluate the growth inhibitory activities of those structures

269 identified by the initial screen, a series of secondary assays were performed using various
270 concentrations of the ANB. For each agent dose response curves were drawn from which the
271 compound's IC_{50} was determined (Table 2) (Fig. 4). Out of the six compounds tested against
272 promastigotes, CB1954, NH10 and NH11 had IC_{50} values $<1 \mu M$. These, plus NH9, also
273 displayed sub-micromolar IC_{50} 's against amastigote parasites with two (CB1964 and NH11)
274 yielding values $<100 \text{ nM}$. All compounds that display leishmanicidal activity against either of
275 the *L. major* forms were assayed for cytotoxicity against differentiated THP-1 cells (Table 2)
276 (Fig. 4). In all cases, the ANBs screened had no growth inhibitory effect against this
277 mammalian line at concentrations up to $100 \mu M$. Comparison of the mammalian cell IC_{50}
278 with the equivalent value obtained against amastigote parasites revealed that CB1964 and
279 NH11 displayed >2000 and >1667 -fold selective toxicity toward the intracellular pathogen,
280 respectively. This clearly demonstrates that the anti-parasitic activity displayed by these two
281 compounds was specifically due to growth inhibition of *L. major* and not due to induction of
282 host cell toxicity.

283 To demonstrate that LmNTR played a role in prodrug activation in the parasite itself, the
284 susceptibility of *LmNTR*^{+/-} heterozygous promastigotes to CB1954, NH10 and NH11 was
285 evaluated: attempts (16 independent transformations) to generate *L. major* *LmNTR*^{-/-} null
286 mutants failed to produce recombinant cells leading us to speculate that this activity is
287 essential in insect stage parasites (48). This screening demonstrated that cells expressing
288 lower levels of the oxidoreductase were between 2- and 4-fold more resistant to the ANB
289 than controls: wild type *L. major* had IC_{50} values of $0.42 \pm 0.01 \mu M$, $0.49 \pm 0.04 \mu M$ and
290 $0.91 \pm 0.07 \mu M$ against CB1954, NH10 and NH11, respectively, while *LmNTR*^{+/-}
291 heterozygotes displayed IC_{50} values of $1.62 \pm 0.02 \mu M$, $1.00 \pm 0.07 \mu M$ and $1.77 \pm 0.01 \mu M$
292 towards these agents.

293

294 **Discussion**

295 Several high throughput phenotypic screening approaches are now available to facilitate
296 the search for novel drugs targeting the medically relevant, intracellular stage of *Leishmania*.
297 Of these, the luciferase-based systems have proven to be sensitive, rapid, reproducible and
298 versatile with tagged parasites now being used to follow the fate of the pathogen during the
299 course of an animal model infection (49-52). Here, we constructed a luciferase expressing *L.*
300 *major* line making use of an existing *T. cruzi* integration vector, modifying the expression
301 construct such that it would integrate into the leishmanial ribosomal array (39).
302 Characterisation of the recombinant line established that expression of the reporter had no
303 effect on *L. major* promastigote growth, metacyclogenesis and proliferation of the
304 intracellular amastigote form. Using luciferase expressing *L. major*, growth inhibition assays
305 were established with nifurtimox employed as selective compound for promastigote or
306 amastigote cells in a standardized 96-well plate format.

307 Nitrobenzamide-based compounds that contain an aziridinyl ring or mustard substituent
308 have been evaluated as potential therapies targeting hypoxic cancers and the trypanosomal
309 infections African sleeping sickness and Chagas' disease (53-55). These structures invariably
310 function as prodrugs with their toxicity dependent on reduction of the nitro group(s) to its
311 hydroxylamine derivatives, reactions catalysed by FMN-type I NTRs (18,21). This
312 bioreductive trigger promotes an electronic reconfiguration on the compound's aromatic ring
313 leading to presentation of cytotoxic moieties to the cell (55). For the treatment of hypoxic
314 cancers, the FMN-type I NTR activity must be introduced into mammalian cells using gene-
315 or antibody-based approaches before addition of the nitrobenzamide whereas in
316 trypanosomes an essential endogenous enzyme can be exploited to catalyse the above nitro
317 reduction (18,21,55). As *Leishmania* also express a FMN-type I NTR, it is postulated that

318 nitroaromatic compounds may have potential for the treatment of various forms of
319 leishmaniasis (24).

320 Following the identification that CB1954, the archetypal ANB, displayed potent anti-
321 tumor activity against the Walker 256 carcinoma, a series of derivatives have been
322 synthesised differing in the number/location of nitro groups and other substituents attached to
323 a central benzyl ring core (40,56). Using a small library (14 compounds) of these structures in
324 biochemical screens, we demonstrated that compounds having two nitro groups located at the
325 2- and 4- positions on an aromatic ring backbone relative to the aziridinyl ring were readily
326 metabolised by LmNTR. The location of an amide (or amine)-containing substituent at the 5-
327 (group Ia) or 6- (group II) position did not affect this *in vitro* activity. In contrast,
328 replacement of the 4-nitro group with an H or SO₂Me substituent (group Ib) generated
329 compounds that were not metabolized by LmNTR. These biochemical studies revealed that
330 LmNTR exhibits a slightly different substrate preference compared to *Trypanosoma brucei*
331 NTR (TbNTR). In contrast to LmNTR the trypanosomal enzyme was only able to metabolise
332 compounds having the group Ia configuration. This highlights the importance of the 4-nitro
333 group during prodrug activation with both LmNTR and TbNTR unable to metabolise
334 compounds with a group Ib arrangement (39).

335 When tested against *Leishmania*, all group Ia structures showed potent growth inhibitory
336 properties toward promastigote and/or amastigote parasites thus mirroring the biochemical
337 observations. Several of these ANBs yielded IC₅₀'s below 1 μM against the intracellular *L.*
338 *major*, including CB1954 and NH11 that had sub 100 nM values. In contrast, none of the
339 group Ib ANBs had an effect on parasite growth, consistent with observations made using the
340 trypanosomal NTRs and mammalian cells expressing *E. coli* FMN-type I NTRs, again
341 highlighting the importance of the 4-nitro group and its reduction products make in mediating
342 cytotoxicity (39,40,57). Surprisingly, of the group II compounds only NH6 displayed any

343 leishmanicidal activity. NH6 had a lower potency compared with its structural isomer
344 CB1954 (CB1954 was 42- and 16-fold more effective at inhibiting amastigote and
345 promastigote parasite growth than NH6, respectively). The decreased potency displayed by
346 group II compounds may reflect the ability of these structures (or their reduction products) to
347 access regions of the cell where activation (or downstream leishmanicidal processes) take
348 place. This could be in part due to the spatial arrangement of these substituents (located at the
349 6- position on the benzyl ring) which may hinder the presentation of the adjacent azirindinyl
350 cytotoxic moiety to biomolecular targets within the cellular environment.

351 To conclusively demonstrate the link between LmNTR and leishmanicidal activities, the
352 susceptibility of *L. major* *LmNTR*^{+/-} heterozygous cell line to the most effective compounds
353 (CB1954, NH10 and NH11) was investigated (49). In this context, parasites with lower levels
354 of NTR displayed relative resistance to all compounds tested, mirroring observations made
355 for the trypanocidal, NTR-activated nitroaromatic prodrugs (35,39,45,47). This, in
356 conjunction with the observation that none of the group Ia compounds displayed cytotoxicity
357 to the macrophage-like cells within which the intracellular *L. major* were cultured, suggests
358 that the group Ia ANBs specifically target the parasite itself rather than by promoting death to
359 the mammalian cell.

360 We have now identified two leishmanicidal ANB-based compounds (CB1954 and NH11)
361 that display significant potency toward the *L. major* forms that replicate inside the
362 mammalian host. These structures were shown to mediate their anti-parasitic activities
363 following specific activation in a reaction catalysed by a FMN-type I NTR, an activity
364 present in the pathogen but absent from the mammalian macrophage cells. Indeed these
365 compounds display little or no cytotoxicity to the mammalian macrophage-like cell having
366 >1660-fold selectivity when targeting intracellular parasites although when using *in vivo*
367 rodent models CB1954 does exhibit hepatotoxicity, neurotoxicity and causes gastrointestinal

tract disturbances (58, 59): the toxicity/pharmacokinetics of other ANBs screened here have not been determined. Despite this group Ia compounds, particularly CB1954 and NH11, warrant further attention in the development of novel leishmanicidal therapies and ideally represent one component of a new combinatorial treatment.

372

373 **Acknowledgments**

374 Andrew Voak was a recipient of a Queen Mary University of London Graduate Training
375 Studentship

376

377 **References**

- 378 1. **Stuart, K., R. Brun, S. Croft, A. Fairlamb, R. E. Gurtler, J. McKerrow, S. Reed,**
379 **and R. Tarleton.** 2008. Kinetoplastids: related protozoan pathogens, different
380 diseases. *J. Clin. Invest.* **118**: 1301-1310.
- 381 2. **Cruz, I., C. Canavate, J. M. Rubio, M.A. Morales, C. Chicharro, F. Laguna, M.**
382 **Jimenez-Mejias, G. Sirera, S. Videla, and J. Alvar.** 2002. *Leishmania* in discarded
383 syringes from intravenous drug users. *Trans. R. Soc. Trop. Med. Hyg.* **96 Suppl 1**:
384 S185-189.
- 385 3. **Gonzalez, C., O. Wang, S. E. Strutz, C. Gonzalez-Salazar, V. Sanchez-Cordero,**
386 **and S. Sarkar.** 2010. Climate change and risk of leishmaniasis in North America:
387 predictions from ecological niche models of vector and reservoir species. *PLoS Negl.*
388 *Trop. Dis.* **4**: e585.
- 389 4. **Pavli, A., and H.C. Maltezos.** 2010. Leishmaniasis, an emerging infection in
390 travelers. *Int. J. Infect. Dis.* **14**: e1032-1039.
- 391 5. **Croft, S. L., S. Sundar, and A. H. Fairlamb.** 2006. Drug resistance in leishmaniasis.
392 *Clin. Microbiol. Rev.* **19**: 111-126.
- 393 6. **Haldar, A.K., P. Sen, and S. Roy.** 2011. Use of antimony in the treatment of
394 leishmaniasis: current status and future directions. *Mol. Biol. Int.* **2011**, 571242.
- 395 7. **Croft, S. L., and P. Olliaro.** 2011. Leishmaniasis chemotherapy - challenges and
396 opportunities. *Clin. Microbiol. Infect.* **17**: 1478-1483.
- 397 8. **Grunberg, E., and E. H. Titsworth.** 1973. Chemotherapeutic properties of
398 heterocyclic compounds: monocyclic compounds with five-membered rings. *Annu.*
399 *Rev. Microbiol.* **27**: 317-346.
- 400 9. **Takahashi, M., S. Iizuka, T. Watanabe, M. Yoshida, J. Ando, K. Wakabayashi,**
401 **and A. Maekawa.** 2000. Possible mechanisms underlying mammary carcinogenesis
402 in female Wistar rats by nitrofurazone. *Cancer Lett.* **156**: 177-184.
- 403 10. **Padda, R. S., C. Wang, J. B. Hughes, R. Kutty, and G. N. Bennett.** 2003.
404 Mutagenicity of nitroaromatic degradation compounds. *Environ. Toxicol. Chem.* **22**:
405 2293-2297.

- 406 11. **Hiraku, Y., A. Sekine, H. Nabeshi, K. Midorikawa, M. Murata, Y. Kumagai, and**
407 **S. Kawanishi.** 2004. Mechanism of carcinogenesis induced by a veterinary
408 antimicrobial drug, nitrofurazone, via oxidative DNA damage and cell proliferation.
409 *Cancer Lett.* **215**: 141-150.
- 410 12. **McCalla, D. R.** 1983. Mutagenicity of nitrofurans derivatives: review. *Environ.*
411 *Mutagen.* **5**: 745-765.
- 412 13. **Matsumoto, M., H. Hashizume, T. Tomishige, M. Kawasaki, H. Tsubouchi, H.**
413 **Sasaki, Y. Shimokawa, and M. Komatsu.** 2006. OPC-67683, a nitro-dihydro-
414 imidazooxazole derivative with promising action against tuberculosis in vitro and in
415 mice. *PLoS Med.* **3**: e466.
- 416 14. **Tweats, D., B. Bourdin Trunz, and E. Torrele.** 2012. Genotoxicity profile of
417 fexinidazole - a drug candidate in clinical development for human African
418 trypanomiasis (sleeping sickness). *Mutagenesis* **27**: 523-532.
- 419 15. **Trunz, B. B., R. Jedrysiak, D. Tweats, R. Brun, M. Kaiser, J. Suwinski, and E.**
420 **Torrele.** 2011. 1-Aryl-4-nitro-1H-imidazoles, a new promising series for the
421 treatment of human African trypanosomiasis. *Eur. J. Med. Chem.* **46**: 1524-1535.
- 422 16. **Stover, C. K., P. Warrener, D. R. VanDevanter, D. R. Sherman, T. M. Arain, M.**
423 **H. Langhorne, S. W. Anderson, J. A. Towell, Y. Yuan, D. N. McMurray, B. N.**
424 **Kreiswirth, C. E. Barry, and W. R. Baker.** 2000. A small-molecule
425 nitroimidazopyran drug candidate for the treatment of tuberculosis. *Nature* **405**: 962-
426 966.
- 427 17. **Adagu, I. S., D. Nolder, D. C. Warhurst, and J. F. Rossignol.** 2002. *In vitro*
428 activity of nitazoxanide and related compounds against isolates of *Giardia*
429 *intestinalis*, *Entamoeba histolytica* and *Trichomonas vaginalis*. *J. Antimicrob.*
430 *Chemother.* **49**: 103-111.
- 431 18. **Denny, W. A.** 2003. Prodrugs for Gene-Directed Enzyme-Prodrug Therapy (Suicide
432 Gene Therapy). *J. Biomed. Biotechnol.* **2003**: 48-70.
- 433 19. **Korba, B. E., A. B. Montero, K. Farrar, K. Gaye, S. Mukerjee, M. S. Ayers, and**
434 **J. F. Rossignol.** 2008. Nitazoxanide, tizoxanide and other thiazolides are potent
435 inhibitors of hepatitis B virus and hepatitis C virus replication. *Antiviral. Res.* **77**: 56-
436 63.
- 437 20. **Arya, S. C., and N. Agarwal.** 2009. Nitrofurantoin: the return of an old friend in the
438 wake of growing resistance. *BJU Int.* **103**: 994-995.
- 439 21. **Chen, Y., and L. Hu.** 2009. Design of anticancer prodrugs for reductive
440 activation. *Med. Res. Rev.* **29**, 29-64.
- 441 22. **Torrele, E., B. Bourdin Trunz, D. Tweats, M. Kaiser, R. Brun, G. Mazue, M. A.**
442 **Bray, and B. Pecoul.** 2010. Fexinidazole-a new oral nitroimidazole drug candidate
443 entering clinical development for the treatment of sleeping sickness. *PLoS Negl.*
444 *Trop. Dis.* **4**: e923.
- 445 23. **Wyllie, S., S. Patterson, L. Stojanovski, F. R. Simeons, S. Norval, R. Kime, K. D.**
446 **Read, and A. H. Fairlamb.** 2012. The anti-trypanosome drug fexinidazole shows
447 potential for treating visceral leishmaniasis. *Sci. Transl. Med.* **4**: 119re111.
- 448 24. **Patterson, S., S. Wyllie, L. Stojanovski, M. R. Perry, F. R. Simeons, S. Norval,**
449 **M. Osuna-Cabello, M. De Rycker, K. D. Read, and A. H. Fairlamb.** 2013 The R
450 enantiomer of the anti-tubercular drug PA-824 as a potential oral treatment for
451 visceral leishmaniasis. *Antimicrob. Agents. Chemother.* in press.
452 doi:10.1128/AAC.00722-13

- 453 25. **Peterson, F. J., R. P. Mason, J. Hovsepien, and J. L. Holtzman.** 1979. Oxygen-
454 sensitive and -insensitive nitroreduction by *Escherichia coli* and rat hepatic
455 microsomes. *J. Biol. Chem.* **254**: 4009-4014.
- 456 26. **Gavin, J. J., F. F. Ebetino, R. Freedman, and W. E. Waterbury.** 1966. The aerobic
457 degradation of 1-(5-nitrofurfurylideneamino)-2-imidazolidinone (NF-246) by
458 *Escherichia coli*. *Arch. Biochem. Biophys.* **113**: 399-404.
- 459 27. **McCalla, D. R., A. Reuvers, and C. Kaiser.** 1971. Breakage of bacterial DNA by
460 nitrofurans derivatives. *Cancer Res.* **31**, 2184-2188.
- 461 28. **Whitmore, G. F., A. J. Varghese, and S. Gulyas.** 1986. Reaction of 2-
462 nitroimidazole metabolites with guanine and possible biological consequences. *IARC*
463 *Sci. Publ.*, 185-196.
- 464 29. **Whitmore, G. F., and A. J. Varghese.** 1986. The biological properties of reduced
465 nitroheterocyclics and possible underlying biochemical mechanisms. *Biochem.*
466 *Pharmacol.* **35**: 97-103.
- 467 30. **Streeter, A. J., and B. A. Hoener.** 1988. Evidence for the involvement of a nitrenium
468 ion in the covalent binding of nitrofurazone to DNA. *Pharm. Res.* **5**: 434-436.
- 469 31. **Hall, B. S., C. Bot, and S. R. Wilkinson.** 2011. Nifurtimox activation by
470 trypanosomal type I nitroreductases generates cytotoxic nitrile metabolites. *J. Biol.*
471 *Chem.* **286**: 13088-13095.
- 472 32. **Hall, B. S., and S.R. Wilkinson.** 2012. Activation of benznidazole by trypanosomal
473 type I nitroreductases results in glyoxal formation. *Antimicrob. Agents. Chemother.*
474 **56**: 115-123.
- 475 33. **McCalla, D. R., C. Kaiser, and M. H. Green.** 1978. Genetics of nitrofurazone
476 resistance in *Escherichia coli*. *J. Bacteriol.* **133**: 10-16.
- 477 34. **Whiteway, J., P. Koziarz, J. Veall, N. Sandhu, P. Kumar, B. Hoecher, and I. B.**
478 **Lambert.** 1998. Oxygen-insensitive nitroreductases: analysis of the roles of *nfsA* and
479 *nfsB* in development of resistance to 5-nitrofurans derivatives in *Escherichia coli*. *J.*
480 *Bacteriol.* **180**: 5529-5539.
- 481 35. **Wilkinson, S. R., M. C. Taylor, D. Horn, J. M. Kelly, and I. Cheeseman.** 2008. A
482 mechanism for cross-resistance to nifurtimox and benznidazole in trypanosomes.
483 *Proc. Natl. Acad. Sci. USA* **105**: 5022-5027.
- 484 36. **Moreno, S. N., R. P. Mason, R. P. Muniz, F. S. Cruz, and R. Docampo.** 1983.
485 Generation of free radicals from metronidazole and other nitroimidazoles by
486 *Trichomonas foetus*. *J. Biol. Chem.* **258**: 4051-4054.
- 487 37. **Moreno, S. N., R. P. Mason, and R. Docampo.** 1984. Reduction of nifurtimox and
488 nitrofurantoin to free radical metabolites by rat liver mitochondria. Evidence of an
489 outer membrane-located nitroreductase. *J. Biol. Chem.* **259**: 6298-6305.
- 490 38. **Chen, S., R. Knox, K. Wu, P. S. Deng, D. Zhou, M. A. Bianchet, and L. M.**
491 **Amzel.** 1997. Molecular basis of the catalytic differences among DT-diaphorase of
492 human, rat, and mouse. *J. Biol. Chem.* **272**: 1437-1439.
- 493 39. **Bot, C., B. S. Hall, N. Bashir, M. C. Taylor, N. A. Helsby, and S. R. Wilkinson.**
494 2010. Trypanocidal activity of aziridinylnitrobenzamide prodrugs. *Antimicrob.*
495 *Agents Chemother.* **54**: 4246-4252.
- 496 40. **Helsby, N. A., G. J. Atwell, S. Yang, B. D. Palmer, R. F. Anderson, S. M. Pullen,**
497 **D. M. Ferry, A. Hogg, W. R. Wilson, and W. A. Denny.** 2004.
498 Aziridinyldinitrobenzamides: synthesis and structure-activity relationships for
499 activation by *E. coli* nitroreductase. *J. Med. Chem.* **47**: 3295-3307.


- 500 41. **Kapler, G. M., C. M. Coburn, and S. M. Beverley.** 1990. Stable transfection of the
501 human parasite *Leishmania major* delineates a 30-kilobase region sufficient for
502 extrachromosomal replication and expression. *Mol. Cell. Biol.* **10**:1084-1094.
- 503 42. **da Silva, R., and D. L. Sacks.** 1987. Metacyclogenesis is a major determinant of
504 *Leishmania* promastigote virulence and attenuation. *Infect. Immun.* **55**: 2802-2806.
- 505 43. **Fleit, H. B., and C. D. Kobasiuk.** 1991. The human monocyte-like cell line THP-1
506 expresses Fc gamma RI and Fc gamma RII. *J. Leukoc. Biol.* **49**: 556-565.
- 507 44. **Tsuchiya, S., M. Yamabe, Y. Yamaguchi, Y. Kobayashi, T. Konno, and K. Tada.**
508 1980. Establishment and characterization of a human acute monocytic leukemia cell
509 line (THP-1). *Int. J. Cancer* **26**: 171-176.
- 510 45 **Hall, B. S., X. Wu, L. Hu, and S. R. Wilkinson.** 2010. Exploiting the drug-
511 activating properties of a novel trypanosomal nitroreductase. *Antimicrob. Agents*
512 *Chemother.* **54**: 1193-1199.
- 513 46. **Race, P. R., A. L. Lovering, S. A. White, J. I. Grove, P. F. Searle, C. W.**
514 **Wrighton, and E. I. Hyde.** 2007. Kinetic and structural characterisation of
515 *Escherichia coli* nitroreductase mutants showing improved efficacy for the prodrug
516 substrate CB1954. *J. Mol. Biol.* **368**: 481-492.
- 517 47. **Bot, C., B. S. Hall, G. Alvarez, R. Di Maio, M. Gonzalez, H. Cerecetto, and S. R.**
518 **Wilkinson.** 2013. Evaluating 5-nitrofurans as trypanocidal agents. *Antimicrob.*
519 *Agents Chemother.* **57**: 1638-1647.
- 520 48. **Voak, A. A., V. Gobalakrishnapilla, K. Seifert, E. Balczon, L. Hu, B. S. Hall, and**
521 **S. R. Wilkinson.** 2013. An essential type I nitroreductase from *Leishmania major* can
522 be used to activate leishmanicidal prodrugs. *J. Biol. Chem.* in press. doi:
523 10.1074/jbc.M113.494781
- 524 49. **de La Llave, E., H. Lecoecur, A. Besse, G. Milon, E. Prina, and T. Lang.** 2011. A
525 combined luciferase imaging and reverse transcription polymerase chain reaction
526 assay for the study of *Leishmania* amastigote burden and correlated mouse tissue
527 transcript fluctuations. *Cell. Microbiol.* **13**: 81-91.
- 528 50. **Michel, G., B. Ferrua, T. Lang, M. P. Maddugoda, P. Munro, C. Pomares, E.**
529 **Lemichez, and P. Marty.** 2011. Luciferase-expressing *Leishmania infantum* allows
530 the monitoring of amastigote population size, *in vivo*, *ex vivo* and *in vitro*. *PLoS Negl.*
531 *Trop. Dis.* **5**: e1323.
- 532 51. **Thalhofer, C. J., J. W. Graff, L. Love-Homan, S. M. Hickerson, N. Craft, S. M.**
533 **Beverley, and M. E. Wilson.** 2010. *In vivo* imaging of transgenic *Leishmania*
534 parasites in a live host. *J. Vis Exp.*
- 535 52. **Roy, G., C. Dumas, D. Sereno, Y. Wu, A. K. Singh, M. J. Tremblay, M.**
536 **Ouellette, M. Olivier, and B. Papadopoulou.** 2000. Episomal and stable expression
537 of the luciferase reporter gene for quantifying *Leishmania* spp. infections in
538 macrophages and in animal models. *Mol. Biochem. Parasitol.* **110**: 195-206.
- 539 53. **Jameson, M. B., D. Rischin, M. Pegram, J. Gutheil, A. V. Patterson, W. A.**
540 **Denny, and W. R. Wilson.** 2010. A phase I trial of PR-104, a nitrogen mustard
541 prodrug activated by both hypoxia and aldo-keto reductase 1C3, in patients with solid
542 tumors. *Cancer Chemother. Pharmacol.* **65**: 791-801.
- 543 54. **Patel, P., J. G. Young, V. Mautner, D. Ashdown, S. Bonney, R. G. Pineda, S. I.**
544 **Collins, P. F. Searle, D. Hull, E. Peers, J. Chester, D. M. Wallace, A. Doherty, H.**
545 **Leung, L. S. Young, and N. D. James.** 2009. A phase I/II clinical trial in localized
546 prostate cancer of an adenovirus expressing nitroreductase with CB1954. *Mol. Ther.*
547 **17**: 1292-1299.

- 548 55. **Wilkinson, S. R., C. Bot, J. M. Kelly, and B. S. Hall.** 2011. Trypanocidal activity of
549 nitroaromatic prodrugs: current treatments and future perspectives. *Curr. Top. Med.*
550 *Chem.* **11**: 2072-2084.
- 551 56. **Cobb, L. M., T. A. Connors, L. A. Elson, A. H. Khan, B. C. Mitchley, W. C. Ross,**
552 **and M. E. Whisson.** 1969. 2,4-dinitro-5-ethyleneiminobenzamide (CB 1954): a
553 potent and selective inhibitor of the growth of the Walker carcinoma 256. *Biochem.*
554 *Pharmacol.* **18**: 1519-1527.
- 555 57. **Helsby, N. A., D. M. Ferry, A. V. Patterson, S. M. Pullen, and W. R. Wilson.**
556 2004. 2-Amino metabolites are key mediators of CB 1954 and SN 23862 bystander
557 effects in nitroreductase GDEPT. *Br. J. Cancer* **90**: 1084-1092.
- 558 58. **Chung-Faye, G., D. Palmer, D. Anderson, J. Clark, M. Downes, J. Baddeley, S.**
559 **Hussain, P. I. Murray, P. Searle, L. Seymour, P. A. Harris, D. Ferry, and D. J.**
560 **Kerr.** 2001. Virus-directed, enzyme prodrug therapy with nitroimidazole reductase: a
561 phase I and pharmacokinetic study of its prodrug, CB 1954. *Clin. Cancer Res.* **7**:
562 2662-2666
- 563 59. **Tang, M.H., N. A. Helsby, M. A. Goldthorpe, K. M. Thompson, S. Al-Ali, and M.**
564 **D. Tingle.** 2007. Hepatic nitroreduction, toxicity and toxicokinetics of the anti-tumour
565 prodrug CB 1954 in mouse and rat. *Toxicology.* **240**: 70-85.
- 566

567

Tables

568

Table 1: Structure of aziridinyl nitrobenzamide compounds.

Compound	Group	Structure
CB1954	Ia	R1=NO ₂ ; R2=CONH ₂ ; R3=H
NH1	Ia	R1=NO ₂ ; R2=CONH(CH ₂) ₂ Nmorpholide; R3=H
NH2	Ia	R1=NO ₂ ; R2=CONH(CH ₂) ₂ CO ₂ Me; R3=H
NH9	Ia	R1=NO ₂ ; R2=oxazole; R3=H
NH10	Ia	R1=NO ₂ ; R2=CONHCH ₂ CHCH ₂ ; R3=H
NH11	Ia	R1=NO ₂ ; R2=CONH(CH ₃)CH ₃ ; R3=H
NH12	Ia	R1=NO ₂ ; R2=R3=H
NH3	Ib	R1=H; R2=R2=CONH ₂ ; R3=H
NH4	Ib	R1=SO ₂ Me; R2=CONH ₂ ; R3=H
NH5	Ib	R1=SO ₂ Me; R2=NHCH ₂ CH(OH)CH ₂ OH; R3=H
NH6	II	R1=NO ₂ ; R2=H; R3=CONH ₂
NH7	II	R1=NO ₂ ; R2=H; R3=NHCH ₂ CH(OH)CH ₂ OH
NH8	II	R1=NO ₂ ; R2=H; R3=CONH(CH ₂) ₂ Nmorpholide

569

570

571 **Table 2: Susceptibility of *L. major* and THP-1 cells to aziridinyl nitrobenzamides^a**

compounds	group	<i>L. major</i> IC ₅₀ (μM)		differentiated THP-1 IC ₅₀ (μM) ^c	selective toxicity ^d
		promastigotes ^b	amastigotes ^c		
nifurtimox		7.50 ± 0.40	0.58 ± 0.06	>100	>172
CB1954	Ia	0.42 ± 0.01	0.05 ± 0.02	>100	>2000
NH1	Ia	7.32 ± 0.63	11.15 ± 2.01	>100	>9
NH2	Ia	1.48 ± 0.18	2.95 ± 1.14	>100	>34
NH9	Ia	>10	0.67 ± 0.05	>100	>149
NH10	Ia	0.49 ± 0.04	0.56 ± 0.05	>100	>178
NH11	Ia	0.91 ± 0.07	0.06 ± 0.01	>100	>1667
NH12	Ia	>10	1.32 ± 0.02	>100	>76
NH3-5	Ib	>10	>10	nd	nd
NH6	II	6.85 ± 0.18	2.10 ± 0.22	>100	>48
NH7-8	II	>10	>10	nd	nd

572

573 ^and, not determined. ^bData are means from 4 experiments ± standard deviation. ^cData are
574 means from 3 experiments ± standard deviation. ^dThe therapeutic index of a compound was
575 calculated as a ratio of the IC₅₀ against differentiated THP-1 cells to the IC₅₀ against
576 amastigote parasites. Nifurtimox was used as a reference compounds.

577

Figure 1




Figure 1. Luciferase expression in insect- and mammalian-stage *L. major*. (A). The luciferase activity, in relative light units, of six recombinant *L. major* promastigote clones (Luc1-6) was determined and compared to that of the parental line (wt). 1×10^6 cells were used in each analysis. (B). *L. major* wild type (square; solid line) and *LmNTR^{+/-}BLA* heterozygote (diamond; dotted line) promastigote parasites growth was monitored until cultures were in the stationary phase of growth. From day 5 onward, the number of metacyclic form parasites in wild type (triangle; solid line) and *LmNTR^{+/-}BLA* heterozygote (triangle; dotted line) promastigote culture was determined following purification by agglutination. The data is expressed as % metacyclics load in the total *L. major* population. All curves shown are derived from a single data set and are representative of experiments performed in triplicate. (C). Purified *L. major* metacyclic parasites engineered to express luciferase were used to infect differentiated THP-1 cells. Over a 4 day post infection period, extracts were generated from each cell line and the luciferase activity determined. Following background correction, the luciferase activity was plotted against time. All curves shown are derived from a single data set and are representative of experiments performed in triplicate. (D). Correlation between promastigote load (between 625 to 160,000 cells) and luciferase activity. Three independent readings were taken for each parasite load and the values are means \pm standard deviation. (E). Correlation between luciferase activity and *L. major* amastigote load using various parasite/macrophage cell ratios (0:1 to 80:1).

Figure 2




Figure 2. Susceptibility of *L. major* to nifurtimox. Dose response curves of luciferase expressing *L. major* promastigotes (A) and amastigotes (B) to nifurtimox were determined from which IC_{50} values were calculated. The data are the means from three independent experiments \pm standard deviation.

Figure 3




Figure 3. Activity of LmNTR toward different aziridinyl nitrobenzamides. (A) SDS-PAGE gel (10 %) stained with Coomassie blue. Lane 1, size standards; lane 2, purified HIS-tagged LmNTR. (B) The activity of purified recombinant LmNTR was assessed by using various ANBs as substrates (100 μ M) at a fixed concentration of NADH (100 μ M). Enzyme activity, expressed in nmoles of NADH oxidised per minute per mg LmNTR ($\text{nmol.mg}^{-1}\text{min}^{-1}$), was then calculated using an ϵ value of $1,220 \text{ M}^{-1} \text{ cm}^{-1}$ with the assumption that four molecules of NADH are oxidised per molecule of ANB reduced (46). NFX (nifurtimox) was used as control. For nifurtimox, the change in absorbance at 435 nm was followed and enzyme activity determined using an ϵ value of $19,000 \text{ mM}^{-1}\text{cm}^{-1}$, again assuming that four molecules of NADH were oxidised per molecule of nitrofurantoin reduced (47). The enzyme activity values are the means of data from three assays \pm standard deviations. The ANB group designations as listed in Table 1 are noted.

Figure 4




Figure 4. Dose response curves of *L. major* and mammalian cells to aziridinyl nitrobenzamides. Various concentrations of leishmanicidal ANBs were tested against *L. major* and THP-1 cells. The growth inhibitory effect of each treatment was evaluated and dose response curves constructed for promastigotes (●), amastigotes (■) and differentiated THP-1 cells (▲). In all cases, drug treatments were performed in triplicate and the plots shown represent the average growth inhibition obtained at each concentration \pm standard deviation. The curves for compounds CB1954, NH6 and NH10 are shown.