



***DOCK6* Mutations are Responsible for a Distinct Autosomal Recessive Variant of Adams-Oliver Syndrome Associated with Brain and Eye Anomalies**

Journal:	<i>Human Mutation</i>
Manuscript ID:	humu-2014-0525.R1
Wiley - Manuscript type:	Brief Report
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Sukalo, Maja; University Hospital Magdeburg, Institute of Human Genetics Tilsen, Felix; University Hospital Magdeburg, Institute of Human Genetics Kayserili, Hulya; Istanbul Medical Faculty, Medical Genetics Department Müller, Dietmar; Klinikum Chemnitz, Institut für Medizinische Genetik Tuysuz, Beyhan; Istanbul University, Department of Pediatric Genetics Ruddy, Deborah; Guy's Hospital, Department of Clinical Genetics Wakeling, Emma; North West London Hospitals NHS Trust, North West Thames Regional Genetics Service Ørstavik, Karen; Oslo University Hospital, Department of Medical Genetics Snape, Katie; St George's Healthcare NHS Trust, Department of Clinical Genetics Trembath, Richard; Queen Mary University of London, Barts and The London School of Medicine and Dentistry De Smedt, Maryse; Leuven University Hospital, Department of Medical Genetics van der Aa, Nathalie; Antwerp University Hospital, Department of Medical Genetics Skalej, Martin; University Hospital Magdeburg, Institute of Neuroradiology Mundlos, Stefan; Universitätsmedizin Berlin and Max Planck Institute for Molecular Genetics Berlin, Institute for Medical and Human Genetics Charité Wuyts, Wim; Antwerp University Hospital, Department of Medical Genetics; University of Antwerp, Department of Medical Genetics Southgate, Laura; Queen Mary University of London, Barts and The London School of Medicine and Dentistry; King's College London, Division of Genetics and Molecular Medicine Zenker, Martin; University Hospital Magdeburg, Institute of Human Genetics</p>
Key Words:	Adams-Oliver syndrome, <i>DOCK6</i> , autosomal recessive, brain anomalies, eye anomalies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review

BRIEF REPORT***DOCK6* Mutations are Responsible for a Distinct Autosomal Recessive Variant of Adams-Oliver Syndrome Associated with Brain and Eye Anomalies**

Maja Sukalo^{1†*}, Felix Tilsen^{1†}, Hülya Kayserili², Dietmar Müller³, Beyhan Tüysüz⁴, Deborah M. Ruddy⁵, Emma Wakeling⁶, Karen Helene Ørstavik⁷, Katie M. Snape⁸, Richard Trembath^{5,9}, Maryse De Smedt¹⁰, Nathalie van der Aa¹¹, [Martin Skalej](#)¹², Stefan Mundlos¹³, Wim Wuyts^{11,14#}, Laura Southgate^{9,15#}, Martin Zenker^{1#}

† These authors contributed equally as first authors

These authors contributed equally as senior authors

¹Institute of Human Genetics, University Hospital Magdeburg, Magdeburg, Germany

²Medical Genetics Department, Istanbul Medical Faculty, Istanbul, Turkey

³Institut für Medizinische Genetik, Klinikum Chemnitz, Chemnitz, Germany

⁴Department of Pediatric Genetics, Istanbul University, Istanbul, Turkey

⁵Department of Clinical Genetics, Guy's Hospital, London, UK

⁶North West Thames Regional Genetics Service, North West London Hospitals NHS Trust, Harrow, UK

⁷Department of Medical Genetics, Oslo University Hospital, Oslo, Norway

⁸Department of Clinical Genetics, St George's Healthcare NHS Trust, London, UK

⁹Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK

¹⁰Department of Medical Genetics, Leuven University Hospital, Leuven, Belgium

¹¹Department of Medical Genetics, Antwerp University Hospital, Antwerp, Belgium

¹²[Institute of Neuroradiology, University Hospital Magdeburg, Magdeburg, Germany](#)

1
2
3 ¹³Institute for Medical and Human Genetics Charité, Universitätsmedizin Berlin and Max
4
5 Planck Institute for Molecular Genetics Berlin, Berlin, Germany

6
7 ¹⁴Department of Medical Genetics, University of Antwerp, Antwerp, Belgium

8
9 ¹⁵Division of Genetics and Molecular Medicine, King's College London, London, UK

10
11 Contract grant sponsor: German Research Foundation, DFG ZE 524/2-3

12
13
14
15
16 *Corresponding Author:

17
18 Maja Sukalo

19
20 Institute of Human Genetics

21
22 University Hospital Magdeburg

23
24 Leipziger Straße 44

25
26 39120 Magdeburg

27
28
29 Germany

30
31 E-mail: Maja.Sukalo@med.ovgu.de

Abstract

Adams-Oliver syndrome (AOS) is characterized by the association of aplasia cutis congenita with terminal transverse limb defects, often accompanied by additional cardiovascular or neurological features. Both autosomal dominant and recessive disease transmission have been observed, with recent gene discoveries indicating extensive genetic heterogeneity. Mutations of the *DOCK6* gene were first described in autosomal recessive cases of AOS and only five *DOCK6*-related families have been reported to date. Recently, a second type of autosomal recessive AOS has been attributed to *EOGT* mutations in three consanguineous families. Here, we describe the identification of 13 *DOCK6* mutations, the majority of which are novel, across 10 unrelated individuals from a large cohort comprising 47 sporadic cases and 31 AOS pedigrees suggestive of autosomal recessive inheritance. *DOCK6* mutations were strongly associated with structural brain abnormalities, ocular anomalies, and intellectual disability, thus suggesting that *DOCK6*-linked disease represents a variant of AOS with a particularly poor prognosis.

Keywords: Adams-Oliver syndrome, *DOCK6*, autosomal recessive, brain anomalies, eye anomalies.

1
2
3 First described in 1945, Adams-Oliver syndrome (AOS) is characterized by the
4 combination of terminal transverse limb defects (TTLD) and aplasia cutis congenita (ACC)
5 typically located in the midline parietal and/or occipital region of the scalp [Adams and
6 Oliver, 1945]. Structures underlying these defects (skull bones, meninges, sinus) may also be
7 involved. AOS is often associated with additional congenital vascular anomalies such as cutis
8 marmorata telangiectatica congenita (CMTC), reported in around 20% of patients, pulmonary
9 hypertension, and lesions of presumed vascular etiology in other organs. Moreover, around
10 20% of patients with AOS have congenital cardiac defects including – amongst others – aortic
11 valve anomalies, septal defects and tetralogy of Fallot [Snape et al., 2009]. The spectrum of
12 congenital anomalies observed in AOS has led to the hypothesis that disturbed vasculogenesis
13 may underlie this disorder [Swartz et al., 1999]. AOS is emerging as a very heterogeneous
14 disorder, both clinically and genetically. To date, three genes have already been identified as
15 causative for the autosomal dominant form, namely *ARHGAP31* (MIM *610911; AOS1;
16 #100300) [Southgate et al., 2011], *RBPJ* (MIM *147183; AOS3; #614814) [Hassed et al.,
17 2012], and *NOTCH1* (MIM *190198; AOS5; #616028) [Stittrich et al., 2014]. Two genes,
18 *DOCK6* (MIM *614194; AOS2; #614219) [Shaheen et al., 2011] and *EOGT* (MIM *614789;
19 AOS4; #615297) [Shaheen et al., 2013], have been reported in pedigrees with autosomal
20 recessive transmission of AOS. Each of these genes apparently accounts for only a minor
21 proportion of patients. It is therefore likely that further AOS genes will be identified in the
22 future.

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Homozygous or compound heterozygous *DOCK6* mutations have so far been reported in only four inbred Arab families [Shaheen et al., 2011, 2013] and in a single sporadic patient [Lehman et al., 2014], respectively. The *DOCK6* protein belongs to the conserved dedicator of cytokinesis family and has a role in remodeling the actin cytoskeleton by acting as a guanine nucleotide exchange factor (GEF) for two members of the Rho GTPase family, Cdc42 and Rac1 [Miyamoto et al., 2007]. This regulation of Cdc42 and Rac1 complements

1
2
3 the GTPase-activating protein (GAP) activity of the gene product of *ARHGAP31*
4 [Tcherkezian et al., 2006], mutations of which underlie some autosomal dominant cases of
5 AOS [Southgate et al., 2011; Isrie et al., 2014], thus pointing at abnormal cytoskeleton
6 remodeling as one of the basic pathogenic mechanisms leading to AOS.
7
8
9

10
11 To further understand the role of *DOCK6* in the etiology of this disorder and to
12 establish possible phenotype correlations, we performed a comprehensive mutation screen of
13 this gene in a large and heterogeneous patient cohort. The study cohort consisted of 88
14 patients from 78 unrelated families recruited by the partners of the AOS Collaborative Group.
15 The presence of both ACC and TTLD in at least one affected family member served as
16 minimal clinical inclusion criteria for this study, with the exception of one case that has been
17 previously published as a variant of AOS with cognitive impairment, but without scalp defect
18 [Brancati et al., 2008]. Additional physical abnormalities were reported in a considerable
19 proportion of patients and included cerebral (n=19), ocular (n=13), neurodevelopmental
20 (n=25), and cardiac anomalies (n=14). Either sporadic cases of AOS (n=47) or those with a
21 pedigree constellation suggestive of autosomal recessive disease transmission (n=31) were
22 included. Parental consanguinity and/or the presence of multiple affected children of
23 clinically unaffected parents were regarded as possible indicators of autosomal recessive
24 inheritance. Families with parent-child transmission of the phenotype suggesting autosomal
25 dominant inheritance were excluded. *DOCK6* mutation screening was performed by PCR and
26 conventional sequencing of all 48 coding exons and flanking intronic regions (Supp. Materials
27 and Methods). The study was approved by the institutional review boards of the participating
28 centers, University of Magdeburg/Erlangen, Guy's and St Thomas' Hospitals London, and
29 University of Antwerp. Written informed consent was obtained from the patients and/or the
30 parents. Mutations, unclassified variants and phenotype data were submitted to the Leiden
31 Open Variation Database (<http://databases.lovd.nl/>).
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 In this cohort we detected 10 unrelated individuals with biallelic sequence changes in
4 *DOCK6* that were classified as probable pathogenic mutations. Seven of those patients were
5 offspring of consanguineous parents, two originated from non-consanguineous families with
6 multiple affected children, and one was a sporadic case with no known parental consanguinity
7 (Supp. Figure S1). The overall proportion of *DOCK6*-related AOS across our complete cohort
8 was 13%, with a frequency of 29% (9/31) among the families suggestive of autosomal
9 recessive inheritance and 2% (1/47) in sporadic cases with no parental consanguinity. Our
10 findings thus underscore the importance of *DOCK6* as a gene for autosomal recessive AOS.
11 They also suggest that a small proportion of apparently sporadic cases are in fact recessive
12 with *DOCK6* as the underlying etiology.
13
14
15
16
17
18
19
20
21
22
23
24

25 The mutations observed in these 10 families included nonsense (n=1), missense (n=4),
26 frameshift (n=4), and splice site mutations (n=3), as well as one larger intragenic deletion-
27 insertion resulting in deletion of exons 42 to 47. The latter was identified through the failure
28 to amplify the terminal exons by PCR and confirmed by focused MLPA and breakpoint
29 sequencing (Supp. Figure S1, family 9). Eleven of these 13 mutations were novel and two
30 have been previously described as causative of AOS [Shaheen et al., 2011, 2013] (Figure 1A
31 and Supp. Table S1). Seven index patients had homozygous mutations consistent with self-
32 stated parental consanguinity, while the remaining three had compound heterozygous
33 changes. Of the four missense mutations observed in this cohort, three were homozygous in
34 affected children from consanguineous families (c.3047T>C, p.Leu1016Pro; c.3154G>A,
35 p.Glu1052Lys; c.4786C>T, p.Arg1596Trp) and one (c.788T>A, p.Val263Asp) occurred in
36 compound heterozygosity with a splice site mutation on the second allele. All four missense
37 variations were classified as likely causative mutations on the basis of conservation of the
38 affected residue, as assessed by various online prediction tools (Supp. Table S2). Moreover, in
39 the consanguineous family harboring the missense mutation p.Leu1016Pro (c.3047T>C,
40 family 1), previous homozygosity mapping using a SNP array had been consistent with
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 linkage to the *DOCK6* locus in the index patient, demonstrating a 22 Mb stretch of
4
5 autozygosity on chromosome 19 (data not shown). In one pedigree (family 6), segregation of
6
7 compound heterozygosity for the missense mutation p.Val263Asp and a splice site mutation
8
9 on the second allele (c.5939+2T>C) was confirmed in the two affected siblings (Table 1,
10
11 Supp. Figure S1). Of the three splice site mutations observed in this study, one
12
13 (c.4106+5G>T) is outside of the canonical splice site dinucleotide. Unfortunately, no
14
15 appropriate material could be obtained to prove the splicing effect on the mRNA level.
16
17 However, compound heterozygosity for this change and a frameshift mutation on the other
18
19 allele was found to segregate with the phenotype in family 7 (Table 1, Supp. Figure S1).
20
21 Furthermore, splice prediction tools consistently calculated that this change likely abrogated
22
23 splice donor function at this site (Supp. Table S3), thus supporting the likely pathogenic role
24
25 of this variation. To date, six distinct *DOCK6* mutations have been reported to underlie the
26
27 AOS type 2 (Figure 1A, Supp. Table S1), with loss of function or expression of the *DOCK6*
28
29 protein suggested as the basic pathogenic mechanism [Shaheen et al., 2011, 2013; Lehman et
30
31 al., 2014]. Taken together with previous reports, this study demonstrates that *DOCK6*
32
33 mutations are distributed over the entire gene with no obvious clustering to certain domains of
34
35 the encoded protein (Figure 1A). A deleterious effect on the gene product is plausible for
36
37 most of these changes, as they are predicted to lead to either a truncated protein or nonsense-
38
39 mediated mRNA decay. However, the precise functional consequences of the novel missense
40
41 mutations presented here remain to be explored.
42
43
44
45
46

47 In addition to the pathogenic mutations described above, we also identified 16
48
49 heterozygous *DOCK6* sequence variations in our cohort, which remained as unclassified due
50
51 to either uncertain clinical significance or annotation in dbSNP (build 139) as rare variants
52
53 (MAF <0.01) (Supp. Table S4). These variants included predicted amino acid substitutions
54
55 (n=8), synonymous alterations in the coding sequence (n=5), and intronic substitutions within
56
57 20 bp of the splice site (n=3). None of these variations were unambiguously classified as
58
59
60

1
2
3 disease-causing by prediction tools. Thirteen unrelated sporadic cases harbored a single
4
5 heterozygous unclassified *DOCK6* variant, while two patients were found to have two or
6
7 more variants. Of these, one case had inherited both variants (c.885C>T, p.(=) and
8
9 c.2104G>A, p.Gly702Ser) from the mother on the same allele (data not shown). Another
10
11 patient was found to harbor three unclassified variants (c.885C>T, p.(=); c.1289G>A,
12
13 p.Arg430His; c.1833-19C>G), the segregation of which could not be studied. Notably, this
14
15 patient was previously reported in the literature as a variant subtype of AOS associated with
16
17 cerebral anomalies, seizures and severe MR, but without ACC of the scalp [Brancati et al.,
18
19 2008]. While most of these variations are more likely to be non-pathogenic (Supp. Table S5),
20
21 we cannot fully exclude any contribution to the observed phenotype. Our mutation screening
22
23 strategy did not assess mutations of the promoter and intronic changes. We also did not
24
25 systematically screen for larger genomic deletions/duplications. Therefore, it remains possible
26
27 that additional pathogenic variants may have been missed in this cohort and that the given
28
29 figure of the contribution of *DOCK6*-related disease is somewhat underestimated. However,
30
31 for the *DOCK6* mutation-negative patients originating from consanguineous families we can
32
33 state that five had a previous SNP array analysis showing no suggestive stretch of
34
35 homozygosity at the *DOCK6* locus (data not shown). In two out of four further subjects who
36
37 had no previous homozygosity mapping, *DOCK6* sequencing revealed at least one
38
39 heterozygous SNP, whilst for two cases, sequencing results were uninformative in excluding
40
41 homozygosity at the *DOCK6* gene locus. Thus, at least for our consanguineous families we
42
43 can conclude that genes other than *DOCK6* are very likely involved in the pathogenesis of
44
45 AOS. Mutations of the *EOGT* gene may account for part of our *DOCK6*-negative AOS cases
46
47 [Shaheen et al., 2013]; however mutation screening of this gene was not within the scope of
48
49 this study. It also remains to be seen whether further recessive AOS genes will be identified in
50
51 due course. Moreover, considering the inclusion criteria for this study, it is possible that a
52
53 proportion of our cohort may in fact represent dominant *de novo* mutations or, in the case of
54
55
56
57
58
59
60

1
2
3 affected siblings with asymptomatic parents, autosomal dominant inheritance with incomplete
4
5 penetrance.
6

7
8 The main clinical findings of the *DOCK6*-positive individuals from our cohort are
9
10 summarized in Table 1. Detailed clinical data could be obtained from 10 patients originating
11
12 from eight families. The patients' ages ranged between one week and 20 years (median 4.3
13
14 years). All except one affected individual from these families had ACC of the scalp and
15
16 TTLD of variable expression; patient 7.2 presented only with mild hypoplasia of toenails
17
18 along with a congenital heart defect, impaired vision and mild cognitive impairment, whereas
19
20 his sister presented with classic AOS features including ACC and TTLD. Across our *DOCK6*-
21
22 positive cohort, the limb defects ranged from minimal hypoplasia of terminal phalanges to
23
24 severe transverse reduction defects (Figure 1B). Notably, aside from ACC typically located
25
26 on the scalp vertex, four patients had additional areas of ACC on the abdomen. Further
27
28 associated anomalies, primarily related to the nervous system, were present in all individuals
29
30 carrying homozygous or compound-heterozygous *DOCK6* mutations. Specifically, all patients
31
32 from whom sufficient data could be obtained were reported with developmental delay or
33
34 mental retardation, ranging from mild to severe (Table 1). A broad range of additional
35
36 neurological abnormalities were reported in most cases, including cerebral palsy, spasticity,
37
38 contractures, and epilepsy. Only one patient aged ≥ 4 years had achieved the ability to walk
39
40 without support. Behavioral abnormalities including autistic behavior or temper tantrums
41
42 were reported in two patients. Brain MRI or CT had been performed for seven patients and
43
44 was abnormal in all cases. The most frequent changes observed on brain imaging included
45
46 ventriculomegaly, periventricular leukomalacia/calcifications, and hypoplasia/atrophy of the
47
48 corpus callosum (Table 1). Images from five affected individuals are exemplarily shown in
49
50 Figure 1C. Patient 4.1 underwent cerebral ultrasonography at 3 months of age which also
51
52 showed ventriculomegaly. A further patient (6.2) was previously reported with ventricular
53
54 dilatation, partial agenesis of the corpus callosum, and periventricular leukomalacia on
55
56
57
58
59
60

1
2
3 autopsy [Orstavik et al., 1995]. Where available, measurements of head circumference were
4
5 in the microcephalic range for all eight patients. Ocular anomalies including microphthalmia,
6
7 retinal detachment, and visual impairment were reported in all patients for whom clinical
8
9 information was obtainable. In contrast, cardiac anomalies were observed in only three cases.
10

11
12 Taken together, the most striking phenotypic attribute of *DOCK6*-related AOS in the
13
14 presented cohort is the strong association with important neurodevelopmental and ocular
15
16 anomalies. The pattern of neurological impairment and most of the reported morphological
17
18 changes (microcephaly, ventricular dilatation, periventricular calcifications, cortical changes)
19
20 are suggestive of a disruptive vascular pathogenesis rather than a primary maldevelopment of
21
22 the brain. Lesions classified as calcifications according to density analysis, may represent
23
24 primary calcifications but can in fact also have resulted from previous microbleeds. Likewise,
25
26 the main ocular anomalies observed in our *DOCK6*-positive patients, namely microphthalmia
27
28 and retinal detachment, are compatible with a disruptive vasculogenesis. The high prevalence
29
30 of brain and eye abnormalities as well as the pattern of cerebral and ocular involvement is in
31
32 line with previous case reports (Table 1). However the data on the previously reported
33
34 patients do not provide specific detail to definitely state that brain involvement is a constant
35
36 feature in AOS type 2. While *DOCK6* mutations are generally a rare cause of AOS, in our
37
38 cohort they accounted for 8/25 (32%) cases presenting with major neurodevelopmental
39
40 defects and for 9/19 (47%) cases with documented brain abnormalities. Taken together, these
41
42 data suggest that *DOCK6* mutations are particularly responsible for a variant of AOS
43
44 characterized by ACC, TTLD plus cerebral and ocular abnormalities. The existence of such a
45
46 variant was previously postulated nearly 20 years ago [Orstavik et al., 1995] and our study
47
48 now confirms that *DOCK6* is indeed the gene responsible for the disease in that family
49
50 (family 6). The strong association of *DOCK6* mutations with anomalies of the brain and eye
51
52 implies that deleterious effects on angiogenesis caused by *DOCK6* deficiency also affect
53
54 development of these particular structures. In their review, Snape et al. concluded that
55
56
57
58
59
60

1
2
3 abnormal brain and ocular findings are more common in autosomal recessive AOS [Snape et
4 al., 2009]. It is becoming clear that the individuals with *DOCK6* mutations account for a
5
6
7 substantial part for this observation. By contrast, among five patients with *EOGT* mutations,
8
9
10 only one patient was reported to have brain anomalies and no abnormal ocular findings were
11
12 reported in any subject [Shaheen et al., 2013]. **Nonetheless, across our complete cohort,**
13
14 **approximately two-thirds of the AOS patients with major neurodevelopmental disorders and**
15
16 **about half of the cases with structural brain anomalies could not be explained by *DOCK6***
17
18 **mutations, thus suggesting that the association with a neurological phenotype is not specific to**
19
20 **AOS type 2.**

21
22
23 In summary, by presenting 10 novel families with *DOCK6* mutations, we substantially
24
25 expand the clinical and mutational spectrum of AOS type 2. Our findings provide independent
26
27 corroboration that mutations in *DOCK6* are responsible for nearly one third of autosomal
28
29 recessively inherited AOS and that this genetic entity also accounts for a minority of sporadic
30
31 cases. AOS type 2 is particularly if not consistently associated with cerebral and ocular
32
33 anomalies in addition to ACC and TTLD. In patients with such a constellation of symptoms
34
35 *DOCK6* should therefore be the primary candidate gene for molecular investigation.

36 37 38 39 40 41 **Acknowledgements**

42
43 The authors thank the participating families and the clinicians who recruited patients
44
45 to the European AOS Consortium (M. Aglam, L. Al-Gazali, D. Amor, S. Beekman, F.
46
47 Brancati, G. Breisch-Lotz, W. Coerdts, B. De Vries, J. Desir, L. Fayol, L. Graul-Neumann, Z.
48
49 Gucev, U. Hehr, M. Holder, D. Horn, E. Jacquemin, S. Janssens, A. Kariminejad, E. Kinning,
50
51 Y. Lacassie, C. Li, V. López González, S. Lynch, L. McGregor, V. Meiner, I. Navarro, S.
52
53 Mercier, S. Odent, M. Patel, L. Pinson, K. Prescott, W. Reardon, C. Rusu, M. Splitt, C.
54
55 Tertilt, E. Thompson, J. Tolmie, J. van den Ende, L. van Maldergem, A. Weber, M.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Whiteford, M. Willems). We also thank Ilka Kramer, Susan Engelberg, and Nicole Epperlein for excellent technical support, and [J. Westvik for his kind collaboration](#).

For Peer Review


References

- Adams FH, Oliver CP. 1945. Hereditary deformities in man due to arrested development. *J Hered* 36:3-7.
- Brancati F, Garaci FG, Mingarelli R, Dallapiccola B. 2008. Abnormal neuronal migration defect in the severe variant subtype of Adams-Oliver syndrome. *Am J Med Genet A* 146A:1622-3.
- Cote JF, Vuori K. 2002. Identification of an evolutionarily conserved superfamily of DOCK180-related proteins with guanine nucleotide exchange activity. *J Cell Sci* 115:4901-13.
- Hassed SJ, Wiley GB, Wang S, Lee JY, Li S, Xu W, Zhao ZJ, Mulvihill JJ, Robertson J, Warner J, Gaffney PM. 2012. RBPJ mutations identified in two families affected by Adams-Oliver syndrome. *Am J Hum Genet* 91:391-5.
- Isrie M, Wuyts W, Van Esch H, Devriendt K. 2014. Isolated terminal limb reduction defects: extending the clinical spectrum of Adams-Oliver syndrome and ARHGAP31 mutations. *Am J Med Genet A* 164A:1576-9.
- Lehman A, Stittrich AB, Glusman G, Zong Z, Li H, Eydoux P, Senger C, Lyons C, Roach JC, Patel M. 2014. Diffuse angiopathy in Adams-Oliver syndrome associated with truncating DOCK6 mutations. *Am J Med Genet A* 164A:2656-62.
- Miyamoto Y, Yamauchi J, Sanbe A, Tanoue A. 2007. Dock6, a Dock-C subfamily guanine nucleotide exchanger, has the dual specificity for Rac1 and Cdc42 and regulates neurite outgrowth. *Experimental cell research* 313:791-804.
- Orstavik KH, Stromme P, Spetalen S, Flage T, Westvik J, Vesterhus P, Skjeldal O. 1995. Aplasia cutis congenita associated with limb, eye, and brain anomalies in sibs: a variant of the Adams-Oliver syndrome? *Am J Med Genet* 59:92-5.

- 1
2
3 Prothero J, Nicholl R, Wilson J, Wakeling EL. 2007. Aplasia cutis congenita, terminal limb
4 defects and falciform retinal folds: confirmation of a distinct syndrome of vascular
5 disruption. *Clin Dysmorphol* 16:39-41.
6
7
8
9
10 Shaheen R, Faqeih E, Sunker A, Morsy H, Al-Sheddi T, Shamseldin HE, Adly N, Hashem M,
11 Alkuraya FS. 2011. Recessive mutations in DOCK6, encoding the guanidine
12 nucleotide exchange factor DOCK6, lead to abnormal actin cytoskeleton organization
13 and Adams-Oliver syndrome. *Am J Hum Genet* 89:328-33.
14
15
16
17
18 Shaheen R, Aglan M, Keppler-Noreuil K, Faqeih E, Ansari S, Horton K, Ashour A, Zaki MS,
19 Al-Zahrani F, Cueto-Gonzalez AM, Abdel-Salam G, Temtamy S, et al. 2013.
20 Mutations in EOGT confirm the genetic heterogeneity of autosomal-recessive Adams-
21 Oliver syndrome. *Am J Hum Genet* 92:598-604.
22
23
24
25
26
27 Snape KM, Ruddy D, Zenker M, Wuyts W, Whiteford M, Johnson D, Lam W, Trembath RC.
28
29 2009. The spectra of clinical phenotypes in aplasia cutis congenita and terminal
30 transverse limb defects. *Am J Med Genet A* 149A:1860-81.
31
32
33
34 Southgate L, Machado RD, Snape KM, Primeau M, Dafou D, Ruddy DM, Branney PA,
35 Fisher M, Lee GJ, Simpson MA, He Y, Bradshaw TY, et al. 2011. Gain-of-function
36 mutations of ARHGAP31, a Cdc42/Rac1 GTPase regulator, cause syndromic cutis
37 aplasia and limb anomalies. *Am J Hum Genet* 88:574-85.
38
39
40
41
42
43 Stittrich AB, Lehman A, Bodian DL, Ashworth J, Zong Z, Li H, Lam P, Khromykh A, Iyer
44 RK, Vockley JG, Baveja R, Silva ES, et al. 2014. Mutations in NOTCH1 Cause
45 Adams-Oliver Syndrome. *Am J Hum Genet* 95:275-84.
46
47
48
49
50 Swartz EN, Sanatani S, Sandor GG, Schreiber RA. 1999. Vascular abnormalities in Adams-
51 Oliver syndrome: cause or effect? *Am J Med Genet* 82:49-52.
52
53
54
55
56
57
58
59
60 Tcherkezian J, Triki I, Stenne R, Danek EI, Lamarche-Vane N. 2006. The human orthologue
of CdGAP is a phosphoprotein and a GTPase-activating protein for Cdc42 and Rac1
but not RhoA. *Biol Cell* 98:445-56.

Figure Legend

Figure 1: (A) DOCK6 protein with known functional domains and distribution of mutations. The protein contains two DOCK homology regions, DHR-1 and DHR-2. DHR-1 spans about 200 amino acids at the N-terminal end of the protein, whereas DHR-2 is located towards the C-terminus and has an approximate length of 500 amino acids [Cote and Vuori, 2002]. All currently known mutations are displayed according to their location in the DOCK6 protein. Red represents nonsense mutations (n=3), black indicates frameshift mutations (n=5), missense mutations are shown in green (n=4), splice site mutations are colored in orange (n=4) and the blue line represents one large deletion insertion at the C-terminal end of the DOCK6 protein spanning exons 42 to 47. Novel mutations reported in this paper are written in italics. (B) Clinical photographs of three *DOCK6*-positive individuals with AOS from this cohort showing areas of alopecia on the vertex resulting from aplasia cutis congenita and terminal defects of the digits of varying severity. (C) Brain imaging of AOS patients with *DOCK6* mutations. Cranial MRI of **patient 2.1** at age 1 year: T2-weighted axial section showing enlarged lateral ventricles and cerebral atrophy particularly affecting the frontal lobe, and contrast enhanced T1-weighted median sagittal section illustrating thin corpus callosum and enlarged basal subarachnoid spaces. CT scan of **patient 3.1** at age 6 years: Axial sections showing ventriculomegaly and periventricular calcifications, and orbital section showing right microphthalmia with interocular hyperdensities representing retinal detachment and cystic malformation of the anterior chamber. T1-weighted MRI of **patient 5.1** at age 3 years: Axial sections showing irregularly shaped and slightly dilated lateral ventricles. Axial CT scan of **patient 6.1** in neonatal period showing ventricular dilatation and multiple periventricular calcifications. Brain imaging of **patient 10.1**: CT scan at age 2 years showing periventricular calcifications, and T2-weighted MRI axial section age 3 years showing irregularly shaped, slightly enlarged ventricles and mild atrophy of the brain. MRI, magnetic resonance imaging, CT, computed tomography.




(A) DOCK6 protein with known functional domains and distribution of mutations.

(B) Clinical photographs of three *DOCK6*-positive individuals with AOS from this cohort showing areas of alopecia on the vertex resulting from aplasia cutis congenita and terminal defects of the digits of varying severity.

(C) Brain imaging of AOS patients with *DOCK6* mutations.

[RGB color]

195x299mm (300 x 300 DPI)



(A) DOCK6 protein with known functional domains and distribution of mutations.
(B) Clinical photographs of three *DOCK6*-positive individuals with AOS from this cohort showing areas of alopecia on the vertex resulting from aplasia cutis congenita and terminal defects of the digits of varying severity.
(C) Brain imaging of AOS patients with *DOCK6* mutations.
 [CMYK color]
 195x299mm (300 x 300 DPI)

Table 1: Mutation and phenotype data of *DOCK6*-positive individuals from this cohort (families 1-10) compared to previously published cases (families 11-15).

Family	Patient	Mutations	Gender	Age	Parental consanguinity	Intra uterine growth restriction	Scalp defect	TTLDD [hands/feet]	Congenital heart defect	Brain anomalies	Microcephaly	Ocular anomalies	Cognitive impairment	Neurology	Additional features	Reference
1	1.1	[p.L1016P] + [p.L1016P]	F	5y	+	na	+	+/+	na	na	+	MO, RD, VO, ACA	DD	SE	high palate	-
2	2.1	[p.T455Sfs*24] + [c.4491+1G>A]	M	10y	-	-	+	+/+	na	VD/BA, CCH	+	NS	sev ID	SE, CP	CMTC, single umbilical artery, cryptorchidism	-
3	3.1	[p.Q434Rfs*21] + [p.Q434Rfs*21]	M	20y	+	+	+	+/+	-	VD/BA, PVL	+	MO, RD, ACA	sev ID	SE, CP	CMTC, abdominal skin defect	-
4	4.1	[p.R1596W] + [p.R1596W]	F	3m	+	-	+	+/+	PDA	VD/BA	+	MO	na	-	knee dislocation	-
5	5.1	[p.E1052K] + [p.E1052K]	M	9y	+	+	+	+/+	-	VD/BA, CCH, PVL	+	MO, RD	mod ID	SE	cryptorchidism	1
6	6.1	[p.V263D] + [c.5939+2T>C]	F	na	-	-	+	+/+	VSD	VD/BA, PVL	+	MO, RD, VO	sev ID	SE, CP	abdominal skin defects, absence of right patella	2
	6.2	[p.V263D] + [c.5939+2T>C] ^a	M	1w [†]	-	+	+	+/+	na	VD/BA, CCH	na	RD	na	na	abdominal skin defect, patella fixed to skin	2
7	7.1	[p.F635Pfs*32] + [c.4106+5G>T]	F	7y	-	-	+	+/+	-	NS	+	NS	sev ID	SE	abdominal skin defect	-
	7.2	[p.F635Pfs*32] + [c.4106+5G>T]	M	8y	-	+	-	-/+	TAPVD	na	na	NS	mild ID	-	hypothyroidism	-
8	8.1	[p.E162*] + [p.E162*]	F	na	+	na	+	+/+	na	na	na	na	na	na		-
9	9.1	[c.5235+205_6102-15delins10] + [c.5235+205_6102-15delins10]	F	7y	+	-	+	+/+	na	PVL	na	na	na	na		-
10	10.1	[p.R841Sfs*6] + [p.R841Sfs*6]	F	na	+	-	+	+/+	na	VD/BA, CCH, PVL	+	na	na	SE		-
11	11.1	[p.T455Sfs*24] + [p.T455Sfs*24]	F	11m	+	na	+	+/+	-	VD/BA, PVL	+	OA	sev ID	SE, CP		3
12	12.1	[p.D416*] + [p.D416*]	F	3.5y	+	na	+	+/+	-	na	+	-	DD	na		3
13	13.1	[p.R841Sfs*6] + [p.R841Sfs*6]	M	1y	+	na	+	+/+	AVD	VD/BA, PVL	na	-	na	na	abdominal skin defect	4
	13.2	[p.R841Sfs*6] + [p.R841Sfs*6]	F	na	+	na	+	+/+	na	VD/BA, PVL	na	na	na	SE	gastroschisis	4
14	14.1	[c.4107-1G>C] + [c.4107-1G>C]	F	2y	+	na	+	+/+	na	PVL, PA	na	OA	na	SE		4
15	15.1	[p.L1064Vfs*60] + [p.E1494*]	F	2y	-	+	+	+/+	TOF, PLSVC	PVL, PE	+	RD	sev ID	SE	placental vasculopathy, neonatal thrombocytopenia, small bowel infarction	5

1
2
3
4
5 ^aGenotype was not directly confirmed as patient is deceased but is assumed to be the same as in affected sibling.

6 F, female; M, male; y, year(s); m, month(s); w, weeks(s); †, deceased; na, no data available; +, present; -, not present; TTLD, terminal transverse
7 limb defects; PDA, patent ductus arteriosus; VSD, ventricular septal defect; TAPVD, total anomalous pulmonary venous connection; AVD, aortic
8 valve dysplasia; TOF, tetralogy of Fallot; PLSVC, persistent left superior vena cava; VD/BA, ventricular dilatation / brain atrophy; CCH, corpus
9 callosum hypoplasia/atrophy; PVL, periventricular lesions (calcification, gliosis); NS, abnormality present, not further specified; PA, pachygyria;
10 PE, porencephaly; MO, microphthalmia; RD, retinal detachment; VO, vitreous opacities/membranes; ACA, anterior chamber abnormality; OA,
11 optic atrophy; DD, developmental delay; ID, intellectual disability; sev, severe; mod, moderate; SE, seizures / epilepsy; CP, cerebral palsy /
12 spasticity; [CMTC, cutis marmorata telangiectatica congenita](#). References: (1) Prothero et al. (2007); (2) Orstavik et al. (1995); (3) Shaheen et al.
13 (2011); (4) Shaheen et al. (2013); (5) Lehman et al. (2014).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Sukalo et al. (2015), *Human Mutation*, Supporting Material

***DOCK6* Mutations are Responsible for a Distinct Autosomal Recessive Variant of Adams-Oliver Syndrome Associated with Brain and Eye Anomalies**

Maja Sukalo^{1†*}, Felix Tilsen^{1†}, Hülya Kayserili², Dietmar Müller³, Beyhan Tüysüz⁴, Deborah M. Ruddy⁵, Emma Wakeling⁶, Karen Helene Ørstavik⁷, Katie M. Snape⁸, Richard Trembath^{5,9}, Maryse De Smedt¹⁰, Nathalie van der Aa¹¹, Martin Skalej¹², Stefan Mundlos¹³, Wim Wuyts^{11,14#}, Laura Southgate^{9,15#}, Martin Zenker^{1#}

¹Institute of Human Genetics, University Hospital Magdeburg, Magdeburg, Germany. ²Medical Genetics Department, Istanbul Medical Faculty, Istanbul, Turkey. ³Institut für Medizinische Genetik, Klinikum Chemnitz, Chemnitz, Germany. ⁴Department of Pediatric Genetics, Istanbul University, Istanbul, Turkey. ⁵Department of Clinical Genetics, Guy's Hospital, London, UK. ⁶North West Thames Regional Genetics Service, North West London Hospitals NHS Trust, Harrow, UK. ⁷Department of Medical Genetics, Oslo University Hospital, Oslo, Norway. ⁸Department of Clinical Genetics, St George's Healthcare NHS Trust, London, UK. ⁹Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK. ¹⁰Department of Medical Genetics, Leuven University Hospital, Leuven, Belgium. ¹¹Department of Medical Genetics, Antwerp University Hospital, Antwerp, Belgium. ¹²Institute of Neuroradiology, Otto von Guericke University Magdeburg, Magdeburg, Germany. ¹³Institute for Medical and Human Genetics Charité, Universitätsmedizin Berlin and Max Planck Institute for Molecular Genetics Berlin, Berlin, Germany. ¹⁴Department of Medical Genetics, University of Antwerp, Antwerp, Belgium. ¹⁵Division of Genetics and Molecular Medicine, King's College London, London, UK. † These authors contributed equally as first authors. # These authors contributed equally as senior authors.

Supp. Material and Methods

We designed oligonucleotide primers for each of the 48 exons of *DOCK6* using the Primer3 software version 4.0.0 (<http://primer3.ut.ee/>) [Untergrasser et al., 2012]. Primer sequences and PCR conditions are available upon request. Sequencing was carried out using the BigDye Terminator Cycle Sequencing Kit v3.1 on an ABI 3500xl automated capillary sequencer (Applied Biosystems, Cheshire, UK). Obtained sequences were compared with the reference sequence (NM_020812.3) using the Sequence Pilot software v4.0.1 (JSI Medical Systems GmbH, Kippenheim, Germany). Pathogenicity of all observed sequence variants was assessed using various online prediction tools. For splice site prediction we utilized the following bioinformatic tools:

- BDGP (Berkeley Drosophila Genome Project)
last updated 28 July 2014, Human or other, minimum scores for splice sites: 0.1
http://www.fruitfly.org/seq_tools/splice.html [Reese et al., 1997]
- NetGene2
Version 2.4, Human
<http://www.cbs.dtu.dk/services/NetGene2/> [Brunak et al., 1991]

Missense mutations were rated using the following:

- PolyPhen-2 (Polymorphism Phenotyping v2)
Version 2.2.2, NP_065863.2
<http://genetics.bwh.harvard.edu/pph2/> [Adzhubei et al., 2010]
- SIFT Human Protein (Sorting Intolerant From Tolerant)
page last modified: August 2011, Ensembl 63, ENSP00000294618
http://sift.jvvi.org/www/SIFT_enst_submit.html [Ng and Henikoff, 2003]

Sukalo et al. (2015), *Human Mutation*, Supporting Material

- MutPred (Mutation Prediction)
last modified 02 Feb 2014, NP_065863.2
<http://mutpred.mutdb.org> [Li et al., 2009]
- GERP (Genomic Evolutionary Rate Profiling)
hg19,
<http://mendel.stanford.edu/SidowLab/downloads/gerp/index.html> [Cooper et al., 2005]


All variants were checked regarding their appearance/frequency in EVS (Exome Variant Server, Gene ID: 57572, GRCh37, <http://evs.gs.washington.edu/EVS/>, [Exome Variant Server, 2015]) and TGP (1000 Genomes Project, <http://www.1000genomes.org/home>, [1000 Genomes Project Consortium et al., 2010]). Protein conservation across species was checked by Standard Protein BLAST (Basic Local Alignment Search Tool, Database: Reference proteins (refseq_protein), Algorithm: blastp (protein-protein BLAST), <http://blast.ncbi.nlm.nih.gov/Blast.cgi>, [Altschul et al., 1990]). Segregation of the variants across family members was checked if appropriate material was available. Designation of mutations follows the guidelines of the Human Genome Variation Society (last modified March 2014; <http://www.hgvs.org/mutnomen/>) [den Dunnen and Antonarakis, 2000] and was verified by Mutalyzer (Version 2.0 beta-24; <https://mutalyzer.nl/>) [Wildeman et al., 2008].

A focused MLPA assay including probes for *DOCK6* exons 40 through 48 was developed (probe sequences available upon request) for copy number determination of the terminal *DOCK6* exons in a family where PCR amplification of exons 42 to 48 failed in the index patient.


We have established a collection of all mutations and selected unclassified variants in the *DOCK6* gene (<http://databases.lovd.nl/shared/variants/DOCK6>) at the Leiden Open Variation Database (LOVD, version 3.0) [Fokkema et al., 2011], as well as all available phenotype data of patients that were designated as AOS2 both clinically and genotypically (<http://databases.lovd.nl/shared/individuals/DOCK6>). To date, the database contains 17 different *DOCK6* mutations and phenotype data from 18 individuals.

Supp. Figure S1: Pedigrees and electropherograms of *DOCK6*-positive AOS patients and their families.

Family 1




Family 2




c.1362_1365del / p.(Thr455Serfs*24)

Pro Ala Thr **Leu Thr Val Thr**
 CCA GCC ACG **CTA ACT** GTC ACA




c.4491+1G>A


Ile Gly His **+1**
 ATC GGC CAC **g** t g a g a g g g g g



RNA analysis




Family 3




c.1296_1297delinsT / p.(Gln434Argfs*21)

Arg Arg Arg Gly Pro **Gln Asp Arg**
 CGC CGT CGG **GGG** CCC CAG GAC CGG




Family 4




c.4786C>T / p.(Arg1596Trp)

Pro Asp Leu Arg Leu Thr Trp
 CCG GAC CTT CGG CTG ACC TGG




Family 5




c.3154G>A / p.(Glu1052Lys)

Cys Ser His Glu His Tyr Val
 TGC AGC CAC GAG CAC TAC GTG




Family 6




c.788T>A / p.(Val263Asp)

Arg Ile Leu Val Lys Cys Leu
 AGG ATC TTG GTC AAG TGT CTG




c.5939+2T>C

Cys Lys ... +2
 TGC AAG AA g t a g g c g c a a




Family 7




c.1902_1905del / p.(Phe635Profs*32)

His His Leu Leu Phe Thr Phe
 CAT CAC CTG CTG TTC ACC TTC




c.4106+5G>T

Val Asp ... +5
 GTG GAC AA gtaggtgtggg




Family 8




c.484G>T / p.(Glu162*)

Ser Gly Asp Glu Arg Ser Gly
 TCT GGA GAC GAG AGG TCC GGC




Sukalo et al. (2015), *Human Mutation*, Supporting Material


Family 9




MLPA



c.5235+205_6102-15delinsCATGGGGCTG hom



Family 10



c.2520dupT / p.(Arg841Serfs*6)

Tyr Ala Phe **Arg** Leu Pro Gly
TAC GCC TTT CGC CTT CCT GGC



Pedigrees representing *DOCK6* mutation positive families. Affected individuals are indicated by filled symbols. All available genotype data is added and sequence electropherograms are shown. Family 2: RNA analysis is additionally displayed. Family 6: in the index patient the wild type allele is drastically under-represented regarding the heterozygous c.5939+2T>C splice site mutation. We assume this to be caused by a technical artefact due to very poor DNA quality. Family 9: MLPA results (multiplex ligation-dependent probe amplification) are shown.

hom, homozygous; *het*, heterozygous; *WT*, wild type; ?* patient not listed in table 1 due to lack of clinical data and not genetically analyzed due to lack of adequate material.

Sukalo et al. (2015), *Human Mutation*, Supporting Material

Supp. Table S1: Mutations in the *DOCK6* gene causing Adams-Oliver syndrome.

Location	Nucleotide Alteration	Predicted effect ^a	Family ID	EVS	MAF (TGP)	Reference(s)
Exon 05	c.484G>T	p.(Glu162*)	8	-	-	this paper
Exon 07	c.788T>A	p.(Val263Asp)	6	-	-	this paper
Exon 11	c.1245dupT	p.(Asp416*)	12	-	-	Shaheen et al. [2011]
Exon 12	c.1296_1297delinsT	p.(Gln434Argfs*21)	3	-	-	this paper
Exon 12	c.1362_1365del	p.(Thr455Serfs*24)	2, 11	0.26 %	-	this paper + Shaheen et al. [2011]
Exon 17	c.1902_1905del	p.(Phe635Profs*32)	7	-	-	this paper
Exon 21	c.2520dupT	p.(Arg841Serfs*6)	10, 13	-	-	this paper + Shaheen et al. [2013]
Exon 25	c.3047T>C	p.(Leu1016Pro)	1	-	-	this paper
Exon 26	c.3154G>A	p.(Glu1052Lys)	5	-	-	this paper
Exon 26	c.3190_3191del	p.(Leu1064Valfs*60)	15	0.02 %	-	Lehman et al. [2014]
Intron 32	c.4106+5G>T	r.spl.? p.?	7	-	-	this paper
Intron 32	c.4107-1G>C	<i>p.Thr1370Metfs*19</i>	14	-	-	Shaheen et al. [2013]
Exon 35	c.4480G>T	p.(Glu1494*)	15	-	-	Lehman et al. [2014]
Intron 35	c.4491+1G>A	r.spl.? p.?	2	-	-	this paper
Exon 38	c.4786C>T	p.(Arg1596Trp)	4	-	-	this paper
Intron 41 – Intron 47	c.5235+205_6102-15 delinsCATGGGGCTG	p.? ^b	9	-	-	this paper
Intron 46	c.5939+2T>C ^c	r.spl.? p.?	6	0.02 %	-	this paper

^aItalic letters indicate that the effect of splicing mutations was demonstrated on the mRNA level.

^bMLPA analysis revealed deletion of exons 42 to 47.

^cThis alteration is also listed in dbSNP (rs201387914) with unknown pathogenicity and frequency. Online tools predict destruction of the donor splice site.

Mutation nomenclature refers to GenBank reference sequence NM_020812.3. Nucleotide numbering reflects cDNA numbering with +1 corresponding to the A of the ATG translation initiation codon in the reference sequence, according to journal guidelines (www.hgvs.org/mutnomen). The initiation codon is codon 1.

EVS (Exome Variant Server): frequency of alterations was calculated according to the total allele count; MAF (minor allele frequency); TGP (1000 Genomes Project): no entries for these alterations.

Sukalo et al. (2015), *Human Mutation*, Supporting Material

Supp. Table S2: Prediction of pathogenicity and conservation of *DOCK6* missense mutations.

Nucleotide alteration	Predicted effect	PolyPhen-2 HumVar [sensitivity/specificity]	SIFT [score/median information content]	MutPred	GERP	BLAST Alignment
c.788T>A	p.V263D	probably damaging (0.998) ----- 0.18 / 0.98	damaging ----- 0.00 / 2.75	0.735	5.05	Human 263 PPREHFGQRIILVKCLSLKFEIEI Mmulatta 249 ---EHFGQRIILVKCLSLKFEIEI Mmusculus 263 PPREHFGQRIILVKCLSLKFEIEI Trubripes 266 VPKEHCGQRIMVKCLSLKFEIEI Drerio 263 VPKEHSGQRIMVKCLSLKFEIEI Dmelanogaster 271 IPVEHMGHRIQVNCLELRLELEV Celegans 261 LPEQEETPKLIVKVEKAAADPFF Xtropicalis 264 VPKEHFGFRLLVKFLSLKFEIEI
c.3047T>C	p.L1016P	probably damaging (0.977) ----- 0.58 / 0.94	damaging ----- 0.00 / 2.71	0.756	4.84	Human 1016 LSLVDRGFVFSLVRAHYKQVATR Mmulatta 1002 LSLVDRGFVFSLVRAHYKQVATR Mmusculus 1080 -----SLVRAHYKQVATR Trubripes 1088 -SLMDRGFVFNILIRSYKQIANK Drerio 1083 -----LVRSYKQIINN Dmelanogaster 1052 -----GFVFLIKTYTKVLISK Celegans 1027 -----RTFVMKLVHKYLIIFAES Xtropicalis 1044 LSLMDRGFVFNILIRSYKQVMWK
c.3154G>A	p.E1052K	probably damaging (0.999) ----- 0.09 / 0.99	damaging ----- 0.00 / 2.71	0.492	4.81	Human 1052 RMEFTRILCSHEHYVTLNLP-----CP Mmulatta 1038 RMEFTRILCSHEHYVTLNLP-----CP Mmusculus 1116 RMEFTRILCSHEHYVTLNLP-----CP Trubripes 1124 RMEFTRIVGSHHYVTLNLP-----ST Drerio 1119 RMEFTRILCSHEHYVTLNLP-----AT Dmelanogaster 1088 KIDELRIVCSHEHFVALNLPFGTSYTM Celegans 1063 KIDFVRVCSMEHYLIVNI-L-----SD Xtropicalis 1078 ----SHCLGKAGFYFSCSLHC-----GH
c.4786C>T	p.R1596W	probably damaging (0.999) ----- 0.09 / 0.99	damaging ----- 0.00 / 2.71	0.715	4.99	Human 1596 IARGYQGSDDLRLTWLQNMAGKH Mmulatta 1583 IARGYQGSDDLRLTWLQNMAGKH Mmusculus 1660 IARGYQGSDDLRLTWLQNMAGKH Trubripes 1669 IARGYQNSPDLRLTWL----- Drerio 1659 IARGYQNSPDLRLTWLQNMAGKH Dmelanogaster 1618 IARGYQNSPDLRLTWL----- Celegans 1560 -----PDLRLTWLQNMAGKH Xtropicalis 1573 IARGYQNSPDLRLTWLQNMAGKH

Sukalo et al. (2015), *Human Mutation*, Supporting Material

Various online prediction tools were used to evaluate mutation effects. **PolyPhen-2**: score >0.909, probably damaging; score 0.447 – 0.908, possibly damaging; score ≤0.446, benign; sensitivity: True Positive Rate, the chance that the mutation is classified as damaging when it is indeed damaging; specificity: 1 – False Positive Rate, the chance that the benign mutation is correctly classified as benign [Adzhubei et al., 2013]. **SIFT**: score = normalized probability that the amino acid change is tolerated; ≤0.05, damaging; >0.05, tolerated; median information content: maximum 4.32, indicates complete conservation at this position; minimum 0.00, indicates a position where all 20 amino acids are tolerated; ideally between 2.75 and 3.5 [Ng and Henikoff, 2003]. **MutPred**: general score; ranges between 1.000 (deleterious mutation) and 0.000 (benign). **GERP**: ranges from 6.17 (highly conserved amino acid residue) to -12.3 (not conserved). **BLAST Alignment**: multiple protein alignment of human DOCK6 and its orthologues; numbers indicate position of affected amino acid residue. Black shading indicates identical amino acid residues; grey shading indicates similar residues (according to BLOSUM62 matrix). Human: *Homo sapiens*, ENSP00000294618.6. Mmulatta: *Macaca mulatta*, ENSMMUP00000012229.2. Mmusculus: *Mus musculus*, ENSMUSP00000034728.7. Trubripes: *Takifugu rubripes*, ENSTRUP00000027209.1. Drerio: *Danio rerio*, ENSDARP00000077379.4. Dmelanogaster: *Drosophila melanogaster*, FBpp0077762.3. Celegans: *Caenorhabditis elegans*, F46H5.4.1. Xtropicalis: *Xenopus tropicalis*, ENSXETP00000036553.2.

Supp. Table S3: Splice site prediction c.4106+5G>T.

		Donor splice site predictions	
		BDGP	NetGene2
	+5		
Wild type	CGCGTGGACAAgtag g tgtgggcaggagggt	0.77	0.881
Mutant	CGCGTGGACAAgtag t tgtgggcaggagggt	0.15	0.341

This donor splice site alteration at position +5 in intron 32 was detected in compound heterozygosity with a frameshift mutation in family 7. The online tools BDGP (Berkeley Drosophila Genome Project) and NetGene2 predict this nucleotide exchange to impair the regular splice donor site at the border between exon 32 and intron 32. The indicated probability scores refer to the authentic splice site.

Sukalo et al. (2015), *Human Mutation*, Supporting Material

Supp. Table S4: Unclassified variants in the *DOCK6* gene.

Location	Nucleotide Alteration ^a	Predicted effect	dbSNP (139)	EVS	MAF (TGP)
Exon 02	c.100C>G	p.(His34Asp)	rs201065561	-	0.0004/2
Exon 09	c.885C>T	p.(=)	rs146599144	0.46 %	0.0048/24
Exon 12	c.1289G>A	p.(Arg430His)	rs143655255	0.30 %	0.0028/14
Exon 12	c.1358C>T	p.(Thr453Met)	-	-	-
Exon 13	c.1445C>T	p.(Pro482Leu)	-	-	-
Intron 16	c.1833-19C>G	r.spl.? p.?	rs188183013	0.22 %	0.0028/14
Exon 19	c.2104G>A	p.(Gly702Ser)	rs199838752	0.08 %	0.0020/10
Exon 23	c.2767G>A	p.(Val923Ile)	rs143194982	0.02 %	0.0002/1
Exon 30	c.3873C>T	p.(=)	rs200843111	0.03 %	-
Exon 31	c.3913C>T	p.(Arg1305Cys)	rs112911897	0.70 %	0.0050/25
Exon 37	c.4732C>T	p.(Leu1578Phe)	-	-	-
Exon 38	c.4899G>A	p.(=)	rs72985308	0.20 %	0.0010/5
Exon 41	c.5229C>A	p.(=)	rs56243833	0.21 %	0.0026/13
Exon 44	c.5640C>T	p.(=)	rs200959822	0.11 %	0.0004/2
Intron 44	c.5688+9G>C	r.spl.? p.?	-	0.09 %	0.0002/1
Intron 45	c.5833-16C>G	r.spl.? p.?	rs199764395	0.05 %	0.0002/1

Only variants within 20 bp of the exons and $MAF \leq 0.01$ were included. All variants show no alteration of splice site prediction in BDGP and NetGene2. None of these variants appeared as homozygous in TGP (1000 Genomes Project). The numbering for the nucleotide changes are based on cDNA sequence in accordance with the GenBank entry NM_020812.3 (GRCh37). ^aFor cDNA numbering, +1 corresponds to the A of the ATG translation initiation codon in the reference sequence.

EVS: Exome Variant Server, frequency of alterations was calculated according to the total allele count; MAF: minor allele frequency [frequency of alternative nucleotide / total allele count of alternative nucleotide].

Sukalo et al. (2015), Human Mutation, Supporting Material

Supp. Table S5: Prediction of pathogenicity and conservation of DOCK6 unclassified missense variants.

Nucleotide alteration	Predicted effect	PolyPhen-2 HumVar [sensitivity/specificity]	SIFT [Score/Median information content]	MutPred	GERP	BLAST Alignment
c.100C>G	p.H34D	benign (0.017) ----- 0.95 / 0.54	tolerated ----- 0.34 / 2.77	0.155	1.86	Human 34 KQV---SRERSGSPHSSRR---CSSSLGV
						Mmulatta 20 KQV---SRERSGSPSSRR---CSSSLGV
						Mmusculus 34 KQV---SRERSGSPHSSRR---SSSSLGV
						Trubripes 32 KQV---SREY-GSPQLSKK[7]VSHVTQL
						Drerio 36 KQV---SREYGSPQMSKKR---AGAHQPV
						Dmelanogaster 33 KNV---SGCHLSKAMDPSL---CGSSISP
						Celegans 35 KHV[7]HRLSEGDNGLDLA---VSMMEKI
Xtropicalis 34 KQV---AREYGGSPQLSKK---RGGQASV						
c.1289G>A	p.R430H	possibly damaging (0.512) ----- 0.82 / 0.81	tolerated ----- 0.12 / 2.78	0.334	3.11	Human 430 GERPAWT---DRRRRGPOD---RASSGD
						Mmulatta 416 GERRSAWT---DRRRRGPOD-----
						Mmusculus 428 --RRPTWA---ERRRRGPOD---RGYSGD
						Trubripes 435 KKGHGTWN---ERKKG-GE---RMSIAD
						Drerio 431 ----GTWN---ERKKG-GE---RMSVGE
						Dmelanogaster 431 SLDRKSSTSEFDQLRRKAND[5]-----
						Celegans 423 PMMMSQCT---TASGAVLTT---AGQSQD
Xtropicalis 431 -ERKGTWN---ERKKA-GE---RLSVGD						
c.1358C>T	p.T453M	probably damaging (0.974) ----- 0.59 / 0.93	damaging ----- 0.00 / 2.78	0.303	4.15	Human 453 DACSF-SGFRPAT---LTVTNFFKQEA
						Mmulatta 439 DACSF-SGFRPAT---LTVTNFFKQEA
						Mmusculus 451 DACSF-SSFRPAT---LTVTNFFKQEA
						Trubripes 458 DTCNF-ATFRPAT---LTVTNFFKQEG
						Drerio 454 DMCNF-TNFRPAT---LTVTNFFKQEG
						Dmelanogaster 482 DFANVVENFREIT---ITVPSFFKQEA
						Celegans 465 -----GTSAA[37]LKFSSFRQEG
Xtropicalis 454 ETCGL-HTFRPAT---LTVTNFFKQEG						
c.1445C>T	p.P482L	probably damaging (0.991) ----- 0.50 / 0.95	damaging ----- 0.00 / 2.76	0.416	4.42	Human 482 DLFKFLADMRRPSSLLRRLRPVT
						Mmulatta 468 DLFKFLADMRRPSSLLRRLRPVT
						Mmusculus 480 -----RPSLLRRLRPVT
						Trubripes 487 --YKFLADMRRPSSVLRRLRPVT
						Drerio 483 DLYKFLADMRRPSSVLRRLRPVT
						Dmelanogaster 511 DLYKTLPELKR-----
						Celegans 531 DIYRICSEMRRRTNGKVHK-KMFN
Xtropicalis 482 -----RRPSTALRRLRPVT						

Sukalo et al. (2015), *Human Mutation*, Supporting Material

Supp. Table S5: Prediction of pathogenicity and conservation of *DOCK6* unclassified missense variants. (*continued*)

Nucleotide alteration	Predicted effect	PolyPhen-2 HumVar [sensitivity/specificity]	SIFT [Score/Median information content]	MutPred	GERP	Alignment (numbers indicate position of mutated protein)		
						Species	Position	
c.2104G>A	p.G702S	benign (0.211)	damaging	0.389	3.62	Human	702	DVALPGMRWVDG H KGVFSVELTA
		0.88 / 0.74	0.02 / 2.74			Mmulatta	688	DVALPGMRWVDG H KGVFSVELTA
c.2767G>A	p.V923I	benign (0.029)	tolerated	0.327	3.53	Mmusculus	700	DVALPGMRWVDG C HKGVSVELTA
		0.94 / 0.59	0.12 / 2.71			Trubripes	707	DVQLPGMKWVDN H KGVFSVEVT-
c.3913C>T	p.R1305C	benign (0.259)	damaging	0.559	2.50	Drerio	702	DVQLPGMKWVDN H KGVFNVEVKA
		0.87 / 0.75	0.00 / 2.71			Dmelanogaster	747	NVHLPGIKWLDN H RAVFSINVEA
c.4732C>T	p.L1578F	possibly damaging (0.453)	damaging	0.594	4.21	Celegans	743	NNALPNLKWVDN H KPTFSCS---
		0.83 / 0.80	0.02 / 2.71			Xtropicalis	702	DVQLPGMKWVDN H KPVFSVDLVA
						Human	923	LALQWVSSSA V REAILQHAWFF
						Mmulatta	909	--LQWVSSSA V REAILQHAWFF
						Mmusculus	987	LALQWVSSGSA V RELVLQHAWFF
						Trubripes	981	LALQWVSSNS T VREAILQHAWFF
						Drerio	986	LALQWVSS T STVREASLQCAWFF
						Dmelanogaster	963	LAL H WVVASG K AADLAMSNSWFL
						Celegans	943	LLEVWLRARG S LRDVSLVHSWFL
						Xtropicalis	916	LVLQWVSSA A V-----
						Human	1305	AFEYKGGKAF E RINSLTFKK--SLD
						Mmulatta	1291	AFEYKGGKAF E RINSLTFKK--SLD
						Mmusculus	1369	AFEYKGGKAF E RINSLTFKK--S--
						Trubripes	1373	CFEYKGRKA L E R INSLTFKK--S D
						Drerio	1368	CFEYKGGKAF E RINSLTFKK--S D
						Dmelanogaster	1329	-----L K R T NT Q S F R K T G S T D
						Celegans	1290	----- A S A R S P--D K T--S L -
						Xtropicalis	1282	CF Q YKGGKAF E RINSLTFKK--SLD
						Human	1578	KMKEHQEDPE M LIDLMYRIARG Y
						Mmulatta	1565	-----E D PE M LIDLMYRIARG Y
						Mmusculus	1642	KMKEHQEDPE M LIDLMYRIARG Y
						Trubripes	1651	KMKEHQ Q DPE M LIDLMYRIAK G Y
						Drerio	1641	KMKEHQ Q DPE M LIDLMYRIAK G Y
						Dmelanogaster	1600	KMKE Y Q E DPE M LIDLM N RIAK G Y
						Celegans	1539	R M RE H V N D Y E M TIDLM Y Q L VE G Y
						Xtropicalis	1555	KMKEHQ Q DPE M LIDLM Y -----

Legend: see above (Supp. Table S2).

Sukalo et al. (2015), *Human Mutation*, Supporting Material

Supp. References

- 1000 Genomes Project Consortium, Abecasis GR, Altshuler D, Auton A, Brooks LD, Durbin RM, Gibbs RA, Hurles ME, McVean GA. 2010. A map of human genome variation from population-scale sequencing. *Nature* 467:1061-73. Erratum in: *Nature* 473:544.
- Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. 2010. A method and server for predicting damaging missense mutations. *Nat Methods* 7:248-9.
- Adzhubei I, Jordan DM, Sunyaev SR. 2013. Predicting Functional Effect of Human Missense Mutations Using PolyPhen-2. *Curr Protoc in Hum Genet* 76:7.20:7.20.1–7.20.41.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403-10.
- Brunak S, Engelbrecht J, Knudsen S. 1991. Prediction of Human mRNA Donor and Acceptor Sites from the DNA Sequence. *J Mol Biol* 220:49-65.
- Cooper GM, Stone EA, Asimenos G; NISC Comparative Sequencing Program, Green ED, Batzoglou S, Sidow A. 2005. Distribution and intensity of constraint in mammalian genomic sequence. *Genome Res* 15:901-13.
- den Dunnen JT, Antonarakis SE. 2000. Mutation nomenclature extensions and suggestions to describe complex mutations: a discussion. *Hum Mutat* 15:7-12.
- Exome Variant Server, NHLBI GO Exome Sequencing Project (ESP), Seattle, WA (URL: <http://evs.gs.washington.edu/EVS/>) [Jan, 2015].
- Fokkema IF, Taschner PE, Schaafsma GC, Celli J, Laros JF, den Dunnen JT. 2011. LOVD v.2.0: the next generation in gene variant databases. *Hum Mutat* 32:557-63.
- Lehman A, Stittrich AB, Glusman G, Zong Z, Li H, Eydoux P, Senger C, Lyons C, Roach JC, Patel M. 2014. Diffuse angiopathy in Adams-Oliver syndrome associated with truncating DOCK6 mutations. *Am J Med Genet A* 164A:2656-62.
- Li B, Krishnan VG, Mort ME, Xin F, Kamati KK, Cooper DN, Mooney SD, Radivojac P. 2009. Automated inference of molecular mechanisms of disease from amino acid substitutions. *Bioinformatics* 25: 2744-50.
- Ng PC, Henikoff S. 2003. SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res* 31:3812-4.
- Reese MG, Eeckman FH, Kulp D, Haussler D. 1997. Improved Splice Site Detection in Genie. *J Comp Biol* 4:311-23.
- Shaheen R, Faqeih E, Sunker A, Morsy H, Al-Sheddi T, Shamseldin HE, Adly N, Hashem M, Alkuraya FS. 2011. Recessive mutations in DOCK6, encoding the guanidine nucleotide exchange factor DOCK6, lead to abnormal actin cytoskeleton organization and Adams-Oliver syndrome. *Am J Hum Genet* 89:328-33.
- Shaheen R, Aglan M, Keppler-Noreuil K, Faqeih E, Ansari S, Horton K, Ashour A, Zaki MS, Al-Zahrani F, Cueto-Gonzalez AM, Abdel-Salam G, Temtamy S, et al. 2013. Mutations in EOGT confirm the genetic heterogeneity of autosomal-recessive Adams-Oliver syndrome. *Am J Hum Genet* 92:598-604.
- Untergrasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012. Primer3 - new capabilities and interfaces. *Nucleic Acids Res* 40:e115.
- Wildeman M, van Ophuizen E, den Dunnen JT, Taschner PE. 2008. Improving sequence variant descriptions in mutation databases and literature using the Mutalyzer sequence variation nomenclature checker. *Hum Mutat* 29:6-13.