

## **Marked overlap of four genetic syndromes with dyskeratosis congenita confounds clinical diagnosis**

Amanda J. Walne<sup>1\*</sup>, Laura Collopy<sup>1</sup>, Shirleny Cardoso<sup>1</sup>, Alicia Ellison<sup>1</sup>, Vincent Plagnol<sup>2</sup>, Canan Albayrak<sup>3</sup>, Davut Albayrak<sup>3</sup>, Sara Sebnem Kilic<sup>4</sup>, Turkan Patiroglu<sup>5</sup>, Haluk Akar<sup>5</sup>, Keith Godfrey<sup>6</sup>, Tina Carter<sup>7</sup>, Makia Marafie<sup>8</sup>, Ajay Vora<sup>9</sup>, Mikael Sundin<sup>10,11</sup>, Thomas Vulliamy<sup>1</sup>, Hemanth Tummala<sup>1</sup>, Inderjeet Dokal<sup>1</sup>

<sup>1</sup>Centre for Genomics and Child Health, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, Barts NHS Trust, London, UK

<sup>2</sup>University College London Genetics Institute, London, UK

<sup>3</sup>Department of Paediatric Haematology, Ondokuz Mayıs University, Samsun, Turkey

<sup>4</sup>Department of Paediatric Immunology Uludag University, Bursa, Turkey

<sup>5</sup>Department of Paediatric Immunology Erciyes University Medical Facility, Kayseri, Turkey

<sup>6</sup>Department of Paediatric Dermatology and NIHR Southampton Biomedical Research Centre, University Hospital, Southampton and University of Southampton, Southampton, UK

<sup>7</sup>Department of Oncology and Haematology, Princess Margret Hospital, WA, Australia

<sup>8</sup>Clinical Cancer and Community Genetics, Kuwait Medical Genetics centre, Kuwait

<sup>9</sup>Department of Haematology, Sheffield Children's NHS foundation Trust, Sheffield, UK;

<sup>10</sup>Section of Paediatric Haematology/Immunology/SCT, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm Sweden

<sup>11</sup>Division of Paediatrics, CLINTEC, Karolinska Institutet, Stockholm, Sweden.

Running head: DC or not DC? A Clinical Dilemma

\*Corresponding author: Amanda Walne, Centre for Genomics and Child Health, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, 4 Newark Street, London E1 2AT, UK

Telephone +44 207 882 2458, Fax +44 207 882 2195

Email a.walne@qmul.ac.uk

Word count abstract: 250

Word count text: 3953

Number of figures and tables: 5 figures and 2 tables

Number of references: 38

Acknowledgments: The authors would like to thank all the clinicians and patients who have helped us over the years, particularly Drs Babik, Balci, Ghoulam and Leblanc. We would also like to thank the staff at Barts and The London Genome Centre for Sanger sequencing analysis. Financial support is provided by The Medical Research Council-MR/K000292/1, Children with Cancer- 2013/144 and Blood Wise-14032 (AJW, LC, SC, AE, TV, HT and ID). KMG is supported by the National Institute for Health Research through the NIHR Southampton Biomedical Research Centre.

## ABSTRACT

Dyskeratosis congenita is a highly pleotropic genetic disorder. This heterogeneity can lead to difficulties in making an accurate diagnosis and delays in appropriate management. The aim of this study was to determine the underlying genetic basis in patients presenting with features of dyskeratosis congenita and who were negative for mutations in the classical dyskeratosis congenita genes. By whole exome and targeted sequencing, we identified biallelic variants in genes that are not associated with dyskeratosis congenita in 17 individuals from 12 families. Specifically these were homozygous variants in *USBI* (eight families), homozygous missense variants in *GRHL2* (two families) and identical compound heterozygous variants in *LIG4* (two families). All patients had multiple somatic features of dyskeratosis congenita but not the characteristic short telomeres. Our case series shows that biallelic variants in *USBI*, *LIG4* and *GRHL2*, the genes mutated in poikiloderma with neutropenia, *LIG4*/Dubowitz syndrome and the recently recognised ectodermal dysplasia/short stature syndrome, respectively, cause features that overlap with dyskeratosis congenita. Strikingly these genes also overlap in their biological function with the known dyskeratosis congenita genes that are implicated in telomere maintenance and DNA repair pathways. Collectively these observations demonstrate the marked overlap of dyskeratosis congenita with four other genetic syndromes confounding accurate diagnosis and subsequent management. This has important implications for establishing a genetic diagnosis when a new patient presents in the clinic. Patients with clinical features of dyskeratosis congenita need to have genetic analysis of *USBI*, *LIG4* and *GRHL2* in addition to the classical dyskeratosis congenita genes and telomere length measurements.

## INTRODUCTION

Dyskeratosis congenita (DC) is a highly heterogeneous genetic and clinical syndrome. The classical presentation of DC is characterized by the mucocutaneous triad of abnormal skin pigmentation (hyper/hypopigmentation with atrophy and telangiectasia, termed poikiloderma), oral leukoplakia and nail dystrophy. Individuals with DC frequently develop bone marrow (BM) failure, are at a high risk of developing cancer and can develop disease features in virtually every system in the body. Pathologically DC is characterized by selective exhaustion of highly proliferative cells<sup>1</sup> that have critically short telomeres and exhibit an abnormal DNA damage response.<sup>2</sup>

The Dyskeratosis Congenita Registry (DCR, London) is a collection of patients that have a clinical diagnosis of DC or an overlapping phenotype as defined by Dokal et al.<sup>3</sup> The diagnostic inclusion criteria are:

- (1) All three mucocutaneous features (abnormal skin pigmentation, nail dystrophy and oral leukoplakia).
- (2) One mucocutaneous feature plus BM failure and two other somatic features of DC.
- (3) Aplastic anaemia (AA), myelodysplastic syndrome (MDS) or idiopathic pulmonary fibrosis (IPF) associated with a pathogenic telomerase variant.
- (4) Hoyeraal Hreidarsson syndrome (HHS, growth retardation, developmental delay, microcephaly, cerebellar hypoplasia, BM failure and immunodeficiency).
- (5) Two or more features seen in DC plus very short telomeres (<1<sup>st</sup> centile).

Based on these criteria the phenotype of DC is highly variable. Furthermore, not all physical malformations are present at the point of diagnosis, with many additional features developing with age or, in the case of severe disease, death may occur before more typical clinical presentations become apparent.

To date pathogenic mutations have been identified in 11 genes that cause DC according to the criteria above: dyskeratosis congenita 1, dyskerin (*DKC1*), telomerase RNA component (*TERC*), telomerase reverse transcriptase (*TERT*), NOP10 ribonucleoprotein (*NOP10*), NHP2 ribonucleoprotein (*NHP2*), TERF1 (TRF1)-interacting nuclear factor 2 (*TINF2*), WD repeat containing, antisense to TP53 (*WRAP53*, also known as *TCAB1*), CST telomere maintenance complex component 1 (*CTC1*), regulator of telomere elongation helicase 1 (*RTEL1*), adrenocortical dysplasia homolog (mouse) (*ACD*) and poly(A)-specific ribonuclease (*PARN*). *DKC1*, *TERC*, *TERT*, *NOP10* and *NHP2* all encode components of the telomerase complex which is involved in telomere elongation. *TINF2* and *ACD* encode components of the shelterin complex which is involved in telomere protection. *CTC1*

encodes a member of the CST complex which facilitates recruitment and docking of telomerase on to the telomere. *WRAP53* is involved in telomerase trafficking and *RTEL1* is involved in telomere replication, and DNA replication and repair.<sup>3-7</sup> The final gene, *PARN*, is not exclusively involved in telomere biology but is mutated in a small number of cases of DC, and is an exoribonuclease involved in the control of mRNA stability and the maturation of TERC snoRNA.<sup>8-10</sup> As the majority of DC genes are involved in telomere biology and DC patients usually have short telomeres, this has led to DC being classed as a telomeropathy.<sup>11</sup> However these mutations do not explain all the cases that are present in the DCR (Data from the DCR, London, April 2016).

The advent of next generation sequencing, particularly whole exome sequencing (WES), has become an invaluable tool for determining the causal variant(s) underlying a specific disease. By sequencing all the coding regions and comparing variants in many individuals with an overlapping phenotype it is possible to elucidate potential disease genes in a way that would not have been possible five years ago.<sup>12</sup> This can give the clinical diagnosis a genetic basis. The main problem in obtaining a definitive clinical diagnosis is the degree of phenotypic overlap that exists between many different diseases. The diagnosis, in many cases, depends on the interpretation of the initial clinician and this is often dependent on their speciality. Although subsequent opinions may be sought, the preliminary diagnosis tends to remain with the patient until proven otherwise. An accurate genetic diagnosis can aid clinical diagnosis and subsequent patient management. This has the potential to suggest alternative treatment avenues that may not have been considered and also aid the selection of healthy sibling donors for hematopoietic stem cell transplantation, should this become necessary.

Here we report 12 families with an initial diagnosis of DC who were found to have biallelic variants in one of three genes linked to a disease distinct from DC. The genes identified were *USB1* (U6 snRNA biogenesis 1) associated with poikiloderma with neutropenia (PN), *GRHL2* (grainyhead like transcription factor 2) which is associated with ectodermal dysplasia/short stature syndrome (ECTDS) and *LIG4* (Ligase IV, DNA, ATP-dependent) which is predominantly associated with LIG4 syndrome and occasionally Dubowitz syndrome. The key presenting features of PN are poikiloderma with non-cycling neutropenia, recurrent infections, short stature and nail abnormalities<sup>13</sup>. The typical clinical features associated with ECTDS are short stature, nail dystrophy, abnormal oral pigmentation, and keratoderma and hyperkeratosis of the hands and feet<sup>14</sup>. LIG4 syndrome is characterized by immune deficiency and developmental and growth delay<sup>15</sup>. Patients can also display unusual facial features, microcephaly, pancytopenia and various skin abnormalities. Dubowitz syndrome is characterised by growth failure/short stature, characteristic facial features, microcephaly, mild mental retardation and eczema<sup>16</sup>. All of these disorders have a high degree of phenotypic overlap with DC, thus leading to difficulties in making an accurate clinical diagnosis. The identification of biallelic *USB1*, *GRHL2* and

*LIG4* mutations in 12 different families, initially diagnosed to have DC, therefore demonstrate the marked phenotypic overlap of ‘classical DC’ with four other genetic syndromes.

## **METHODS**

### **Patient samples**

Exome capture or targeted gene screening was performed on a series of genetically uncharacterized index cases in the Dyskeratosis Congenita Registry (held at Barts and The London). Presenting features included some/all of the classic mucocutaneous abnormalities (abnormal skin pigmentation, nail dystrophy, leukoplakia), with or without bone marrow failure. Peripheral-blood samples were obtained with written consent under the approval of our local research ethics committee (London – City and East). Genomic DNA was extracted from these peripheral blood samples for use in all the subsequent analyses (Puregene, Qiagen).

### **Telomere length measurement**

Telomere lengths were measured using the MMqPCR method (modified from Cawthon<sup>17</sup>), on a LightCycler 480 real time thermocycler (Roche). Briefly, in each well, amplification of telomeric DNA (T) and a single copy gene (S) were quantified against standard curves obtained from dilution of a reference DNA sample. The T/S ratio, obtained in triplicate for each sample, is proportional to the telomere length. This ratio was normalized to the T/S ratio of a second reference sample that was run on every plate to give a relative T/S ratio.

### **Whole exome sequencing (WES)**

50ng genomic DNA was subjected to library preparation and exome capture using the Nextera Rapid Capture Exome kit (Illumina). Sequencing was performed using the Illumina HiSeq 2000 system and 100bp paired end reads were generated and the data processed through the Illumina pipeline. Variants were called as described previously.<sup>8</sup> All relevant variants were confirmed by Sanger sequencing.

### **Targeted gene screening**

We designed an NGS assay covering the coding regions and some 5’UTR’s from 31 genes associated with genetic bone marrow failure syndromes (Supplementary Table 1). We used the Illumina TruSeq custom amplicon kit for library preparation and capture according to manufacturer’s instructions. The resultant targeted fragments were indexed by dual barcodes and then sequenced on the Illumina MiSeq platform. Read alignment was performed automatically using the online MSR: TruSeq amplicon software. Annotation of the variants was performed using ANNOVAR.

## **Variant filtering**

Exome data from our patients were jointly called with 2500 WES internal control samples (UCL-ex consortia) with unrelated conditions to minimise artificial batch effects. Where there was a family history we assumed an autosomal recessive mode of inheritance as all parents were reportedly asymptomatic. Filtering was performed using the following criteria: All variants had to pass the Illumina filter, have a read depth of >10X, be present in the general population at a frequency of <0.0002 as reported on Exome Aggregation Consortium database, ExAC ([www.exac.broadinstitute.org](http://www.exac.broadinstitute.org)), 1000 genome project ([www.1000genomes.org](http://www.1000genomes.org)) and from the UCL-ex consortia. Variants that were predicted to be tolerated and benign by SIFT (<http://sift.jcvi.org/>) and Polyphen (<http://genetics.bwh.harvard.edu/pph2/>) were then removed from further analysis. All relevant variants were confirmed by Sanger sequencing.

## **RESULTS**

### **Telomere length analysis**

It has been well established that in the majority of patients with “classical” DC who have mutations in components of the telomere maintenance pathway have short telomeres, usually below the 1<sup>st</sup> centile.<sup>18, 19</sup> In fact this measure is often used as a diagnostic screening tool to perform a differential diagnosis of DC from other bone marrow failure syndromes such as Fanconi anemia.<sup>20</sup> Telomere lengths were therefore measured in our patients by MMqPCR and this was used as an indicator of whether we could be looking for a mutation in a known telomere biology related gene or not. As none of the patients reported in this study had short telomeres when compared with controls (Figure 1), this suggested that we were looking for mutations outside the spectrum of those usually associated with DC but they could still be associated with other bone marrow failure diseases or else have a previously unidentified association to another pathology.

### **Identification of causal genetic variants**

Analysis of our data has identified significant biallelic variants in three genes as detailed below. Five families underwent WES (Families 4, 9-12) and the remaining underwent targeted sequencing either using a 31 bone marrow failure disease gene panel (Supplementary Table 1) or direct sequencing.

#### ***USB1* (U6 snRNA biogenesis 1)**

Biallelic *USB1* mutations have been previously described in patients with DC and the overlapping disease poikiloderma with neutropenia (PN).<sup>21, 22</sup> Since our previous publication,<sup>21</sup> we have identified homozygous *USB1* variants in an additional eight families with features of DC. In seven of the eight

families, the variant was identified by targeted sequencing and the remaining family (two cases) underwent WES. The only variant that was homozygous and shared by both affected siblings was in *USB1* so this was deemed to be causal. Five out of the eight index cases had the previously documented recurring homozygous variant c.531delA, p.His179MetfsTer86<sup>13</sup> (Figure 2A) which is not described on ExAc, (accessed April 2016). The sixth index case was found to have the homozygous variant c.673C>T, p.Gln225Ter (Figure 2B), which has been reported previously as part of a compound heterozygote.<sup>23</sup> The remaining two families (Family 7 and Family 8) have the same homozygous missense mutation c.623A>G, p.His208Arg (Figure 2C). This is the first report of a homozygous missense being associated with disease in *USB1* as to date all 19 disease-causing variants reported in *USB1* are predicted to be loss of function (splicing, nonsense or frame shift).<sup>13</sup> Segregation analysis confirmed an autosomal recessive mode of inheritance and homozygous variants were confirmed in an additional four affected siblings within these families (Figure 2A). In silico analysis of mutations on USB1 crystal structure (PDB id: 4W7H, Figure 2D) revealed that both p.His179MetfsTer86 and p.Gln225Ter truncates several  $\alpha$ -helices and  $\beta$ -sheets in USB1 (Figure 2E and 2F). Furthermore the homozygous missense change p.His208Arg identified in families 7 and 8 affects the highly conserved histidine residue H208, (Supplementary Figure 1 and Figure 2G and 2H). This change is predicted to be probably damaging by polyphen2 (<http://genetics.bwh.harvard.edu/pph2/>; HumVar score 1.0) and disease causing by Mutation taster (<http://www.mutationtaster.org>). This H208 residue is implicated in the catalytic mechanism of USB1, where it facilitates displacement of uridine nucleoside on RNA substrate to stabilise U6 small nuclear RNA, which in turn plays a critical role in RNA splicing.<sup>24</sup>

Table 1 summarises the clinical presentation of these families and the similarity between the clinical presentation of DC and PN is further highlighted in Figure 3. The degree of overlap between the reticular pigmentation observed on the legs (Figure 3A-C), nail dystrophy of the toes (Figure 3D-F) is marked. Similar pigmentation is seen on the trunk (Figure 3G and 3H) and nail dystrophy and blistering is shown on the hands (Figure 3I). In addition to highlighting the similarity between the cutaneous presentation of DC and PN, panels B, C, E, F and H, I also emphasise the difference in presentation that can be caused by mutations within the same gene. The blood results detailed in Table 1 show that although there is neutropenia in all the patients, the overall blood pathology is more global rather than being restricted to just one lineage. This investigation has expanded the repertoire of *USB1* variants identified to date, by adding a recurrent novel homozygous missense variant.

### ***GRHL2* (grainyhead like transcription factor 2)**

The index case from both families 9 and 10<sup>25</sup> underwent WES, and as both families were consanguineous, we assumed an autosomal recessive mode of inheritance. Using the filtering



approach described, it was noted that both families had homozygous non synonymous variants in the gene *GRHL2* c.1445T>A p.Ile482Lys in Family 9 and c.1213C>A p.Pro405Thr in Family 10 (other homozygous calls that fulfilled the filtering criteria are detailed in Supplementary Table 2). Segregation analysis confirmed that both variants were inherited in an autosomal recessive manner (Figure 4A). Both of these single base substitutions affect highly conserved nucleotides (Supplementary Figure 2). As neither variant is reported on ExAc (accessed April 2016) and both variants are predicted by polyphen2 to be probably damaging (HumVar scores p.P405T - 0.995 and p.I482K - 0.917) and disease causing by Mutation Taster, *GRHL2* is the disease causing gene in these two families. *GRHL2* is a member of a highly conserved family of transcription factors that are essential for epithelial development. At the molecular level, the GRHL transcription factors regulate the expression of proteins involved in cell proliferation, differentiation, adhesion and polarity. These factors adopt a DNA-binding immunoglobulin fold homologous to the DNA-binding domain of key tumour suppressor p53.<sup>26</sup> Recently Petrof and colleagues described homozygous mutations in *GRHL2* in two families with ectodermal dysplasia/short stature syndrome (ECTDS)<sup>14</sup>. One of these was p.Ile482Lys, which is the same variant as identified in Family 9. All the variants described by us and Petrof et al., affect the DNA binding domain of the protein (Figure 4A). Phenotypically there are many shared features between our two families (Families 9 and 10) and the ECTDS families reported in the literature (Table 2). Another recent study<sup>27</sup> examined the role of *GRHL2* in the developing kidney and clarified the involvement of *GRHL2* in a network that controls the development of lumen expansion and barrier formation in renal epithelia, which may explain the previously reported renal agenesis seen in Family 10.<sup>25</sup> This suggests that homozygous variants in the gene *GRHL2*, previously linked to ECTDS, are disease-causing in these two families.

#### ***LIG4* (Ligase IV, DNA, ATP-dependent)**

Analysis of exome data from the index cases of Families 11 and 12 failed to reveal any homozygous variants that passed the filtering strategy. However, a biallelic analysis showed that they both shared the same compound heterozygous variants in *LIG4* namely c.2440C>T, p.Arg814Ter and c.1270\_1274 delAAAAG, p.Lys424ArgfsTer20 (Figure 4B, other biallelic calls are listed in Supplementary Table 2). Although both variants observed in Families 11 and 12 are reported on ExAc at very low frequency (0.000082 and 0.00014, respectively) in the heterozygous state, the likelihood of them occurring together (as is the case here) by chance is  $2.4 \times 10^{-8}$ . *LIG4* is involved in DNA non-homologous end joining and V(D)J recombination.<sup>28</sup> Biallelic mutations in *LIG4* are rare with only 28 cases being reported to date in the literature,<sup>29</sup> and are usually associated with *LIG4* syndrome and more recently in Dubowitz syndrome (DS).<sup>16</sup> Both *LIG4* syndrome and DS belong to a group of disorders that are associated with impaired DNA damage response mechanisms.<sup>28</sup> *LIG4* syndrome is a hereditary disorder associated with impaired DNA double-strand break repair mechanisms. It is

characterized by growth restriction, developmental delay, microcephaly, facial dysmorphism, pancytopenia, variable immune deficiency and an increased predisposition to leukemia. DS is a rare multiple congenital syndrome characterized by cognitive delay, growth failure, microcephaly, distinctive facial dysmorphism, immune defects, pancytopenia, hematological malignancy and neuroblastoma. The degree of clinical overlap between *LIG4* syndrome, DS and our two patients is marked (Table 2). Furthermore the fact that both of the variants in *LIG4* described here have been previously documented to be pathogenic and have been shown to occur in the same combination<sup>30</sup> suggests that these are the causal genetic defects in these two unrelated individuals.

## DISCUSSION

Historically the diagnosis of disease is defined by the presenting clinical features but it is becoming more apparent that this can give conflicting diagnoses as shown in Tables 1 and 2. Recent advances in whole exome and targeted sequencing have enabled a genetic diagnosis to be obtained more readily than ever before. This now enables the clinician to consider the possibility of diseases with overlapping phenotype based on a genetic result rather than solely relying on the clinical presentation. In this report we have elucidated the underlying genetic causal variant(s) in 12 families (comprising 17 affected individuals) who were given a clinical diagnosis of dyskeratosis congenita. Initially we undertook whole exome sequencing on uncharacterised patients who presented with sufficient clinical features to be diagnosed as DC. As targeted sequencing methodologies improved, we then chose to use this approach as a screening tool to identify patients who had mutations in the known DC and bone marrow failure genes (supplementary Table 1). The data reported here represent a sub-set of all the cases analysed, where we believe the disease-causing gene has been identified and where the identified gene has been previously associated with another genetic syndrome. By using a combination of these two strategies we identified disease-causing variants in the genes *USBI* (eight families, 12 cases), *GRHL2* (two families, three cases) and *LIG4* (two families, two cases). Clinically all patients show a high degree of overlap with the phenotypic profile for DC as described by Dokal et al<sup>3</sup> and the other diseases with which the disease-causing variants have been associated. This presents a problem in defining the disease: is it best to use phenotype or genotype?

Significantly short telomeres are often used as a biomarker for DC. A telomere length below the 1<sup>st</sup> centile is considered diagnostic for DC but this measurement is not routinely reported in patients with other bone marrow failure diseases. Figure 1 shows how telomeres in patients with variants in the known telomere genes (*DKC1*, *TERC*, *TERT* and *TINF2*) are significantly short when compared with a control population. Telomere lengths in all affected cases reported here were not significantly short when compared with controls regardless of the underlying genetic variant (Figure 1). This measure

should perhaps be used where a diagnosis of classical DC is suspected as a screening tool to decide the best course of action to determine a genetic diagnosis. Reduced telomere length has also been associated with an increased risk of cancer and DC is regarded as a cancer prone syndrome. Patients with *USB1* mutations tend to have an earlier presentation of myelodysplasia (MDS) than the general population as most reports describe disease in children. In this study and in our previous study<sup>21</sup> we report 6 patients with MDS and 2 with acute myeloid leukemia suggesting there is an increased cancer risk despite the patients not having short telomeres. Based on the small number of cases in our series, there does not appear to be an increased cancer risk for patients with either *GRHL2* or *LIG4* mutations. It is notable however that in the literature *LIG4* syndrome is described as a disorder with predisposition to leukemia. Although an increased cancer risk is not observed in our families this may be due to the small numbers of patients described herein.

Due to the extensive overlap in clinical features between DC and PN as described in Table 1, telomere length seems to be one way of separating these two diagnostic definitions. Given that these cases do not have short telomeres but do possess DC features (Table 1) present us with perplexing situation in accurately defining these patients. We propose it is timely to combine features of these two entities as a new syndrome, “*USB1* deficiency syndrome”, characterized by bone marrow failure, abnormal skin pigmentation (poikiloderma), nail dystrophy, growth restriction, cancer predisposition and normal length telomeres.

It is also notable that the genes mutated in these overlapping syndromes have similarities with the biological functions of the known DC genes (Figure 5). *USB1* has a role in snRNA processing, as is the case for dyskerin, *NHP2* and *NOP10*.<sup>31</sup> It also functions as an exoribonuclease reminiscent to the function of *PARN* that has been recently found to be mutated in both DC and pulmonary fibrosis.<sup>32</sup> Both *LIG4* and *RTEL1* are involved in the maintenance of genomic stability as they are both involved in pathways that repair double strand breaks. *LIG4* is an ATP-dependent DNA ligase that joins double strand breaks during non-homologous end joining and *RTEL1* (a protein that is mutated in a subset of DC) is an ATP-dependent DNA helicase, which has an important role in telomere-length regulation. Both function via interactions with another shelterin component *TRF2* (Telomere repeat binding factor 2).<sup>33,34</sup> Phenotypically many of the features seen in *LIG4*/Dubowitz syndrome are also seen in DC patients with biallelic *RTEL1* mutations (Table 2).<sup>7</sup> This supports the idea that mutations in *LIG4* can give rise to phenotype that is reminiscent of DC as caused by a gene that has a known association with telomere biology. *GRHL2* affects *TERT* expression by acting on its promoter<sup>35, 36</sup> thereby providing an important biological link to classical DC. While these proteins do have different functions or interactions, we speculate that it is this similarity as suggested in Figure 5 that gives rise to the overlapping phenotypes. It is also notable that many of the DC genes have functions other than telomere maintenance. It is therefore possible that those overlapping shared functions may also be contributing to the similar clinical features observed here. In the future it will be interesting to

determine whether therapeutic agents that work in one disease have a beneficial effect in any of the overlapping syndromes.

The findings of this study raise a potential issue as to what constitutes DC. Taking into account the clinical features, germline genetic mutation (s) and telomere length we can recognise three categories of patients: “Category 1”. Patients with sufficient clinical features (such as abnormal skin pigmentation, nail dystrophy, leukoplakia and bone marrow failure) to be labelled as DC, who have a germline mutation (s) in a telomere biology gene (such as *TERT*, *TERC*, *DKC1*, *TINF2*, *RTEL1* and so forth) and who have short telomeres (described as being less than 1<sup>st</sup> centile). “Category 2”. Patients with some clinical features of DC (but not sufficient to be classified as DC) who are found to have a germline mutation (s) in one of the telomere biology genes and who have short telomeres. “Category 3”. Patients with sufficient clinical features of DC but who harbour a germline mutation (s) in a non-telomere biology gene (such as *USB1*, *LIG4* and *GRHL2*) and have normal length telomeres. Category 1 patients can be considered to represent “classical or pure DC”. Category 1 and Category 2 patients can both be considered to represent disorders of telomeres – “telomeropathies”. Category 3 could be considered as “DC-like” or “DC-overlap”.

In summary, in nine out of the 12 families highlighted in this study, the biallelic variants identified in *USB1*, *GRHL2* and *LIG4* have been shown to be pathogenic in previous studies.<sup>13,14,20,23,24</sup> In the remaining families we have provided strong genetic and in-silico evidence that the identified mutations are pathogenic. This study has demonstrated that the pleotropic clinical phenotype of DC markedly overlaps with the recognised disease entities *LIG4* syndrome, Dubowitz syndrome and PN, as well as the recently recognised ECTDS. In doing so it has substantiated the last category as a significant disease entity. The marked overlap of features of DC with PN, ECTDS, *LIG4* and Dubowitz syndromes has important implications for establishing genetic diagnosis when a new patient presents in the clinic. Specifically in patients with a clinical diagnosis of DC the genetic analysis needs to include *GRHL2*, *LIG4* and *USB1* in addition to the classical DC genes.

All work in this study was conducted with informed, written consent and approval of the local ethics committee

## References

1. Pereboeva L, Westin E, Patel T, et al. DNA damage responses and oxidative stress in dyskeratosis congenita. *PLoS One*. 2013;8(10):e76473.
2. Kirwan M, Beswick R, Walne AJ, et al. Dyskeratosis congenita and the DNA damage response. *Br J Haematol*. 2011;153(5):634-643.
3. Dokal I, Vulliamy T, Mason P, Bessler M. Clinical utility gene card for: Dyskeratosis congenita - update 2015. *Eur J Hum Genet*. 2015;23(4).
4. Gramatges MM, Bertuch AA. Short telomeres: from dyskeratosis congenita to sporadic aplastic anemia and malignancy. *Transl Res*. 2013;162(6):353-363.
5. Savage SA. Human telomeres and telomere biology disorders. *Prog Mol Biol Transl Sci*. 2014;125:41-66.
6. Kocak H, Ballew BJ, Bisht K, et al. Hoyeraal-Hreidarsson syndrome caused by a germline mutation in the TEL patch of the telomere protein TPP1. *Genes Dev*. 2014;28(19):2090-2102.
7. Walne AJ, Vulliamy T, Kirwan M, Plagnol V, Dokal I. Constitutional mutations in RTEL1 cause severe dyskeratosis congenita. *Am J Hum Genet*. 2013;92(3):448-453.
8. Tummala H, Walne A, Collopy L, et al. Poly(A)-specific ribonuclease deficiency impacts telomere biology and causes dyskeratosis congenita. *J Clin Invest*. 2015;125(5):2151-2160.
9. Moon DH, Segal M, Boyraz B, et al. Poly(A)-specific ribonuclease (PARN) mediates 3'-end maturation of the telomerase RNA component. *Nat Genet*. 2015.
10. Dhanraj S, Gunja SM, Deveau AP, et al. Bone marrow failure and developmental delay caused by mutations in poly(A)-specific ribonuclease (PARN). *J Med Genet*. 2015;52(11):738-748.
11. Townsley DM, Dumitriu B, Young NS. Bone marrow failure and the telomeropathies. *Blood*. 2014;124(18):2775-2783.
12. Ng SB, Buckingham KJ, Lee C, et al. Exome sequencing identifies the cause of a mendelian disorder. *Nat Genet*. 2010;42(1):30-35.
13. Koparir A, Gezdirici A, Koparir E, et al. Poikiloderma with neutropenia: genotype-ethnic origin correlation, expanding phenotype and literature review. *Am J Med Genet A*. 2014;164A(10):2535-2540.
14. Petrof G, Nanda A, Howden J, et al. Mutations in GRHL2 result in an autosomal-recessive ectodermal Dysplasia syndrome. *Am J Hum Genet*. 2014;95(3):308-314.
15. Chistiakov DA. Ligase IV syndrome. *Adv Exp Med Biol*. 2010;685:175-185.
16. Stewart DR, Pemov A, Johnston JJ, et al. Dubowitz syndrome is a complex comprised of multiple, genetically distinct and phenotypically overlapping disorders. *PLoS One*. 2014;9(6):e98686.
17. Cawthon RM. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nuc Acids Res*. 2009;37((3)):e21.
18. Alter BP, Baerlocher GM, Savage SA, et al. Very short telomere length by flow fluorescence in situ hybridization identifies patients with dyskeratosis congenita. *Blood*. 2007;110(5):1439-1447.
19. Vulliamy TJ, Kirwan MJ, Beswick R, et al. Differences in disease severity but similar telomere lengths in genetic subgroups of patients with telomerase and shelterin mutations. *PLoS One*. 2011;6(9):e24383.
20. Alter BP, Giri N, Savage SA, Rosenberg PS. Telomere length in inherited bone marrow failure syndromes. *Haematologica*. 2015;100(1):49-54.
21. Walne AJ, Vulliamy T, Beswick R, Kirwan M, Dokal I. Mutations in C16orf57 and normal-length telomeres unify a subset of patients with dyskeratosis congenita, poikiloderma with neutropenia and Rothmund-Thomson syndrome. *Hum Mol Genet*. 2010;19(22):4453-4461.
22. Arnold AW, Itin PH, Pigors M, et al. Poikiloderma with neutropenia: a novel C16orf57 mutation and clinical diagnostic criteria. *Br J Dermatol*. 2010;163:866-869.
23. Clericuzio C, Harutyunyan K, Jin W, et al. Identification of a novel C16orf57 mutation in Athabaskan patients with Poikiloderma with Neutropenia. *Am J Med Genet A*. 2011;155A(2):337-342.
24. Hilcenko C, Simpson PJ, Finch AJ, et al. Aberrant 3' oligoadenylation of spliceosomal U6 small nuclear RNA in poikiloderma with neutropenia. *Blood*. 2013;121(6):1028-1038.

25. Balci S, Engiz O, Ereku A, Gozdasoglu S, Vulliamy T. An atypical form of dyskeratosis congenita with renal agenesis and no mutation in DKC1, TERC and TERT genes. *J Eur Acad Dermatol Venereol*. 2009;23(5):607-608.
26. Mlacki M, Kikulska A, Krzywinska E, Pawlak M, Wilanowski T. Recent discoveries concerning the involvement of transcription factors from the Grainyhead-like family in cancer. *Exp Biol Med (Maywood)*. 2015.
27. Aue A, Hinze C, Walentin K, et al. A Grainyhead-Like 2/Ovo-Like 2 Pathway Regulates Renal Epithelial Barrier Function and Lumen Expansion. *J Am Soc Nephrol*. 2015.
28. Chistiakov DA, Voronova NV, Chistiakov AP. Ligase IV syndrome. *Eur J Med Genet*. 2009;52(6):373-378.
29. Tamura S, Higuchi K, Tamaki M, et al. Novel compound heterozygous DNA ligase IV mutations in an adolescent with a slowly-progressing radiosensitive-severe combined immunodeficiency. *Clin Immunol*. 2015;160(2):255-260.
30. Murray JE, Bicknell LS, Yigit G, et al. Extreme growth failure is a common presentation of ligase IV deficiency. *Hum Mutat*. 2014;35(1):76-85.
31. Meier UT. The many facets of H/ACA ribonucleoproteins. *Chromosoma*. 2005;114(1):1-14.
32. Berndt H, Harnisch C, Rammelt C, et al. Maturation of mammalian H/ACA box snoRNAs: PAPD5-dependent adenylation and PARN-dependent trimming. *RNA*. 2012;18(5):958-972.
33. Chapman JR, Taylor MR, Boulton SJ. Playing the end game: DNA double-strand break repair pathway choice. *Mol Cell*. 2012;47(4):497-510.
34. Arnoult N, Karlseder J. Complex interactions between the DNA-damage response and mammalian telomeres. *Nat Struct Mol Biol*. 2015;22(11):859-866.
35. Chen W, Dong Q, Shin KH, et al. Grainyhead-like 2 enhances the human telomerase reverse transcriptase gene expression by inhibiting DNA methylation at the 5'-CpG island in normal human keratinocytes. *J Biol Chem*. 2010;285(52):40852-40863.
36. Kang X, Chen W, Kim RH, Kang MK, Park NH. Regulation of the hTERT promoter activity by MSH2, the hnRNPs K and D, and GRHL2 in human oral squamous cell carcinoma cells. *Oncogene*. 2009;28(4):565-574.
37. Patiroglu T, Akar HH. Clericuzio-type Poikiloderma with Neutropenia Syndrome in a Turkish Family: a Three Report of Siblings with Mutation in the C16orf57 gene. *Iran J Allergy Asthma Immunol*. 2015;14(3):331-337.
38. Kilic SS, Cekic S. Juvenile Idiopathic Inflammatory Myopathy in a Patient With Dyskeratosis Congenita Due to C16orf57 Mutation. *J Pediatr Hematol Oncol*. 2015;38(2):e75-77.

**Table 1: Overlap between DC, PN and patients with homozygous variants in *USB1***

Syndrome/ Family (OMIM)	DC (#305000, #127500, #224230)	PN (#604173)	Family 1	Family 2 <sup>37</sup>	Family 3 <sup>38</sup>	Family 4	Family 5	Family 6	Family 7	Family 8
Consanguinity			Yes			Yes	‡	Yes		Yes
Country of Origin			Turkey	Turkey	Turkey	Turkey	Turkey	Australia	Afghanistan	Afghanistan
Family history			Yes	Yes		Yes				
Age at report (years)			24,14	12,7,5	7	9, 7	6	10	4	16
Mutation	Various*	USB1	p.H179MfsTer86	p.H179MfsTer86	p.H179MfsTer86	p.H179MfsTer86	p.H179MfsTer86	p.Q225Ter	p.H208R	p.H208R
Nail dystrophy	>60% cases**	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Abnormal skin pigmentation	>60% cases	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
BM failure	>60% cases	Yes	Yes <sup>a</sup>	Yes <sup>b</sup>	Yes <sup>c</sup>	Yes <sup>d</sup>	Yes <sup>e</sup>	Yes <sup>f</sup>	Yes <sup>g</sup>	Yes <sup>h</sup>
Leukoplakia	>40% cases		Yes				Yes			Yes
Developmental delay	>20% cases		Yes	Yes		Yes				
Microcephaly	>20% cases					Yes				
Growth restriction	>20% cases	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Hair loss	>20% cases					Yes		Yes		
Pulmonary disease	>10% cases	Yes	Yes		Yes	Yes	Yes			
Cancer	~10% cases	~10% cases						MDS		
Abnormal dentition	<10% cases	Yes	Yes	Yes				Yes		
Gonadal abnormalities	<10% cases	Yes				Yes				
Deafness/ear abnormalities	<10% cases				Yes			Yes		
Eye abnormalities	<10% cases							Yes		
Abnormal facies	<10% cases	Yes		Yes	Yes			Yes		Yes
Skeletal abnormalities	<10% cases							Yes		
Immune deficiency	<10% cases		Yes				Yes		Yes	Yes
Short telomeres	<1 <sup>st</sup> centile	Unknown	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal

All clinical features reported within a family are detailed. DC-Dyskeratosis congenita; PN-poikiloderma with neutropenia; \* Mutations identified in *CTCI*, *DKC1*, *NHP2*, *NOPI0*, *PARN*, *RTEL1*, *TERC*, *TERT*, *TINF2*, *TPP1* and *WRAP53*; \*\* as calculated from the index cases of the 1<sup>st</sup> 400 families included in the DCR, London; †Term used to cover short stature, intra uterine growth restriction and low birth weight; ‡Parents reported as being from the same village but no statement of consanguinity. The blood counts for the index case at presentation are as follows: a-hemoglobin 106g/l, leucocytes  $2.3 \times 10^9/l$ , neutrophils  $0.3 \times 10^9/l$ , platelets  $246 \times 10^9/l$  and bone marrow showed decreased cellularity; b-hemoglobin 125g/l, leucocytes  $3.2 \times 10^9/l$ , neutrophils  $0.1 \times 10^9/l$ , platelets  $311 \times 10^9/l$  and bone marrow showed decreased cellularity with approximately 10% blasts; c-hemoglobin 113g/l, leucocytes  $3.2 \times 10^9/l$ , neutrophils  $0.8 \times 10^9/l$  platelets  $308 \times 10^9/l$ ; d-hemoglobin 115g/l, leucocytes  $3.4 \times 10^9/l$ , neutrophils  $0.8 \times 10^9/l$ , platelets  $138 \times 10^9/l$ ; e-hemoglobin 121g/l, leucocytes  $3.68 \times 10^9/l$ , neutrophils  $0.7 \times 10^9/l$ , platelets  $269 \times 10^9/l$ ; f-hemoglobin 70g/l, leucocytes  $2.1 \times 10^9/l$ , neutrophils  $1.1 \times 10^9/l$ , platelets  $5 \times 10^9/l$  and bone marrow showed increased cellularity, decreased magakaryocytes, dysmyelopoiesis with nuclear changes and abnormal granularity; g-hemoglobin 115g/l, leucocytes  $4.5 \times 10^9/l$ , neutrophils  $0.9 \times 10^9/l$ , platelets  $204 \times 10^9/l$ ; h-hemoglobin 114g/l, leucocytes  $1.7 \times 10^9/l$ , neutrophils  $0.8 \times 10^9/l$ , platelets  $112 \times 10^9/l$ . MDS- myelodysplastic syndrome.



**Table 2: Clinical overlap between the different genetic syndromes and the families with biallelic variants identified in this study**

Syndrome/Family (OMIM reference)	DC (#305000, #127500, #224230)	ECTDS (#616029)	Family 9	Family 10 <sup>25</sup>	LIG4 syndrome (#606593)	Dubowitz syndrome (#223370)	Family 11	Family 12
Consanguinity			Yes	Yes				
Country of origin			Kuwait	Turkey			Guadeloupe	UK
Family history			Yes					
Age at report (years)			10, 5	27			5	5
Mutated gene (variant identified)	Various*	<i>GRHL2</i> (p.Y398H, p.I482K)	<i>GRHL2</i> (p.I482K)	<i>GRHL2</i> (p.P405T)	<i>LIG4</i> (various)	<i>LIG4</i> (p.R814Ter, p.S205LfsTer29)	<i>LIG4</i> (p.R814Ter, p.K424RfsTer20)	<i>LIG4</i> (p.R814Ter, p.K424RfsTer20)
Nail dystrophy	>60% cases**	Yes	Yes	Yes				
Abnormal skin pigmentation	>60% cases	Yes	Yes	Yes	Yes		1 café au lait spot	
BM failure	>60% cases		No <sup>a</sup>	Yes <sup>b</sup>	Yes	Yes	Yes <sup>c</sup>	Yes <sup>d</sup>
Leukoplakia	>40% cases	Yes	Yes	Yes				
Developmental delay	>20% cases				Yes	Yes		
Microcephaly	>20% cases				Yes	Yes	Yes	Yes
Growth restriction†	>20% cases	Yes	Yes	Yes	Yes	Yes	Yes	Dwarfism
Abnormal dentition	<10% cases	Yes	Yes	Yes		Yes		
Oesophageal stricture	<10% cases	Yes						
Gonadal abnormalities	<10% cases			Yes	Yes	Yes		
Ataxia	<10% cases							
Immune deficiency	<10% cases				Yes		Yes	Yes
Deafness/ear abnormalities	<10% cases	Yes						
Kidney abnormalities	<10% cases		Recurrent infections	Renal agenesis				
Abnormal facies	<10% cases				Yes	Yes	Yes	
Short telomeres	<1 <sup>st</sup> centile	Unknown	Normal	Normal	Unknown	<1 <sup>st</sup> centile <sup>e</sup>	Normal	Normal

All clinical features reported within a family are detailed. ECTDS – Ectodermal dysplasia/short stature syndrome. \* Mutations identified in *CTC1*, *DKC1*, *NHP2*, *NOP10*, *PARN*, *RTKL1*, *TERC*, *TERT*, *TINF2*, *TPP1* and *WRAP53*;\*\* as calculated from the index cases of the 1<sup>st</sup> 400 families in the DCR, London; †Term used to cover short stature, intra uterine growth restriction and low birth weight. The blood counts for the index case at presentation are as follows: a - hemoglobin 141g/l, leucocytes 7.2x10<sup>9</sup>/l, neutrophils 4.4x10<sup>9</sup>/l, platelets 307x10<sup>9</sup>/l (normal blood counts); b-hemoglobin 89g/l, leucocytes 4.9x10<sup>9</sup>/l, platelets 216x10<sup>9</sup>/l and bone marrow showed reduced cellularity with megaloblastic changes; c-hemoglobin 96g/l, neutrophils 0.5x10<sup>9</sup>/l, platelets 20x10<sup>9</sup>/l and bone marrow showed reduced cellularity and reduced megakaryocytes; d-pancytopenia with B-lymphopenia and bone marrow showed reduced cellularity. e- reported in 3 patients only

## Figure Legends

### Figure 1: Telomere lengths in patients with *GRHL2*, *LIG4* and *USB1* mutations are not short

Relative T/S ratios are within the normal range. Open circles – controls (n=130); green diamonds – patients with biallelic *USB1* variants (n=12), blue triangles – patients with homozygous *GRHL2* variants (n=2); Inverted red triangles – patients with compound heterozygous *LIG4* variants (n=2). For comparison short telomeres are seen in patients with mutations in the known telomere associated genes *DKC1* (n=50, open squares), *TERC* (n=66, open triangles), *TERT* (n=36, open inverted triangles) and *TINF2* (n=33, open diamonds). Black line represents the median for each data set.

### Figure 2: Identification and segregation of disease-causing variants in *USB1*

- A) The c.531delA variant is identified in five families with a DC phenotype. Representative sequencing traces are shown for the wild type (+/+), heterozygous (+/-) and homozygous (-/-) forms of the mutation.
- B) Family tree and sequence traces of c.673C>T variant seen in Family 6. A representative heterozygous trace and the homozygous trace are shown.
- C) Family tree and sequence traces of the novel c.623A>G homozygous variant (G/G) as seen in Family 7 and 8. A representative wild type (A/A) trace is also shown. NA – sample not available.
- D) Protein crystal structure of human *USB1* (PDB id: 4W7H) depicted in ribbon form indicating the both transit and terminal lobes indicated in green  $\alpha$  helices. The position of identified residues mutated in *USB1* deficiency patients is indicated as stick model in blue.

E and F) In silico modelling of truncation mutations reveals loss several  $\alpha$  helices and  $\beta$  sheets in both transit and terminal lobes of *USB1* secondary structure.

G and H) The invariant H-x-S motif in the active site is indicated as stick model. The missense change Histidine to Arginine at 208 position (red) introduces an extra hydrogen bond (dotted line black indicated by arrows), with Ser210 (blue), which in turn might cause aberrant oligoadenylation of RNA substrate.

### Figure 3: Similar clinical presentation is observed between DC and PN highlighting the marked overlap between these two syndromes

Panels A, D, G are from a male patient with p.Ala353Val *DKC1* mutation, panels B, E and H are from a male patient with the homozygous *USB1* p.His127MetfsTer86 mutation and panels C, F I are from a child with the homozygous *USB1* p.His208Arg mutation. Panels A-C show similar patterns of reticular pigmentation on the legs. Panels D-F show nail dystrophy on the toes. Panels G and H show reticular pigmentation on the trunk and panel I shows pigmentation on the arm along with nail dystrophy on the fingers. It is notable that the mucocutaneous features of the patient with *USB1* p.His179MetfsTer86 mutation appear more like those of the patient with *DKC1* p.Ala353Val mutation than the one with the *USB1* p.His208Arg mutation.

**Figure 4: Identification of disease-causing variants in two different genes in patients with clinical features of DC**

- A) Segregation and sequence traces for the homozygous *GRHL2* variants identified in Families 9 and 10. Representative heterozygous and homozygous traces are shown for both variants. Schematic representation of the protein shows the identified domains and the location of the variants. Red arrow highlights the variant identified in this study. Black arrow highlights a variant identified in previous study. Green arrow highlights the variant observed in both this and previous study. NA-sample not available.
- B) Segregation and sequence traces for the compound heterozygous *LIG4* variants identified in Families 11 and 12. A representative wild type and heterozygous trace is shown for each. The nucleotides deleted are indicated by a line above the corresponding bases on the wild type (+/+) trace. Schematic representation of the protein showing the precise location of variants in different domains. Black arrows highlight variants identified in previous studies. Green arrows highlight variants observed in both this and previous studies.

**Figure 5: Biological overlap of classical DC proteins (TERT, DKC1, NOP10, NHP2, PARN and RTEL1) and those (GRHL2, USB1 and LIG4) mutated in patients in this study**

Proteins with similar functions are grouped together. The schematic shows that GRHL2 can be grouped broadly with the molecules which have role in telomerase expression, USB1 with small nuclear ribonucleoprotein (snRNP) processing and LIG4 with genomic stability. Previous studies have shown that biallelic mutations in *GRHL2*, *USB1* and *LIG4* are associated with ectodermal dysplasia, poikiloderma with neutropenia (or USB1 deficiency syndrome) and Lig4/Dubowitz syndrome, respectively. Mutations in DKC1, NHP2, NOP10, PARN and RTEL1 have been linked to dyskeratosis congenita.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

