

Response to treatment with TNF α inhibitors in rheumatoid arthritis is associated with high levels of GM-CSF and GM-CSF⁺ T lymphocytes

Jonas Bystrom¹, Felix I. Clanchy², Taher E. Taher¹, Mohammed Al-Bogami¹, Hawzheen Mohammed¹, Saba Alzabin², Pamela Mangat³, Ali S. Jawad¹, Richard O. Williams², Rizgar A. Mageed¹

¹Experimental Medicine and Rheumatology, William Harvey Research Institute, Queen Mary University of London, London; ²Kennedy Institute of Rheumatology, Oxford University, Oxford and ³Department of Rheumatology, Royal Free Hospital, NHS Foundation Trust London, UK

Running title: High GM-CSF levels predict responsiveness to anti-TNF

Address for correspondence and reprint requests:

Dr. Jonas Bystrom,
Centre for Experimental Medicine and Rheumatology,
William Harvey Research Institute,
Barts and the London School of Medicine and Dentistry,
Queen Mary University of London,
Charterhouse Square,
London EC1M 6BQ

Email: j.bystrom@qmul.ac.uk
Tel: +44 (0)20 7882 2473
Fax: +44(0)20 7882 8252

Abstract

TNF α inhibitors are a mainstay treatment option for patients with rheumatoid arthritis (RA) refractory to other treatment options. However, many patients either do not respond, or relapse after initially responding to these agents. This study was carried out to identify biomarkers that distinguish responder from non-responder patients before treatment. The level of cytokines in plasma and those produced by T cells, B cells and monocytes in 97 RA patients treated with TNF α inhibitors was measured before and after 1 and 3 months of treatment by multiplex analyses. The frequency of T cell subsets and intracellular cytokines were determined by flow cytometry. The results reveal that pre-treatment, T cells from patients who went on to respond to anti-TNF α produced significantly more GM-CSF than non-responders. Furthermore, responders produced higher levels of IL-1 β , TNF α and IL-6. Cytokine profiling confirmed the association between high levels of GM-CSF and responsiveness to anti-TNF α . Thus, high plasma levels of GM-CSF pre-treatment had a positive predictive value of 87.5% (61.6% to 98.5% at 95% CI). The study also show that cells from most anti-TNF α responder patients in the current cohort produced higher levels of GM-CSF and TNF α pre-treatment than non-responder patients. Findings from the current study and our previous observations that non-responsiveness to anti-TNF α is associated with high IL-17 levels suggest that RA in responder and non-responder patients could be driven/sustained by different inflammatory pathways.

Introduction

Rheumatoid arthritis (RA) is a debilitating disease characterised by autoimmunity and the production of high levels of pro-inflammatory cytokines resulting in joint inflammation, the production of matrix-degrading enzymes and cartilage and bone destruction [1]. Epidemiological and genetic studies indicate that susceptibility to RA is associated with HLA-DR alleles and genes encoding cytokines and proteins that regulate T cell responses, highlighting the importance of T cells in disease pathogenesis [2].

Knowledge of the principal pro-inflammatory mediators that cause synovial inflammation in RA led to the discovery and use of biological TNF α inhibitors as effective therapies for treating patients [3]. Most treated patients undergo remission, especially if anti-TNF α is administered soon after diagnosis [4,5]. However, a significant proportion of patients established disease either do not respond to the treatment or relapse after an initial response, suggesting that the disease in these patients is not driven by TNF α . Exactly why some RA patients do not respond to TNF α inhibitors while others relapse after an initial response remains unclear. One possible explanation for the inconsistent or incomplete response of RA patients to anti-TNF α is that TNF α , perhaps paradoxically, also has regulatory functions. For example, although there is good evidence that the activity of regulatory T cells improves following treatment with anti-TNF α [6-8], there is also evidence that TNF α modulates TCR-mediated signalling [9,10]. An alternative explanation may be that chronic inflammation and disease processes in non-responder patients are not wholly driven by TNF α . In this regard, a number of laboratories including ours have revealed that T cells from non-responder RA patients produce high levels of IL-17 and have a high frequency of circulating Th17 cells [11,12]. Moreover, our studies have revealed that treating patients with TNF α inhibitors increases the frequency of Th17 cells and IL-17 production, possibly through increasing IL-

12/IL-23 p40 production by monocytes/macrophages [12]. Although the molecular basis and the physiological and therapeutic implications remain to be determined, these findings indicate that TNF α /TNF α blockade have significant impacts on the function and frequency of T cell subsets [13]. This suggestion is consistent with the association between T cell responses and signalling generated by TNF α through its two receptor in T cells [reviewed in 14].

The available evidence suggests that mechanisms driving inflammation in RA could be different in different patients as indicated by variations in the response of individual patients to treatment with anti-TNF α (7,11,12). Therefore, a major challenge has been in the identification of alternative pathways that drive disease in non-responder patients before treatment with TNF α inhibitors starts to enable better, evidence-based therapeutic use of these biological agents. A large number of studies from many laboratories explored the predictive value of many biomarkers including cytokines produced by peripheral blood mononuclear cells or the transcriptome prior to treatment (summarized in Table 1). In the current study we assess the phenotype and response of T cell subsets in RA patients prescribed anti-TNF α before treatment in relation to the patient's subsequent response/lack of response to treatment. The results provide evidence for relationships between response/lack of response to anti-TNF α and the frequency of T cell subsets and patterns of plasma levels of pro-inflammatory cytokines prior to treatment.

Patients and Methods

Patients

Patients were recruited from rheumatology clinics at Barts Health NHS, the Royal Free London NHS Foundation and Imperial College Healthcare Trusts with patients' informed consent. The study was approved by the Ethical Committee (06/Q0605/8; NRES Committee London - City & East) and conducted in compliance with the Declaration of Helsinki 2013. The patients were prescribed anti-TNF α based on NICE guidelines which include failure to respond to treatment with 2 disease-modifying agents one of which is methotrexate and with a DAS28 score >5.1. Patients were treated with adalimumab, certolizumab pegol, golimumab or etanercept in combination with methotrexate. Patients' response to treatment was based on the European League Against Rheumatism (EULAR) response criteria [28]; patients were considered responders when their DAS28 decreased by >1.2 to \leq 3.2 after 3 months of treatment. All other treated patients were considered non-responders. Immediately prior to treatment and after 1 and 3 months of treatment, 25mL blood was drawn and B and T cells and monocytes enriched and phenotypically characterised by FACS, stimulated and cultured. Plasma were separated and stored at -80°C until tested.

Reagents

PerCP-Cy5.5-conjugated anti-TNF α , eFluor® 660 anti-GM-CSF, PE-Cy7 anti-IFN γ , PE anti-IL-17A, FITC-anti-CD45RA, PE-anti-CD45RO, eFluor® 450-anti-CD161 and unconjugated anti-CD28 antibodies (Abs) were from eBioscience. Brilliant Violet 605-anti-CCR6 and APC-anti-IL-10 Abs were purchased from BioLegend. RosetteSep negative-enrichment kits for human T cells, B cells and monocytes were from StemCell Technologies. Ficoll-Paque was from GE Healthcare, RPMI from Lonza and foetal bovine serum from Life Technologies. Lipopolysaccharide (E. coli, serotype 0111:B4), Ionomycin and PMA were

from Sigma-Aldrich and GolgiplugTM was from BD Bioscience and Leucoperm from AbD Serotec.

Cell enrichment and analysis

Blood T and B cells and monocytes were enriched by negative selection with RosetteSep kits using mononuclear cells separated by Ficoll-Paque centrifugation. The purity of enriched T cells was consistently $\geq 95\%$ while for monocyte and B cells purity was 86-98%. The cells were suspended in RPMI containing 10% FCS and antibiotics and analysed by FACS or stimulated *in vitro*. T cells were stimulated with 10 $\mu\text{g}/\text{mL}$ anti-CD3 mAb (clone OKT3) pre-coated onto sterile tissue culture plates and 10 $\mu\text{g}/\text{mL}$ anti-CD28 (clone 28.2) in the medium. B cells were stimulated with 10 $\mu\text{g}/\text{mL}$ goat F(ab')₂ anti-human IgM coated onto wells of the plates and 10 $\mu\text{g}/\text{mL}$ antibody to CD40 (clone G28-5) in the medium. Monocytes were stimulated with 1 $\mu\text{g}/\text{mL}$ LPS (E. coli, serotype 0111:B4). Supernatants were collected after 48hrs, debris removed by centrifugation and cytokines quantified using Meso Scale Discovery (MSD) multiplex kits or ELISA. To verify that levels of TNF α produced by T cells and monocytes were representative of what the cells produce independent of the stimulation protocol, T cells and monocytes from 3 healthy controls (HCs) were also stimulated independently for 48hr with 0.1 $\mu\text{g}/\text{mL}$ PMA and Ionomycin. In addition, T cells were stimulated with 0.1 $\mu\text{g}/\text{mL}$ PMA and Ionomycin in the presence of 1 $\mu\text{l}/\text{mL}$ GolgiplugTM (BD Bioscience) overnight to determine the level of intracellular cytokines. The cells were stained for CD45RA, CD45RO, CCR6 and CD161 by incubation with conjugated mAbs for 20mins at 4°C, permeabilized using Leucoperm (AbD Serotec) and stained for intracellular cytokines with labelled mAbs for IL-17, INF γ , TNF α , GM-CSF and IL-10. The cells were fixed with 0.5% p-formaldehyde in PBS and analysed. FACS was performed using a BD-

LSR Fortessa X20 (BD Biosciences) and cell populations gated and assessed using FACS Diva 6.0 software (BD Bioscience).

Cytokine measurements

For quantification of cytokines, multiplex MSD kits or sandwich ELISA were used. Conditions for cytokine measurements including determining optimal dilutions of different clinical samples were carried out prior to the measurements as reported [12]. Level of the following cytokines/chemokines were determined in culture supernatants of *ex vivo* stimulated T and B cells and monocytes and in plasma: TNF α , IL-17, IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-13, IL-20, IL-22, IL-23p40, GM-CSF, IFN γ and MCP-1.

Statistical analyses

Comparison of patterns of cytokine production was with the Multi Experiment Viewer software (Dana Faber Cancer Institute, Boston, MA). Hierarchical clustering was with Pearson's correlation with complete linkage. Where indicated, statistical analyses were carried out using the GraphPad Prism 6 software. The Mann-Whitney *U* test, Wilcoxon matched-pairs, signed rank test, Fisher's exact test or the Chi² test were used for the analysis of differences between or within groups, as appropriate. Positive and negative predictive values were calculated using the online program https://www.medcalc.net/tests/diagnostic_test.php.

Results

Patient response to therapy

Ninety seven RA patients prescribed anti-TNF α were recruited to the study (Table 2) and their clinical samples used in multiple experiments described in this report. The patients were treated with one of the following 4 anti-TNF α agents: adalimumab, certolizumab pegol, golimumab or etanercept in combination with methotrexate. No notable or consistent differences in cytokine production by enriched cells were seen between patients receiving different anti-TNF α agents. Assessment of patients' response to anti-TNF α was based on the EULAR response criteria at 3 months after treatment. Seventy six patients (78%) responded to the treatment, slightly higher than previously reported [6-8, 11].

Distinct pro-inflammatory T cell cytokine profiles predict responsiveness or lack of responsiveness to treatment with anti-TNF α

In a previous study we determined that prior to treatment with anti-TNF α , non-responder RA patients had high frequencies of IL-17⁺ T cells and, when their T cells stimulated *in vitro*, the cells produced significantly higher levels of IL-17 than responder patients [12]. The current study compared the profile of cytokines produced by enriched T and B cells and monocytes in responder and non-responder patients prior to and after the start of treatment to establish whether distinct inflammatory pathways associate with the disease in responder as compared with non-responder patients. Levels of TNF α , IL-17, IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-13, IL-20, IL-22, IL-23p19, GM-CSF, IFN γ and MCP-1 either in culture supernatants of the stimulated T and B cells and monocytes or detected in plasma collected from most of the patients at the time of sampling were measured. For clarity of our observations, only data for cytokine measurements that consistently showed significant

differences between responder and non-responder patients are presented and discussed in detail in this report.

Monocytes are generally believed to be a main source of TNF α in joints of RA patients although there is evidence that the cytokine is also produced by T cells [29-32]. The current study confirmed that monocytes, T and B cells all produce TNF α when stimulated *in vitro* (Figure 1A). However, T cells stimulated with anti-CD3/CD28 mAbs produced almost 45 folds the level of TNF α produced by LPS-stimulated monocytes ($P<0.0001$; Figure 1A and Table 3). To exclude the possibility that the higher levels of TNF α produced by T cells was merely due to the activation protocol and not capacity of the cells themselves, both lineage cells were independently stimulated with PMA/Ionomycin in parallel experiments. The mean and SEM level of TNF α produced by T cells stimulated with PMA/Ionomycin was $5.9\pm 4.3\text{ng/mL}$, >10-folds higher than what monocytes produced with PMA/Ionomycin at $0.5.8\pm 0.4\text{ng/mL}$ ($P<0.05$; Figure 1B). Before treatment, T cells from responder patients stimulated with anti-CD3/CD28 mAbs tended to produce more TNF α than non-responder patients ($P=0.20$; Figure 1A and Table 3).

T cells from responder patients produced seven folds higher levels of IL-1 β than T cells from non-responders (Figure 1C and Table 3). However, monocytes produced higher levels of IL-1 β than T cells but there was no difference in IL-1 β production by monocytes from responder and non-responder patients. As expected, B cells produced less IL-1 β than monocytes but there was a trend for higher levels in responder patients (Figure 1C and Table 3).

T cells produced significantly higher levels of GM-CSF compared with monocytes and B cells ($P<0.0001$; Figure 1D and Table 3). Importantly, GM-CSF levels produced by T cells were significantly higher in responder compared with non-responder patients before treatment ($P<0.05$; Figure 1D and 1E and Table 3). There was a significant increase in GM-

CSF production by T cells in responder patients after 1 month of treatment but declined at 3 months ($P<0.05$, Figure 1E). Consistent with higher GM-CSF levels produced by T cells from responder patients, there were also a trend for more GM-CSF⁺ T cells in the blood of responder patients before (3.0 ± 2.0 vs. $1.4\pm 1.4\%$) and after 1 month (3.1 ± 2.0 vs. $1.4\pm 1.0\%$) of treatment although these differences were not statistically significant (Figure 1F).

Monocytes, T and B cells produced IL-6 with monocytes producing 10-folds higher levels than T and B cells ($P<0.0001$; Table 3). T cells from responder patients produced >3 folds IL-6 levels more than T cells from non-responder patients before treatment ($P=0.27$, Figure 1G and Table 3).

MCP1 was produced almost exclusively by monocytes with 1.8 folds higher levels in non-responders compared with responders before and after 1 month of treatment. The difference increased to 2.3 folds after 3 months of treatment ($P<0.05$, Figure 1H and Table 3).

Since GM-CSF, TNF α and IL-1 β produced by T cells were higher in responder patients compared with non-responder patients, we carried out a cluster analysis to assess their predictive value of responsiveness to anti-TNF α pre-treatment. A 2-dimensional clustering was carried out with Pearson's correlation and complete linkage analysis (Figure 2A). Two clusters emerged, one containing patients with high levels of GM-CSF, IL1 β and TNF α (connected with a blue bracket) and the second containing patients with low levels of the cytokines (connected with a red bracket). Eighteen of 19 patients within the first cluster (94.7%) were responders and only one was a non-responder. The second cluster contained 19 non-responders (57.6%) and 14 responders (42.4%) (Figure 2C). The relative prevalence of responders to non-responders in each cluster was significantly different (Fisher's Exact Test, $P<0.001$). Inclusion of the other T cell-derived cytokines measured in this study did not improve the clustering and were, therefore, not included in this analysis.

High plasma GM-CSF levels confirms association with responsiveness to anti-TNF α

To explore the feasibility of developing a strategy for identifying likely responders to anti-TNF α before treatment based on the findings from the current study, we also assessed plasma levels of TNF α , IL-1 β , IL-6, GM-CSF and MCP-1 pre-treatment (Figures 2B and 3A-E). There were no differences between responder and non-responder patients in pre-treatment plasma levels of TNF α , IL-6 or MCP-1. However, high pre-treatment levels of GM-CSF were significantly associated with responsiveness to anti-TNF α (6.4 ± 4.6 pg/mL vs. 2.8 ± 2.0 pg/mL, responders vs. non-responders; $P<0.05$; Figure 3D). To assess the potential application of measuring GM-CSF in plasma and/or culture supernatants of T cells before treatment to predict responsiveness, we carried out a Receiver Operated Characteristics (ROC) analysis by plotting the true positive rate (Sensitivity) as a function of the false positive rate (100-Specificity). The analyses revealed that the level of GM-CSF in culture supernatants of activated T cells gave a predictive value, area under curve (AUC), of 0.70, (0.56-0.85 at 95% confidence interval [CI]; $P<0.05$), and for plasma an AUC of 0.73 (0.53-0.90% at 95% CI; $P=0.09$; Figure 2D). Calculation of odds ratio (OR) showed that responder patients had an OR of 3.9 to respond to treatment at GM-CSF levels of >2.5 pg/mL produced by 200 cultured T cells and an OR of 11.7 at plasma levels of >3.5 pg/mL compared with non-responders (Figure 2E). The predictive value of GM-CSF in culture supernatants of T cells had a specificity of 73.3% and a sensitivity of 58.3% (40.8-74.5 at 95% CI). For plasma GM-CSF, specificity was 71.4% (29.0-96.3% at 95% CI) and sensitivity 82.4% (56.6-96.2% at 95% CI). The positive predictive values, at cut-offs as stated above, for GM-CSF in culture supernatants of T cells was 84.0% (63.9% to 95.5% at 95% CI) and for plasma 87.5% (61.6% to 98.5% at 95% CI).

The phenotype and activation status of GM-CSF⁺ T cells in RA patients

There is evidence that GM-CSF can be produced by CD161⁺IFN γ ⁺ ex-Th17 cells in the joint of some RA patients and by peripheral blood Th17 cells in patients with multiple sclerosis [33,34]. Other studies, however, have shown that GM-CSF is produced by T cells other than Th17 cells [35,36]. To assess the existence of links between GM-CSF-producing T cells and Th17 cells in our cohort of patients in relation to responsiveness to anti-TNF α , T cells were enriched from the blood of 7 responder and 3 non-responder patients, stimulated with PMA/Ionomycin for 16 hours, stained for membrane proteins and intracellular cytokines and analysed by FACS. We used CCR6 expression, a membrane protein expressed on the majority of IL-17⁺ T cells, to identify Th17 cells in the patients (Figure 4A). To determine whether GM-CSF⁺ T cells overlapped with Th17 cells we gated on CCR6⁺IL-17⁺ T cells within the enriched/stimulated CD3⁺ T cells. The FACS contour plots first confirmed that almost all of the IL-17⁺ T cells expressed CCR6 (2.2% of all CD3⁺ T cells; Figure 4A). The gated CCR6⁺IL-17⁺ T cells were then analysed for membrane CD161 and intracellular TNF α , GM-CSF and IFN γ expression. This analysis showed that of the CCR6⁺IL-17⁺ T cells, 58.0 \pm 12.7% and 57.0 \pm 16.1% expressed CD161, 19.6 \pm 11.7% and 41.9 \pm 33.7% expressed IFN γ , 79.3 \pm 9.6% and 85.3 \pm 2.5% expressed TNF α but only 12.3 \pm 4.8% and 13.6 \pm 6.6% expressed GM-CSF in responder and non-responder patients, respectively. In contrast, analysis of GM-CSF⁺ T cells showed that almost all co-expressed TNF α ⁺ (example of FACS contour plots in Figure 4B). The GM-CSF⁺TNF α ⁺ T cells (4.9% of all CD3⁺ T cells) were then gated and analysed for membrane CD161 and intracellular IFN γ and IL-17 expression (Figure 4B). This analysis showed that of the GM-CSF⁺TNF α ⁺ T cells 12.1 \pm 7.0% and 20.1 \pm 4.9% expressed CD161, 32.2 \pm 12.2% and 45.4 \pm 14.9% expressed IFN γ and 3.7 \pm 1.0% and 5.0 \pm 1.4% expressed IL-17 in responders and non-responders, respectively (Figure 4B). Differences between responder and non-responder patients were not, however, statistically

significant ($P>0.05$). Further analysis of GM-CSF⁺ T cell phenotypes in 20 patients showed that they were mostly of the CD45RO⁺ effector/memory phenotype (Figure 4C). As revealed in Figure 4B, a relatively small proportion expressed low levels of CD161 but none expressed IL-10 (not shown) indicating that they were unlikely to be regulatory T cells (Tregs) [37,38]. Thus, although a few Th17 cells produced GM-CSF, the majority did not ($P<0.0001$). Likewise, only few cells within the GM-CSF⁺ TNF α ⁺ T cell population produced IL-17 ($p<0.0001$). Furthermore, when the level of IL-17 and GM-CSF from *in vitro* anti-CD3/CD28-activated T cell culture supernatants were compared in a subgroups of the patients, none of the patients simultaneously produced high level of both cytokines (Figure 4D).

Discussion

This study provides evidence that the response/lack of response of RA patients to anti-TNF α therapy is associated with differences in the frequency of T cell subsets in patients. Thus, the results show that responder patients tend to have a high frequency of T cells that produce GM-CSF in contrast to non-responders that have a high frequency of IL-17-producing T cells at the start of treatment with anti-TNF α . Furthermore, clustering analysis of the pro-inflammatory cytokines TNF α , GM-CSF and IL-1 β suggest that the disease in responder and non-responder patients is likely to be driven/sustained through different inflammatory pathways. The study also suggests that T cells could be the major producers of TNF α and GM-CSF in RA patients. Consistent with the association between high frequency of GM-CSF⁺ T cells that produce high levels the cytokine when activated *in vitro* and responsiveness to TNF α inhibitors is that high plasma level of GM-CSF are also detected in the patients. High level of IL-17 prior to treatment, in contrast, has an inverse relationship with responsiveness to anti-TNF α [11,12].

About 30% of RA patients prescribed anti-TNF α do not respond adequately and a further 50% of responders relapse within 12 months of treatment [39]. Previous studies from our laboratory have revealed an association between the lack of response to anti-TNF α and high levels of IL-17-producing T cells pre-treatment [12]. These studies also showed an increase in the frequency of Th17 cells after treatment. This latter observation could be related to the fact that TNF α also has immunomodulatory functions [40]. For example, TNF α has been shown to modulate diabetes, EAE, lupus and arthritis in animal models, possibly, by inhibiting Th17 cells [41-44]. Importantly, findings from the current study show that T cells could be a major source of TNF α *in vivo*. This finding is consistent with studies showing that both macrophages and T cells in the synovial lining tissue produce TNF α [30,32]. Since T

cells and monocytes express TNF α receptors, the cytokine could act directly on these cells and modulate their differentiation and functions [14,44]. Indeed, blockade of TNF α in arthritic mice led to the expansion of collagen-specific Th1 and Th17 cells and our previous studies showed that this effect is likely to be due to increased IL-12/IL-23 p40 production by monocytes and antigen-presenting cells [12,44].

There is clearly an unmet need to identify biomarkers for evidence-based prescription of biological therapies in all RA patients. Identifying such biomarkers has been the subject of much research by many investigators. Several potential biomarkers have been reported (Table 1) but the application of such biomarkers in larger populations beyond the cohorts studied is unknown. For example, anti-cyclic citrullinated peptide antibodies and rheumatoid factors, autoantibodies traditionally associated with RA, were studied but ultimately shown to be of limited value [45,46]. We propose that a combination of reported biomarkers, for example GM-CSF and the inflammatory protein MRP8/14 or certain micro-RNAs, could predict responsiveness to anti-TNF α in most RA patient cohorts more reproducibly [22,26].

GM-CSF is an important cytokine for the maturation of macrophages and CD1⁺ pro-inflammatory dendritic cells [36,47] as well as in myelopoiesis [48]. Interestingly, treatment with GM-CSF has been shown to precipitate arthritis in some cancer patients [49]. Nevertheless, although GM-CSF has well-characterised pro-inflammatory properties, it is not established if GM-CSF⁺ T cells are always pathogenic [50]. Thus, the cytokine is associated with a better prognosis in some animal models of auto-immune diseases. For example, GM-CSF has been shown to enhance the ability of dendritic cells to promote Treg cell activity and to ameliorate experimental thyroiditis and diabetes in mice [51,52]. Although some studies have shown that GM-CSF could be co-produced by IL-17-producing T cells [33,34], our data

and other recent studies indicate that none, or only low levels, of GM-CSF are produced by Th17 cells [34,35,53].

Our findings also provide support for the proposition that arthritis could be driven/sustained by different inflammatory pathways in different patients resulting in a heterogeneous response to TNF α inhibitors. The lack of responsiveness to anti-TNF α in patients with low plasma levels of GM-CSF/low frequency GM-CSF⁺ T cells indicates that disease processes in these patients are unlikely to be dominated by a pathways driven by TNF α /GM-CSF. However, an alternative explanation could be that anti-TNF α ameliorate arthritis in responder patients by binding to autoreactive effector T cells that express membrane TNF α and thereby ameliorate their pro-inflammatory effects. In non-responder patients, in contrast, it is likely that IL-1 β and IL-23 produced by monocytes, when TNF α is blocked, drive Th17 cell differentiation and/or survival and promote chronic inflammation through a different pathway [12].

The current study has also revealed that monocytes from non-responder patients produce higher MCP-1 levels compared with responder patients. A significant difference was seen after 3 months of treatment, which may be accounted for by the possibility that monocytes in anti-TNF α non-responders are resistant to apoptosis [54]. However, further studies of monocytes in responder and non-responder RA patients could elucidate mechanisms that drive inflammation in patient subsets. Such studies may also reveal if there is a relationship between high IL-17 levels in non-responder patients and high level MCP-1 production by monocytes [55]. All in all, our study is consistent with other studies published in the current issue to indicate that assessment of immune cell subsets could identify biomarkers that help in designing evidence-based strategies for prescribing the most effective biological therapies in autoimmune diseases [56,57].

Conclusions

This study shows that different T cell subsets are likely involved in driving arthritis in different RA patients and that depending on which T cell subset predominates, determines which patient responds to anti-TNF α . In addition to its potential clinical relevance, the study provides insights into distinct inflammatory pathways that could drive the disease and help in future patient stratification for treatment. Further in-depth studies of GM-CSF⁺TNF α ⁺ T cells and IL-17-producing T cells, monocyte subsets, their proteome, transcriptome and genotype could provide invaluable new markers for targeted therapies in RA.

Acknowledgment: This study was supported by an Investigator Initiated Research Grant No. WS872934 from Pfizer.

Conflict of Interest: The authors declare no financial gains and no conflict of interest.

References

1. McInnes IB, Schett G (2011) The pathogenesis of rheumatoid arthritis. *N Engl J Med* 365(23):2205-2219.
2. Suzuki A, Kochi Y, Okada Y, Yamamoto K (2011) Insight from genome-wide association studies in rheumatoid arthritis and multiple sclerosis. *FEBS Lett* 585(23):3627-3632.
3. Elliott MJ, Maini RN, Feldmann M, Long-Fox A, Charles P, Katsikis P, Brennan FM, Walker J, Bijl H, Ghayeb J et al (1993) Treatment of rheumatoid arthritis with chimeric monoclonal antibodies to tumor necrosis factor alpha. *Arthritis Rheum* 36(12):1681-1690.
4. Emery P, Breedveld FC, Hall S, Durez P, Chang DJ, Robertson D, Singh A, Pedersen RD, Koenig AS, Freundlich B (2008) Comparison of methotrexate monotherapy with a combination of methotrexate and etanercept in active, early, moderate to severe rheumatoid arthritis (COMET): a randomised, double-blind, parallel treatment trial. *Lancet* 372(9636):375-382.
5. Klarenbeek NB, Allaart CF, Kerstens PJ, Huizinga TW, Dijkmans BA (2009) The BeSt story: on strategy trials in rheumatoid arthritis. *Curr Opin Rheumatol* 21(3):291-298.
6. Nie H, Zheng Y, Li R, Guo TB, He D, Fang L, Liu X, Xiao L, Chen X, Wan B et al (2013) Phosphorylation of FOXP3 controls regulatory T cell function and is inhibited by TNF-alpha in rheumatoid arthritis. *Nat Med* 19(3):322-328.
7. McGovern JL, Nguyen DX, Notley CA, Mauri C, Isenberg DA, Ehrenstein MR (2012) Th17 cells are restrained by Treg cells via the inhibition of interleukin-6 in patients with rheumatoid arthritis responding to anti-tumor necrosis factor antibody therapy. *Arthritis Rheum* 64(10):3129-3138.
8. Evans HG, Roostalu U, Walter GJ, Gullick NJ, Frederiksen KS, Roberts CA, Sumner J, Baeten DL, Gerwien JG, Cope AP et al (2014) TNF-alpha blockade induces IL-10 expression in human CD4+ T cells. *Nat Commun* 5:3199.
9. Cope AP, Liblau RS, Yang XD, Congia M, Laudanna C, Schreiber RD, Probert L, Kollias G, McDevitt HO (1997) Chronic tumor necrosis factor alters T cell responses by attenuating T cell receptor signaling. *J Exp Med* 185(9):1573-1584.
10. Cope AP (2008) T cells in rheumatoid arthritis. *Arthritis Res Ther* 10 Suppl 1:S1.
11. Chen DY, Chen YM, Chen HH, Hsieh CW, Lin CC, Lan JL (2011) Increasing levels of circulating Th17 cells and interleukin-17 in rheumatoid arthritis patients with an inadequate response to anti-TNF- α therapy. *Arthritis Res Ther* 13(4):R126.
12. Alzabin S, Abraham SM, Taher TE, Palfreeman A, Hull D, McNamee K, Jawad A, Pathan E, Kinderlerer A, Taylor PC et al (2012) Incomplete response of inflammatory arthritis to TNFalpha blockade is associated with the Th17 pathway. *Ann Rheum Dis* 71(10):1741-1748.
13. Kawashima M, Miossec P (2005) Effect of treatment of rheumatoid arthritis with infliximab on IFN gamma, IL4, T-bet, and GATA-3 expression: link with improvement of systemic inflammation and disease activity. *Ann Rheum Dis* 64(3):415-418.
14. Bystrom J, Clanchy FI, Taher TE, Mangat P, Jawad AS, Williams RO, Mageed RA (2016) TNF α in the regulation of Treg and Th17 cells in rheumatoid arthritis and other autoimmune inflammatory diseases. *Cytokine pii: S1043-4666(16)30490-2*.

15. Fabre S, Dupuy AM, Dossat N, Guisset C, Cohen JD, Cristol JP, Daures JP, Jorgensen C (2008) Protein biochip array technology for cytokine profiling predicts etanercept responsiveness in rheumatoid arthritis. *Clin Exp Immunol* 153(2):188-195.
16. Kayakabe K, Kuroiwa T, Sakurai N, Ikeuchi H, Kadiombo AT, Sakairi T, Kaneko Y, Maeshima A, Hironuma K, Nojima Y (2012) Interleukin-1 β measurement in stimulated whole blood cultures is useful to predict response to anti-TNF therapies in rheumatoid arthritis. *Rheumatology* 51(9):1639-1643.
17. Wampler Muskardin T, Vashisht P, Dorschner JM, Jensen MA, Chrabot BS, Kern M, Curtis JR, Danila MI, Cofield SS, Shadick N et al (2016) Increased pretreatment serum IFN- β / α ratio predicts non-response to tumour necrosis factor α inhibition in rheumatoid arthritis. *Ann Rheum Dis* 75(10):1757-1762.
18. Hueber W, Tomooka BH, Batliwalla F, Li W, Monach PA, Tibshirani RJ, Van Vollenhoven RF, Lampa J, Saito K, Tanaka Y et al (2009) Blood autoantibody and cytokine profiles predict response to anti-tumor necrosis factor therapy in rheumatoid arthritis. *Arthritis Res Ther* 11(3):R76.
19. Greisen SR, Schelde KK, Rasmussen TK, Kragstrup TW, Stengaard-Pedersen K, Hetland ML, Hørslev-Petersen K, Junker P, Østergaard M, Deleuran B et al (2014) CXCL13 predicts disease activity in early rheumatoid arthritis and could be an indicator of the therapeutic 'window of opportunity'. *Arthritis Res Ther* 16(5):434.
20. Han BK, Kuzin I, Gaughan JP, Olsen NJ, Bottaro A (2016) Baseline CXCL10 and CXCL13 levels are predictive biomarkers for tumor necrosis factor inhibitor therapy in patients with moderate to severe rheumatoid arthritis: a pilot, prospective study. *Arthritis Res Ther* 18:93.
21. Dennis G Jr, Holweg CT, Kummerfeld SK, Choy DF, Setiadi AF, Hackney JA, Haverty PM, Gilbert H, Lin WY, Diehl L et al (2014) Synovial phenotypes in rheumatoid arthritis correlate with response to biologic therapeutics. *Arthritis Res Ther* 16(2):R90.
22. Choi IY, Gerlag DM, Herenius MJ, Thurlings RM, Wijbrandts CA, Foell D, Vogl T, Roth J, Tak PP, Holzinger D (2015) MRP8/14 serum levels as a strong predictor of response to biological treatments in patients with rheumatoid arthritis. *Ann Rheum Dis* 74(3):499-505.
23. Lequerré T, Gauthier-Jauneau AC, Bansard C, Derambure C, Hiron M, Vittecoq O, Daveau M, Mejjad O, Daragon A, Tron F et al (2006) Gene profiling in white blood cells predicts infliximab responsiveness in rheumatoid arthritis. *Arthritis Res Ther* 8(4):R105.
24. Julià A, Erra A, Palacio C, Tomas C, Sans X, Barceló P, Marsal S (2009) An eight-gene blood expression profile predicts the response to infliximab in rheumatoid arthritis. *PLoS One* 4(10):e7556.
25. Folkersen L, Brynedal B, Diaz-Gallo LM, Ramsköld D, Shchetynsky K, Westerlind H, Sundström Y, Schepis D, Hensvold A, Vivar N et al (2016) Integration of known DNA, RNA and protein biomarkers provides prediction of anti-TNF response in rheumatoid arthritis: results from the COMBINE study. *Mol Med* 22(8):322-328.
26. Castro-Villegas C, Pérez-Sánchez C, Escudero A, Filipescu I, Verdu M, Ruiz-Limón P, Aguirre MA, Jiménez-Gomez Y, Font P, Rodríguez-Ariza A et al (2015) Circulating miRNAs as potential biomarkers of therapy effectiveness in rheumatoid arthritis patients treated with anti-TNF α . *Arthritis Res Ther* 17:49.
27. Krintel SB, Dehlendorff C, Hetland ML, Hørslev-Petersen K, Andersen KK, Junker P, Pødenphant J, Ellingsen T, Ahlquist P, Lindegaard HM et al (2016) Prediction of treatment response to adalimumab: a double-blind placebo-controlled study of

- circulating microRNA in patients with early rheumatoid arthritis. *Pharmacogenomics J* 16(2):141-146.
28. Fransen J, van Riel PL (2005) The Disease Activity Score and the EULAR response criteria. *Clin Exp Rheumatol* 23(5 Suppl 39):S93-99.
 29. Chara L, Sánchez-Atrio A, Pérez A, Cuende E, Albarrán F, Turrión A, Chevarria J, Sánchez MA, Monserrat J, de la Hera A et al (2012) Monocyte populations as markers of response to adalimumab plus MTX in rheumatoid arthritis. *Arthritis Res Ther* 14(4):R175.
 30. Church LD, Filer AD, Hidalgo E, Howlett KA, Thomas AM, Rapecki S, Scheel-Toellner D, Buckley CD, Raza K (2010) Rheumatoid synovial fluid interleukin-17-producing CD4 T cells have abundant tumor necrosis factor-alpha co-expression, but little interleukin-22 and interleukin-23R expression. *Arthritis Res Ther* 12(5):R184.
 31. van der Bruggen T, Nijenhuis S, van Raaij E, Verhoef J, van Asbeck BS (1999) Lipopolysaccharide-induced tumor necrosis factor alpha production by human monocytes involves the raf-1/MEK1-MEK2/ERK1-ERK2 pathway. *Infect Immun* 67(8):3824-3829.
 32. Chu C Q, Field M, Feldmann M, Maini R N (1991) Localization of tumor necrosis factor alpha in synovial tissues and at the cartilage-pannus junction in patients with rheumatoid arthritis. *Arthritis Rheum* 34(9):1125-1132.
 33. Kara EE, McKenzie DR, Bastow CR, Gregor CE, Fenix KA, Ogunniyi AD, Paton JC, Mack M, Pombal DR, Seillet C et al (2015) CCR2 defines in vivo development and homing of IL-23-driven GM-CSF-producing Th17 cells. *Nat Commun* 6:8644.
 34. Piper C, Pesenacker AM, Bending D, Thirugnanabalan B, Varsani H, Wedderburn LR, Nistala K (2014) T cell expression of granulocyte-macrophage colony-stimulating factor in juvenile arthritis is contingent upon Th17 plasticity. *Arthritis Rheum* 66(7):1955-1960.
 35. Noster R, Riedel R, Mashreghi MF, Radbruch H, Harms L, Haftmann C, Chang HD, Radbruch A, Zielinski CE (2014) IL-17 and GM-CSF expression are antagonistically regulated by human T helper cells. *Sci Transl Med* 6(241):241ra80.
 36. Reynolds G, Gibbon JR, Pratt AG, Wood MJ, Coady D, Raftery G, Lorenzi AR, Gray A, Filer A, Buckley CD et al (2016) Synovial CD4+ T-cell-derived GM-CSF supports the differentiation of an inflammatory dendritic cell population in rheumatoid arthritis. *Ann Rheum Dis* 75(5):899-907.
 37. Asseman C, Mauze S, Leach MW, Coffman RL, Powrie F (1999) An essential role for interleukin 10 in the function of regulatory T cells that inhibit intestinal inflammation. *J Exp Med* 190(7):995-1004.
 38. Maggi L, Santarlasci V, Capone M, Peired A, Frosali F, Crome SQ, Querci V, Fambrini M, Liotta F, Levings MK et al (2010) CD161 is a marker of all human IL-17-producing T-cell subsets and is induced by RORC. *Eur J Immunol* 40(8):2174-2181.
 39. Buch MH, Bingham SJ, Bryer D, Emery P (2007) Long-term infliximab treatment in rheumatoid arthritis: subsequent outcome of initial responders. *Rheumatology* 46(7):1153-1156.
 40. Jacob CO, McDevitt HO (1988) Tumour necrosis factor-alpha in murine autoimmune 'lupus' nephritis. *Nature* 331(6154):356-358.
 41. Satoh J, Seino H, Abo T, Tanaka S, Shintani S, Ohta S, Tamura K, Sawai T, Nobunaga T, Oteki T et al (1989) Recombinant human tumor necrosis factor alpha suppresses autoimmune diabetes in nonobese diabetic mice. *J Clin Invest* 84(4):1345-1348.

42. Liu J, Marino MW, Wong G, Grail D, Dunn A, Bettadapura J, Slavin AJ, Old L, Bernard CC (1998) TNF is a potent anti-inflammatory cytokine in autoimmune-mediated demyelination. *Nat Med* 4(1):78-83.
43. Jacob N, Yang H, Pricop L, Liu Y, Gao X, Zheng SG, Wang J, Gao HX, Putterman C, Koss MN et al (2009) Accelerated pathological and clinical nephritis in systemic lupus erythematosus-prone New Zealand Mixed 2328 mice doubly deficient in TNF receptor 1 and TNF receptor 2 via a Th17-associated pathway. *J Immunol* 182(4):2532-2541.
44. Notley CA, Inglis JJ, Alzabin S, McCann FE, McNamee KE, Williams RO (2008) Blockade of tumor necrosis factor in collagen-induced arthritis reveals a novel immunoregulatory pathway for Th1 and Th17 cells. *J Exp Med* 205(11):2491-2497.
45. Potter C, Hyrich KL, Tracey A, Lunt M, Plant D, Symmons DP, Thomson W, Worthington J, Emery P, Morgan AW et al (2009) Association of rheumatoid factor and anti-cyclic citrullinated peptide positivity, but not carriage of shared epitope or PTPN22 susceptibility variants, with anti-tumour necrosis factor response in rheumatoid arthritis. *Ann Rheum Dis* 68(1):69-74.
46. Hueber W, Tomooka BH, Batliwalla F, Li W, Monach PA, Tibshirani RJ, Van Vollenhoven RF, Lampa J, Saito K, Tanaka Y et al (2009) Blood autoantibody and cytokine profiles predict response to anti-tumor necrosis factor therapy in rheumatoid arthritis. *Arthritis Res Ther* 11(3):R76.
47. Cornish AL, Campbell IK, McKenzie BS, Chatfield S, Wicks IP (2009) G-CSF and GM-CSF as therapeutic targets in rheumatoid arthritis. *Nat Rev Rheumatol* 5(10):554-559.
48. Hamilton JA (2008) Colony-stimulating factors in inflammation and autoimmunity. *Nat Rev Immunol* 8(7):533-544.
49. de Vries E G, Willemse P H, Biesma B, Stern A C, Limburg P C, Vellenga E (1991) Flare-up of rheumatoid arthritis during GM-CSF treatment after chemotherapy. *Lancet* 338(8765):517-518.
50. Herndler-Brandstetter D, Flavell RA (2014) Producing GM-CSF: a unique T helper subset? *Cell research* 24(12):1379-1380.
51. Ganesh BB, Cheatem DM, Sheng JR, Vasu C, Prabhakar BS (2009) GM-CSF-induced CD11c+CD8a--dendritic cells facilitate Foxp3+ and IL-10+ regulatory T cell expansion resulting in suppression of autoimmune thyroiditis. *Int Immunol* 21(3):269-282.
52. Gaudreau S, Guindi C, Menard M, Besin G, Dupuis G, Amrani A (2007) Granulocyte-macrophage colony-stimulating factor prevents diabetes development in NOD mice by inducing tolerogenic dendritic cells that sustain the suppressive function of CD4+CD25+ regulatory T cells. *J Immunol* 179(6):3638-3647.
53. Sheng W, Yang F, Zhou Y, Yang H, Low P Y, Kemeny D M, Tan P, Moh A, Kaplan MH, Zhang Y et al (2014) STAT5 programs a distinct subset of GM-CSF-producing T helper cells that is essential for autoimmune neuroinflammation. *Cell Res* 24(12):1387-1402.
54. Meusch U, Klingner M, Baerwald C, Rossol M, Wagner U (2013) Deficient spontaneous in vitro apoptosis and increased tmTNF reverse signaling-induced apoptosis of monocytes predict suboptimal therapeutic response of rheumatoid arthritis to TNF inhibition. *Arthritis Res Ther* 15(6):R219.
55. Shahrara S, Pickens SR, Mandelin AM 2nd, Karpus WJ, Huang Q, Kolls JK, Pope RM (2010) IL-17-mediated monocyte migration occurs partially through CC chemokine ligand 2/monocyte chemoattractant protein-1 induction. *J Immunol* 184(8):4479-4487.

56. Gazeau P, Alegria GC, Devauchelle-Pensec V, Lemerle J, Bendaoud B, Saraux A, Cornec D, Renaudineau Y (2017) Switched memory B cells influence response to abatacept in rheumatoid arthritis. *Clinic Rev Allerg Immunol* (In press).
57. Schreiber K, Nocturne G, Cornec D, Daïen CI (2017) Lymphocytes as biomarkers of therapeutic response in rheumatic autoimmune diseases, a realistic goal? *Clinic Rev Allerg Immunol* (In press).

Table 1 Reported studies identifying predictive biomarkers of RA patients' responsiveness to TNF α inhibitors.

Biomarkers studied

Cytokines	Clinical sample	Protocol and outcome	Reference
12 Cytokines/Chemokines	Serum	Protein biochip array; high levels of MCP-1 and EGF associate with the response to etanercept	15
TNF α , IL-1 β , IL-6	White blood cells (WBCs)	LPS-stimulated WBCs; high levels of IL-1 β by predicted responsiveness to TNF α inhibitors	16
IFN β /IFN α	Serum	Functional reporter cell assay; increased ratio of IFN β to IFN α strongly associated with non-response to TNF α inhibitors	17
Cytokines+ Autoantibodies	Serum	Antigen array and multiplex cytokine assay; identified multi-parameter proteins that predicted response to etanercept	18
Chemokines/Inflammatory Mediators			
CXCL13	Plasma	ELISA; high baseline levels associated with remission at 2 years following treatment with TNF α inhibitors	19
CXCL10+CXCL13	Serum	ELISA; elevated baseline levels were associated with favourable response to TNF α inhibitors	20
ICAM1+CXCL13	Serum	Electrochemiluminescence; high soluble ICAM and low CXCL13 levels predict good response to anti-TNF α	21
MRP8/14	Serum	ELISA; high levels of the myeloid-related protein (MRP)8/14 complex (endogenous TLR-4 receptor agonist) predicted response to TNF α inhibitors	22

High GM-CSF levels predict responsiveness to anti-TNF

mRNAs

Panel of 8 transcripts	Blood mononuclear cells (BMNCs)	Microarray and quantitative RT-PCR; transcripts for ribosomal components, cell adhesion and inhibition of migration/invasion, cytochromes, proteasome-mediated proteolysis, enzymes and signalling predicted responsiveness to infliximab	23
Panel of 8 transcripts	Whole blood	Microarray; transcripts of genes encoding HLA-DRB3, SH2D1B, GNLY, CAMP, SLC2A3 and IL2RB, MXD4 and TLR5 predicted response to infliximab	24
Panel of 8 transcripts	Whole blood	High-throughput RNA sequencing, DNA genotyping and proteomics; transcripts for SORBS3, AKAP9, CYP4F12, MUSTN, CX3CR1, SLC2A3, C21orf58 and TBC1D8 two SNPs and plasma sICAM1/CXCL13 protein ratio predicted responsiveness to TNF α inhibitors	25
Micro RNA 23 (miR-23) and miR-223	Serum	miRNA polymerase chain reaction (PCR) array; miR-23 predicted response to anti-TNF α /DMARDs combination therapy	26
miR-22 and miR-886.3p	Serum	Micro RNA array; low expression of miR-22 and high expression of miR-886.3p associated with good response to adalimumab and methotrexate.	27

Table 2 Demographic and clinical data on patients included in the study.

	Total	Responders (%)	Non-responders (%)	P value
Number of patients	97	76 (78)	21 (22)	
Age (years)	55.7	55.8±13.9	57.0±14.1	0.49
Gender (female: male)	2.7	2.2	6	0.22
DAS-28 before treatment	5.76	5.70±0.8	5.96±0.6	0.12
DAS-28 after 3 months of treatment		3.5±1.2	5.8±0.7	<0.01

Statistical analyses were carried out using the Mann-Whitney U test except for gender ratios where Chi² test was used. Values provided as the Mean±Standard deviation (SD).

Table 3 Pro-inflammatory cytokine levels produced by *ex vivo* enriched T cells, B cells and monocytes from patients with RA prior to the start of treatment with anti-TNF α .

	T cells	P value	B cells	P value	Monocytes	P value
TNFα	45.2\pm 281.5		0.6\pm 1.4		1.0\pm 0.7	
Responders	62.3 \pm 341.6	0.2	0.8 \pm 1.8	0.78	1.0 \pm 0.8	0.39
Non-responders	6.6 \pm 5.0		0.2 \pm 0.3		1.0 \pm 0.5	
IL-1β	0.31\pm 1.5		0.1\pm0.3		3.9\pm 2.6	
Responders	0.4 \pm 1.8	0.7	0.2 \pm 0.3	0.51	4.0 \pm 3.0	0.83
Non-responders	0.1 \pm 0.1		0.04 \pm 0.1		3.9 \pm 2.0	
IL-6	1.7\pm4.6		1.7\pm3.6		23.7\pm16.8	
Responders	2.5 \pm 6.2	0.27	2.5 \pm 4.5	0.53	22.8 \pm 16.8	0.46
Non-responders	0.8 \pm 1.0		0.5 \pm 1.0		26.0 \pm 17.1	
GM-CSF	2.5\pm1.8		0.2\pm0.3		0.8\pm1.7	
Responders	2.9 \pm 1.9	0.02	0.2 \pm 0.3	0.55	0.6 \pm 1.8	0.06
Non-responders	1.6 \pm 1.1		0.2 \pm 0.4		1.1 \pm 1.4	
CCL2					20.1\pm23.0	
Responders					16.9 \pm 13.8	0.45
Non-responders					29.7 \pm 38.8	

Cytokine levels are presented as Mean \pm SD in ng/mL. For data on experiments summarised in the table cytokines produced by T and B cells and monocytes enriched from the blood of 46 responder and 21 non-responder patients are presented. Numbers in bold are for the Mean \pm SD of cytokines in all patients combined. *P* values <0.05 are considered statistically significant. Statistical analyses were carried out using the Mann-Whitney *U* test.

Figure legends:

Figure 1. Cytokine production by T cells, B cells and monocytes in rheumatoid arthritis

patients prior to and after treatment with anti-TNF α . T cells, B cells and monocytes were enriched by negative selection from the blood of RA patients immediately before treatment with anti-TNF α agents and then after 1 and 3 months. The cells were stimulated for 48 hrs and the level of cytokines produced determined using MSD multiplex kits. Levels of 16 cytokines were determined but only results of cytokines with notable differences are presented: (A) and (B) are for data on TNF α ; (C): IL-1 β and (D): GM-CSF. (A) TNF α produced by T cells stimulated with anti-CD3/anti-CD28 mAbs, B cells stimulated with anti-IgM/anti-CD40 and monocytes stimulated with LPS. (B) Mean \pm standard error of the mean (SEM) for TNF α produced by T cells and monocytes from 3 healthy controls and stimulated for 48hrs with PMA and Ionomycin. (C) and (D) show levels of IL-1 β and GM-CSF produced by T and B cells and monocytes activated and cultured as described in (A). (E) Data on GM-CSF production by T cells as described for (A) before treatment (0 months) and after 1 and then 3 months. (F) Data on the frequency of GM-CSF⁺ T cells before treatment and after 1 and 3 months in 11 responder and 3 non-responder RA patients. Enriched T cells were stimulated with PMA/Ionomycin for 16 hrs and stained for intracellular GM-CSF. (G) IL-6 production by T cells and (H) CCL2 production by monocytes. The data before and after treatment are for 67 patients. Differences between responders and non-responders were assessed using two-tailed Mann Whitney U test. Difference at different time points in the same group was assessed using Wilcoxon matched-pairs signed rank test. * indicates $P < 0.05$; ** indicates $P < 0.01$.

Figure 2. Cluster and Receiver Operating Characteristic (ROC) analyses for predictive

value of cytokines in the response to anti-TNF α . (A, B) Cluster analyses of TNF α , IL-1 β and GM-CSF in RA patients pre-treatment with anti-TNF α in relation to responsiveness/non-

responsiveness. Cluster analyses of GM-CSF, IL-1 β and TNF α levels (**A**) by *in vitro* activated T cells from responder (RES) and non-responder (N_R) patients whose identity codes are indicated by numbers and (**B**) in plasma from the same patients. The cluster analyses were performed using Pearson's correlation with complete linkage. The two clusters in (**A**) were highlighted with a blue (cluster 1) and a red (cluster 2) bracket. (**C**) Summary of data from the cluster in (**A**). Significant differences between responders and non-responders were calculated using Fishers exact test. (**D**) ROC curve analyses of the response to treatment with anti-TNF α . The increasing area under the ROC curve (area under the curve, AUC) corresponds to a higher diagnostic test yield. (**E**) Summary of sensitivity, specificity and odds ratio (OR) for GM-CSF levels by T cells and in plasma before treatment in predicting the response to treatment with anti-TNF α . PPV: positive predictive value; NPV: negative predictive value. Data are presented with 95% CI.


Figure 3. Levels of cytokines in plasma pre-treatment of RA patients with anti-TNF α .




Plasma levels of cytokines were determined immediately before treatment with anti-TNF α using MSD multiplex kits. Levels of cytokines shown are for: (**A**) TNF α ; (**B**) IL-1 β ; (**C**) IL-6; (**D**) GM-CSF and (**E**) MCP-1. Values are presented as the Mean \pm SEM in the plasma of 40 responder and 13 non-responder patients on whom plasma samples were available. Statistical analyses were with the two-tailed Mann Whitney U test. * indicates $P<0.05$; ** indicates $P<0.01$.

Figure 4. GM-CSF⁺ and IL-17⁺ T cells segregate into two different subsets.

Enriched T cells were stimulated with PMA/Ionomycin for 16 hrs and stained for membrane proteins and intracellular cytokines as indicated. (**A**) Contour plot showing the % of CCR6⁺ and CCR6⁺IL-17⁺ in blood T cells and (**B**) TNF α ⁺ and TNF⁺GM-CSF⁺ T cells in the blood of RA patients. CCR6⁺IL-17⁺ in (**A**) were gated and the frequency of T cells expressing intracellular

GM-CSF, TNF α and IFN γ ⁺ and membrane CD161 determined by FACS. **(B)** PMA/Ionomycin-stimulated T cells were stained for intracellular TNF α and GM-CSF and the frequency in responder and non-responder patients determined. The frequency of TNF α ⁺GM-CSF⁺ T cells that co-expressed IFN γ and IL-17 membrane CD161 was determined by multicolour FACS. The data represent the Mean \pm SEM values from 7 responder and 3 non-responder patients. **(C)** Contour plots showing characteristics of RA patients GM-CSF⁺ T cells (expression of CD45RA, CD45RO and CD161) pre-treatment. Percentages of differentially stained cell subsets are given inside the quadrants. Blood samples from 20 RA patients were analysed for these characterisation experiments. **(D)** Linear regression showing the correlation between the level of IL-17 and GM-CSF produced by patients' T cells activated with anti-CD3/anti-CD28 mAbs.





(E)

GM-CSF measured from

	T cells (per 200 cells) (Cut off >2.5pg/ mL)	Plasma (Cut off >3.5pg/mL)
OR	3.9	11.7
PPV	84.0%	87.5%
NPV	42.3%	62.5%
Sensitivity	58.3%	82.4%
Specificity	73.3%	71.4%

