

The fuzzy S^2 structure of M2-M5 systems in ABJM membrane theories

Horatiu Nastase^{a,1}, Constantinos Papageorgakis^{b,2} and Sanjaye Ramgoolam^{c,3}

^a *Global Edge Institute, Tokyo institute of Technology,
Tokyo 152-8550, Japan*

^b *Department of Theoretical Physics, Tata Institute of Fundamental Research,
Homi Bhabha Road, Mumbai 400 005, India*

^c *Centre for Research in String Theory, Department of Physics,
Queen Mary, University of London,
Mile End Road, London E1 4NS, UK*

ABSTRACT

We analyse the fluctuations of the ground-state/funnel solutions proposed to describe M2-M5 systems in the level- k mass-deformed/pure Chern-Simons-matter ABJM theory of multiple membranes. We show that in the large N limit the fluctuations approach the space of functions on the 2-sphere rather than the naively expected 3-sphere. This is a novel realisation of the fuzzy 2-sphere in the context of Matrix Theories, which uses bifundamental instead of adjoint scalars. Starting from the multiple M2-brane action, a U(1) Yang-Mills theory on $\mathbb{R}^{2,1} \times S^2$ is recovered at large N , which is consistent with a single D4-brane interpretation in Type IIA string theory. This is as expected at large k , where the semiclassical analysis is valid. Several aspects of the fluctuation analysis, the ground-state/funnel solutions and the mass-deformed/pure ABJM equations can be understood in terms of a discrete noncommutative realisation of the Hopf fibration. We discuss the implications for the possibility of finding an M2-brane worldvolume derivation of the classical S^3 geometry of the M2-M5 system. Using a rewriting of the equations of the SO(4)-covariant fuzzy 3-sphere construction, we also directly compare this fuzzy 3-sphere against the ABJM ground-state/funnel solutions and show them to be different.

¹E-mail address: nastase.h.aa@m.titech.ac.jp

²E-mail address: costis@theory.tifr.res.in

³E-mail address: s.ramgoolam@qmul.ac.uk

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 2 |
| 2 | The SO(4)-covariant fuzzy three-sphere in BLG and ABJM theories | 4 |
| 2.1 | The SO(4)-covariant (Guralnik-Ramgoolam) fuzzy S^3 | 5 |
| 2.2 | Bagger-Lambert-Gustavsson theory | 7 |
| 2.3 | ABJM theory | 9 |
| 3 | Fuzzy two-sphere structure of M2-M5 ABJM solutions | 10 |
| 3.1 | Ground-state matrices and symmetries | 11 |
| 3.2 | Fuzzy S^2 harmonics from $U(N) \times U(\bar{N})$ with bifundamentals | 13 |
| 3.3 | SU(2) harmonic decomposition of bifundamental matrices | 17 |
| 4 | Action for matter fluctuations on the two-sphere | 19 |
| 4.1 | Matrix fluctuation expansion | 20 |
| 4.2 | Fluctuations and bosonic field theory on the classical S^2 | 25 |
| 4.3 | The fuzzy funnel case | 32 |
| 5 | The CS-Higgs system and the Higgs mechanism | 32 |
| 5.1 | Higgs mechanism | 34 |
| 5.2 | Other contributions from the scalar kinetic term | 36 |
| 6 | Final action for fluctuations | 37 |
| 7 | M5 to D4 through Hopf | 38 |
| 7.1 | Physical radius and D4-brane energy for the funnel | 39 |
| 7.2 | \mathbb{Z}_k reduction of S^3 | 40 |
| 7.3 | Hopf fibration and the classical limit of the fuzzy two-sphere | 41 |
| 7.4 | Hindsight is 20/20 : ABJM structure from fuzzy Hopf | 42 |
| 7.5 | Multi-membrane actions and S^3 at large N | 44 |
| 8 | Discussion and future avenues | 45 |
| 9 | Summary and conclusions | 47 |
| A | Some useful formulae for the parallel scalars | 48 |
| A.1 | Identities for the $s_i s_i$ and r - s fluctuations | 48 |
| A.2 | Expressions and identities for s_i decomposition | 50 |
| B | Technical aspects of Higgsing | 51 |
| B.1 | Derivation of Eq.(5.12) | 51 |
| B.2 | Higgsing from trace over \mathbf{V}^- | 51 |
| C | Towards multiple M5-branes | 52 |

1 Introduction

M2 worldvolume theories have been at the centre of intense recent activity after the discovery of manifestly $\mathcal{N} = 8$ superconformal three dimensional Lagrangians by Bagger-Lambert and Gustavsson (BLG) [1–4], following earlier insights of [5, 6]. Inspired by subsequent related developments [7–11] and independent advances in supersymmetric Chern-Simons-matter theories [12, 13], ABJM [14] have proposed an $\mathcal{N} = 6$ Chern-Simons-matter theory with $U(N) \times U(N)$ gauge group, $SO(6)$ R-symmetry and equal but opposite Chern-Simons (CS) levels $(k, -k)$, to capture the dynamics of the low-energy limit of multiple M2-branes on an M-theory orbifold, $\mathbb{C}^4/\mathbb{Z}_k$. At $k = 1$ the theory is strongly coupled and is conjectured to describe membranes in flat space. The supersymmetry and R-symmetry were argued to be nonperturbatively enhanced to $\mathcal{N} = 8$ and $SO(8)$ respectively.

An important test for any candidate theory of multiple membranes in flat space is that it should reproduce the physics of $M2 \perp M5$ intersections. These are M2-spike solutions of the M5-brane worldvolume theory [15], generalising the D1-spike solutions on D3 worldvolumes. In the case of $D1 \perp D3$ systems, there is a detailed understanding of the physics, both from the D1 and the D3-worldvolume perspectives [16].¹ They can be captured equivalently by a $U(N)$ theory of multiple D1-branes and a $U(1)$ theory of a single D3-brane with N units of magnetic flux. From the point of view of the $U(N)$ theory the $D1 \perp D3$ system is described in terms of a solution involving fuzzy 2-spheres [19–21] through the Myers effect [22]. These are related to families of matrices obeying

$$[X^i, X^j] = 2i\epsilon^{ijk} X^k . \tag{1.1}$$

The X^i enter the physics as ansätze for solving the equations of motion and their commutator action on the space of all $N \times N$ matrices organises these matrices into representations of $SU(2) \simeq SO(3)$. An important aspect of the geometry of the fuzzy 2-sphere involves the construction of fuzzy (matrix) spherical harmonics in $SU(2)$ representations, which approach the space of all classical S^2 spherical harmonics in the limit of large matrices [20]. This construction of fuzzy spherical harmonics allows the analysis of fluctuations in a nonabelian theory of D_p -branes to be expressed at large N in terms of an abelian higher dimensional theory of a $D(p+2)$ brane. At finite N the higher dimensional worldvolume theory becomes a noncommutative $U(1)$ with a UV cutoff [23–26].

The relation (1.1) also appears as an F-flatness condition for a particular mass deformation of $\mathcal{N} = 4$ SYM, called $\mathcal{N} = 1^*$, where all three adjoint chiral multiplets have acquired equal mass. This theory has a discrete set of vacua that satisfy the above condition, which from the string theory point of view can be interpreted as N D3 branes blowing up into concentric D5 branes with flux that adds up to N [27]. The matrix structure of these solutions is the same as for the $D1 \perp D3$ spikes, with the relative transverse directions between the D3 and the D5 forming a fuzzy S^2 at finite N , which can again be seen by analysing small fluctuations around the vacua [28–30]. The lift of this system to M-theory has been discussed in [31]. The natural generalisation of the D3-D5 system is M2-M5, and a theory of multiple membranes should include such solutions.

With the advent of the ABJM action it is possible to investigate both these kinds of M2-M5 configurations in greater detail. An interesting class of mass deformations of the ABJM theory were given in [32, 33]. An important difference between the latter theories and $\mathcal{N} = 1^*$ is that they preserve

¹Similar studies for the $D1 \perp D5$ and $D1 \perp D7$ systems have also been undertaken in [17, 18].

the same $\mathcal{N} = 6$ amount of supersymmetry as the parent theory but still break R-symmetry, down to $SU(2) \times SU(2) \times U(1)$, as well as conformal invariance. The authors of [33] also presented a set of ground-state solutions. Closely related solutions describe M2 \perp M5 funnels in the undeformed ABJM theory [33] and have subsequently been discussed in [34–38]. The explicit form of the matrices for these solutions was interpreted as exhibiting a fuzzy S^3 structure. Our aim here will be to revisit this claim by examining the algebra obeyed by the matrices. Hence, the study of these solutions and their associated fuzzy sphere geometry will be at the central focus of this paper.

Our main result will be that the above ground-state/funnel solutions actually describe fuzzy S^2 's, as opposed to fuzzy S^3 's, albeit realised in a new fashion involving the noncommutative base of the $S^1 \hookrightarrow S^3 \xrightarrow{\pi} S^2$ Hopf fibration. We verify this by analysing small fluctuations around the ground-states at large N, k and showing that they can be organised in terms of a $U(1)$ theory on $\mathbb{R}^{2,1} \times S^2$ consistent with an interpretation as a D4-brane in Type IIA. The calculation involves a combination of techniques familiar from the Matrix Theory literature and the novel Higgs mechanism [7] that relates $U(N) \times U(N)$ CS-matter theories with bifundamental matter to $U(N)$ YM theories with adjoint matter.

It is interesting that the equations of motion defining the ground-state solutions of [33] are the same as the BPS equations for the fuzzy funnel of the ABJM model (once the dependence on the worldvolume coordinate along the funnel is factored out). These equations are in turn a generalisation of the BPS equations of [6] that inspired Bagger and Lambert to define their theory. Our result will imply that the special case of the Bagger-Lambert \mathcal{A}_4 -theory is as of yet unique in describing a fuzzy S^3 ; in all other cases of BLG/ABJM-type three dimensional Lagrangians one is dealing with fuzzy S^2 solutions, despite the original motivation.

A natural definition of the fuzzy S^3 is that it should involve an $SO(4)$ -covariant construction using matrices or some other appropriate algebraic structure, generalising the fuzzy S^2 matrix construction. Accordingly, a fully $SO(4)$ -covariant matrix construction of fuzzy 3-spheres should exhibit matrix spherical harmonics transforming under $SO(4)$, which approach the classical spherical harmonics on S^3 in a large N limit. Such a construction was given by Guralnik and one of the present authors in [39–41]. The Guralnik-Ramgoolam (GR) construction has its mysteries, in that the matrix algebra contains the spherical harmonics only as a subset of all the matrices. A complete construction of a finite N geometry of fuzzy 3-spheres has to take account of the projection which does away with the extra degrees of freedom. This leads to non-associativity [40]. Related developments on fuzzy odd spheres appear in [42–51].

In Section 2, we begin by writing the equations obeyed by the generating matrices of the GR fuzzy S^3 in a new form, appropriate for comparison with the equations of motion of ABJM and BLG. We find that the GR fuzzy S^3 is a solution for the $N = 2$ case of the ABJM theory. This $U(2) \times U(2)$ theory is closely related to the \mathcal{A}_4 -theory of BLG, which is the same as the $SU(2) \times SU(2)$ ABJM theory. However, with our rewriting of the equations obeyed by the matrices of the GR fuzzy S^3 , we will explicitly see that these become different from the solutions of ABJM that were found in [33] for $N > 2$. Given that the mass-deformed theory of [33] does not contain an $SO(4)$ R-symmetry factor the absence of the ‘usual’ fuzzy S^3 should perhaps not come as a surprise.² The fact that the

²We note that for the extreme strongly-coupled region $k = 1, 2$ maximal supersymmetry and $SO(8)$ R-symmetry are expected to be restored. In particular, the $k = 1$ case corresponds to M2-branes in flat space. Since this region is well

GR construction does work for $N = 2$ is closely related to its role in the work of Basu and Harvey [6] and the BLG interpretation of the Basu-Harvey equation in terms of the of the \mathcal{A}_4 3-algebra. Developments on fuzzy M-brane intersections following [6] are reviewed in [53, 54].

In Section 3, we return to the ground-state solutions of ABJM theory at general N given in [33] in terms of matrices $G^\alpha, G_\alpha^\dagger$, and we make the fuzzy S^2 structure apparent. Since the ABJM theory has $U(N) \times U(N)$ gauge symmetry, the Lie algebra is a direct sum which can be viewed as matrices acting on a direct sum of vector spaces $\mathbf{V}^+ \oplus \mathbf{V}^-$. The ABJM action has a global $SU(4)$ symmetry, which is broken by the mass deformation to $SU(2) \times SU(2) \times U(1)$ and the complex scalars decompose as $C^I = (R^\alpha, Q^{\dot{\alpha}})$. One of the $SU(2)$'s acts on the coordinates R^α which are nonzero on the solutions. The other $SU(2)$ acts on the $Q^{\dot{\alpha}}$'s, which are zero on the solutions. The first $SU(2)$ is relevant to the geometry described by the solution and we will call it the geometric $SU(2)$. The generators of this $SU(2)$ can be constructed from bilinears in $G^\alpha, G_\alpha^\dagger$. We describe how the N^2 -dimensional space of matrices acting on \mathbf{V}^+ , called $End(\mathbf{V}^+)$, can be decomposed in terms of representations of this $SU(2)$. Likewise we give the decomposition of $End(\mathbf{V}^-)$. Interestingly these turn out to be different. We complete the $SU(2)$ decompositions with those of $Hom(\mathbf{V}^+, \mathbf{V}^-)$ and $Hom(\mathbf{V}^-, \mathbf{V}^+)$, that is matrices mapping $\mathbf{V}^+ \rightarrow \mathbf{V}^-$ and $\mathbf{V}^- \rightarrow \mathbf{V}^+$ respectively. These matrices are needed to describe the fluctuations of the matter fields which transform in the bifundamental of the gauge group. We obtain that the fuzzy spherical harmonics for all physical fluctuations are consistent with a fuzzy S^2 , and reduce to the usual spherical harmonics, with no other factors, in the ‘classical’ (large N) limit.

In Section 4, we calculate the action for the matter fluctuations to quadratic order around the fuzzy solution, and take their classical limit. In Section 5, we then analyse the gauge fields, for which we use the Higgs mechanism of [7] to turn the nonpropagating 3d CS-type action for $A^{(1)}$ and $A^{(2)}$ into the propagating 5d YM action. The full action for the fluctuations takes the form of an abelian YM theory on $\mathbb{R}^{2,1} \times S^2$ and we explain this result in terms of a D4 wrapping the S^2 in Type IIA string theory. In Section 7 we show how the IIA D4-brane interpretation is expected by considering the M2-M5 system in the large k limit of the M-theory quotient of [14]. The action of the quotient on the S^1 Hopf fibre of the S^3 cross-section of the M2-M5 system leads to a IIA reduction. The structure of the ground-state solutions and their fluctuations are retrospectively determined by considering possible finite matrix realisations of the fuzzy S^2 base of the $S^1 \hookrightarrow S^3 \xrightarrow{\pi} S^2$ Hopf fibration. In this discussion it becomes apparent that the S^1 fibre has (in a sense that we explain) been reduced to two points. The appendices contain some formulae and details of the calculations in Section 4.

Related work pertaining to M5-branes in the more general context of 3-algebra theories includes [55–63].

2 The $SO(4)$ -covariant fuzzy three-sphere in BLG and ABJM theories

In this section, we will rewrite the definition of the GR fuzzy S^3 in a way that can be embedded in BLG/ABJM theory and then see that it only matches the BPS fuzzy sphere solution of BLG and

beyond our semiclassical, large- k analysis, one cannot rule out the emergence of the S^3 when the full quantum theory is taken into account. An example of an emergent S^5 from quantum physics can be found in [52].

ABJM theory in the $r = 1$ case of the Γ -matrix representation.

Notation: We will denote by $M, N, \dots = 1, 8$ the 8 real indices of coordinates transverse to an M2 brane X^I , by $m, n, \dots = 1, 4$ the indices of 4 real coordinates X_m that will form a fuzzy S^3 , with $X_m X^m = 1$, by a, b, \dots the indices of generators of algebras T_a , for both regular Lie algebras and 3-algebras, by $I, J, \dots = 1, 4$ the indices of 4 complex coordinates C^I transverse to an M2, splitting into $\alpha, \beta, \dots = 1, 2$ and $\dot{\alpha}, \dot{\beta}, \dots = 1, 2$, by $A, B, \dots = 1, 6$ the SO(6) indices of $\mathcal{N} = 6$ supersymmetry ϵ^A and by $i, j, \dots = 1, 3$ the indices defining an S^2 , with $X_i X^i = 1$. Also, for SU(N) matrices that can be written as tensor products of Γ -matrix representations, we will denote by r the number of such representations.

2.1 The SO(4)-covariant (Guralnik-Ramgoolam) fuzzy S^3

The SO(4)-covariant matrix construction of the fuzzy S^3 , in terms of tensor products of Γ -matrices, was done by Guralnik and Ramgoolam (GR) in [39–41]. One splits the vector space of spinor representations of SO(4) under $\text{SO}(4) \simeq \text{SU}(2) \times \text{SU}(2)$ as $V = V_+ \oplus V_-$. Take the subspace \mathcal{R}_r of $\text{Sym}(V^{\otimes r})$, acting as the basis vector space for the fuzzy three-sphere representation. \mathcal{R}_r is defined as $\mathcal{R}_r^+ \oplus \mathcal{R}_r^-$, where \mathcal{R}_r^+ and \mathcal{R}_r^- are spaces of $\text{SU}(2) \times \text{SU}(2)$ with labels $(\frac{r+1}{2}, \frac{r-1}{2})$ and $(\frac{r-1}{2}, \frac{r+1}{2})$ respectively, denoting the number of V_+ and V_- factors in each. Then the fuzzy S^3 coordinates X_m (SU(N) matrices) are given by

$$X_m = \mathcal{P}_{\mathcal{R}_r} \hat{X}_m \mathcal{P}_{\mathcal{R}_r} = \mathcal{P}_{\mathcal{R}_r^+} \hat{X}_m \mathcal{P}_{\mathcal{R}_r^-} + \mathcal{P}_{\mathcal{R}_r^-} \hat{X}_m \mathcal{P}_{\mathcal{R}_r^+} = X_m^+ + X_m^-, \quad (2.1)$$

where $\mathcal{P}_{\mathcal{R}_r} = \mathcal{P}_{\mathcal{R}_r^+} + \mathcal{P}_{\mathcal{R}_r^-}$, with $\mathcal{P}_{\mathcal{R}_r^+}, \mathcal{P}_{\mathcal{R}_r^-}$ being projectors onto the subspaces \mathcal{R}_r^+ and \mathcal{R}_r^- respectively and

$$\hat{X}_m = \sum_l \rho_l(\Gamma_m), \quad (2.2)$$

with ρ_l denoting which factor of $\text{Sym}(V^{\otimes r})$ the Γ -matrix acts on. Thus the coordinates of the fuzzy S^3 are represented by the matrices $X_m = X_m^+ + X_m^-$, where

$$X_m^+ = \mathcal{P}_{\mathcal{R}_r^+} \sum_l \rho_l(\Gamma_m P_+) \mathcal{P}_{\mathcal{R}_r^+} \quad \text{and} \quad X_m^- = \mathcal{P}_{\mathcal{R}_r^-} \sum_l \rho_l(\Gamma_m P_-) \mathcal{P}_{\mathcal{R}_r^-}. \quad (2.3)$$

These turn out not to be sufficient to describe the fuzzy S^3 ; one also needs to consider the matrices $Y_m = X_m^+ - X_m^-$. Equivalently, one can instead use the coordinates X_m^+ and X_m^- . Commutation relations for these objects were found in [41] and were simplified in [46]. These can be understood to act as equations of motion for the matrix fields, or alternatively as a defining algebra. The simplified set found in [46] is

$$[J_{mn}, X_n] = 6X_m, \quad [J_{mn}, Y_n] = 6Y_m, \quad (2.4)$$

where X_m are the sphere coordinates, satisfying $X_m X_m = N \mathbb{1}$, while

$$J_{mn} = \alpha[X_m, X_n] + \beta \epsilon_{mnpq} \{X_p, Y_q\}, \quad (2.5)$$

with

$$\alpha = -\frac{2}{(r+1)(r+3)} \quad \text{and} \quad \beta = \frac{r+2}{(r+1)(r+3)} \quad (2.6)$$

and also the constraint

$$[X_m, X_n] = -[Y_m, Y_n]. \quad (2.7)$$

We would now like to write this definition of the fuzzy S^3 in a form that can be straightforwardly compared with the BLG/ABJM theories.

In [50] it was proved that the fuzzy S^3 also satisfies the Matrix model type equation

$$[[X_m, X_n], X_n] = \left((r+1)(r+3) + 4 \right) X_m, \quad (2.8)$$

which means that the X_m equations in (2.4) become

$$\epsilon^{mnpq}[\{X_p, Y_q\}, X_n] = 8 \left(\frac{(r+1)(r+3) + 1}{r+2} \right) X_m. \quad (2.9)$$

The second equation in (2.4) gives the same result with X_m and Y_m interchanged. However, since $X_m = X_m^+ + X_m^-$ and $Y_m = X_m^+ - X_m^-$ while $X_m^+ X_n^+ = X_m^- X_n^- = 0$, we have the more restrictive constraints (as compared to (2.7))

$$\begin{aligned} X_m X_n &= -Y_m Y_n \\ X_m Y_n &= -Y_m X_n. \end{aligned} \quad (2.10)$$

This in turn means that (2.9) becomes

$$\epsilon^{mnpq} X_n Y_p X_q = 2 \left(\frac{(r+1)(r+3) + 1}{r+2} \right) X_m \quad (2.11)$$

and there also exists a corresponding relation with X_m interchanged with Y_m . Then (2.11) and the constraints (2.10) can be taken as a definition of the fuzzy S^3 , where

$$X_m X_m = X_m^+ X_m^- + X_m^- X_m^+ = \frac{(r+1)(r+3)}{2} \equiv N \quad (2.12)$$

in a similar way to the fuzzy S^2 , for which the defining relation is $X_i X_i = R^2$ together with the $SU(2)$ algebra

$$[X_i, X_j] = 2i\epsilon_{ijk} X_k. \quad (2.13)$$

Another way to define the fuzzy S^3 is by writing the equations in terms of variables going from the + space to the - space or vice-versa, *i.e.* X_m^+ and X_m^- . We obtain

$$\begin{aligned} \epsilon^{mnpq} X_n^+ X_p^- X_q^+ &= 2 \left(\frac{(r+1)(r+3) + 1}{r+2} \right) X_m^+ \\ \epsilon^{mnpq} X_n^- X_p^+ X_q^- &= 2 \left(\frac{(r+1)(r+3) + 1}{r+2} \right) X_m^-, \end{aligned} \quad (2.14)$$

which are to be supplemented with the sphere condition (2.12) and the constraints

$$X_m^+ X_n^+ = X_m^- X_n^- = 0, \quad (2.15)$$

since in that case the constraints (2.10) are automatically satisfied.

The relations (2.14) together with the sphere condition (2.12) and the constraints (2.15) give a new definition of the GR fuzzy S^3 that can be easily matched against BLG/ABJM.

2.2 Bagger-Lambert-Gustavsson theory

The fuzzy S^3 solution that was proposed in the model found by Bagger-Lambert [1] and independently Gustavsson [4] is similar to the one defined above and we will see that the two match for the case of the $SO(4)$, or \mathcal{A}_4 -algebra. However, since \mathcal{A}_4 is the only example of a finite dimensional 3-algebra theory with positive definite metric [64, 65], this is somewhat disappointing as we would like to find the S^3 as a solution to the theory for any number N of M2-branes. In retrospect, one needs to look instead at the ABJM theory, but we will see that there one also encounters problems in identifying the GR fuzzy S^3 .

In [6] Basu and Harvey constructed a fuzzy S^3 funnel for the X^m coordinates of M5-branes, as obtained from the hypothetical worldvolume theory of multiple M2-branes, with BPS equation

$$\frac{dX^m}{ds} + \frac{k}{4!} \epsilon^{mnpq} [G, X^n, X^p, X^q] = 0. \quad (2.16)$$

This is solved by the ansatz

$$X^m(s) \propto \frac{1}{\sqrt{s}} G^m \quad (2.17)$$

and the corresponding fuzzy 3-sphere relation

$$G^m \sim \epsilon^{mnpq} [G, G^n, G^p, G^q], \quad (2.18)$$

where G is a fixed matrix and s is the direction along which the membranes extend away from the fivebrane. Subsequently, Bagger and Lambert [1, 3] proposed the fuzzy funnel (BPS) equation

$$\frac{dX^m}{ds} = ik\epsilon^{mnpq} [X^n, X^p, X^q]. \quad (2.19)$$

Here m, n, p, q are directions transverse to the M2s and defining the fuzzy 3-funnel parallel to the M5 by $X_m X_m = 1$. The full set of M2 transverse scalars is decomposed as $X^M = X_a^M T^a$ ($M = 1, \dots, 8$ are all the coordinates transverse to M2, including the 4 coordinates m, n, p, q), where the T^a are generators that define a 3-algebra [2]

$$[T^a, T^b, T^c] = f^{abc} T^d \quad (2.20)$$

and the metric $h^{ab} = \text{Tr}(T^a T^b)$, used to raise and lower indices, is assumed to be positive definite.³ Substituting the 3-algebra (2.20), we obtain the BPS equation for the commuting fields X_a^m

$$\partial_s X_a^m = ik\epsilon^{mnpq} f^{bcd} X_b^n X_c^p X_d^q. \quad (2.21)$$

The corresponding fuzzy S^3 equation, which is an extension of the Basu-Harvey equation (2.18), is

$$R^2 X^m = ik\epsilon^{mnpq} [X^n, X^p, X^q]. \quad (2.22)$$

³If the metric is Lorentzian [56, 66, 67] one has to deal with potential ghosts in the quantum theory that could lead to violation of unitarity. These have been shown to decouple by gauging a certain shift symmetry but the theory is then on-shell equivalent to usual $\mathcal{N} = 8$ SYM in 3d [7, 68–70]. There are various interpretations of the ‘un-gauged’ version [71, 72], which has been shown to be obtained from ABJM through a particular scaling if one also adds a decoupled abelian ‘ghost’ multiplet to the latter [73–76].

It would be natural to think of this equation, with X_m satisfying $X_m X_m = R^2$, as the analogue of (2.11). However, note that the spacetime index m of the commuting fields X_a^m in (2.21) is contracted with ϵ^{mnpq} , while the ‘noncommuting’ (*i.e.* matrix or algebra) index a is contracted with f^{abc}_d , which makes it *a priori* different from (2.11) for general f^{abc}_d . Only in the case of the \mathcal{A}_4 (or $\text{SO}(4)$) algebra, when

$$f^{abcd} = f^{abc}_e h^{de} = \tilde{f} \epsilon^{abcd} \quad (2.23)$$

do we have a similar equation, and in fact that is the only case for which the Bagger-Lambert equation is equivalent to the GR fuzzy 3-sphere.

We can see this equivalence explicitly by using van Raamsdonk’s reformulation [9] of the \mathcal{A}_4 -theory in terms of a usual $\text{SO}(4) \simeq \text{SU}(2) \times \text{SU}(2)$ CS gauge theory with bifundamental matter fields. The reformulation starts with the observation that the objects $\tau^a = (i\sigma_j, \mathbb{1})$ obey

$$\tau^{[a} \tau^{\dagger b} \tau^c] = -\epsilon^{abcd} \tau^d = \epsilon^{dabc} \tau^d. \quad (2.24)$$

This nontrivial relation is obtained because $\tau^a = (i\sigma_j, \mathbb{1})$ have different properties under conjugation: $(i\sigma_j)^\dagger = -i\sigma_j$, whereas $\mathbb{1}^\dagger = \mathbb{1}$. Then define

$$X^M = \frac{1}{2} X_a^M \tau^a = \frac{1}{2} (X_4^M \mathbb{1} + i X_i^M \sigma^i), \quad (2.25)$$

obeying the reality condition

$$(X^M)^* = -\epsilon X^M \epsilon, \quad (2.26)$$

where $\epsilon = -i\sigma_2$ is the antisymmetric 2d symbol. These fields are now in a representation of $\text{SO}(4) \simeq \text{SU}(2)_1 \times \text{SU}(2)_2$, as bifundamental fields with one index in $\text{SU}(2)_1$ and another one in $\text{SU}(2)_2$, *i.e.* $(X^I)^\alpha_{\dot{\alpha}}$, with $\alpha \in \text{SU}(2)_1$ and $\dot{\alpha} \in \text{SU}(2)_2$. Then (2.21) becomes

$$\partial_s X^m = -ik \tilde{f} \epsilon^{mnpq} X^n X^{\dagger p} X^q. \quad (2.27)$$

The corresponding fuzzy S^3 equation

$$R^2 X^m = -ik \epsilon^{mnpq} X^n X^{\dagger p} X^q \quad (2.28)$$

is thus the natural analogue of (2.14) for $r = 1$, since just as X^m and $X^{\dagger m}$

$$X_m^+ = X^m \quad \text{and} \quad X_m^- = X^{\dagger m} \quad (2.29)$$

are bifundamental and conjugate bifundamental fields respectively. The constraints (2.15), originating from the fact that X_m^+ and X_m^- go from V_+ to V_- and vice-versa, are also immediately satisfied due to the bifundamental nature of X^m .

Thus for $r = 1$ the GR fuzzy three-sphere, as seen in the new description (2.14), (2.15), matches with the fuzzy S^3 solution of the \mathcal{A}_4 -theory. This is not too surprising since for $r = 1$ the fuzzy 3-sphere coordinates (2.1) reduce to just $\text{SO}(4)$ Γ -matrices. Yet, as can be easily seen from the above, the identification will not extend to $r > 1$ (or equivalently, 3-algebras other than \mathcal{A}_4 which are not necessarily $\mathcal{N} = 8$), since the $\text{SO}(4)$ relation (2.24) was crucial in turning (2.21) into (2.27), which can in turn be related to the GR fuzzy three-sphere.

2.3 ABJM theory

To have any hope of seeing the $r > 1$ realisation of the GR fuzzy S^3 , one should depart from the \mathcal{A}_4 -theory and go to the ABJM model [14]. These are superconformal CS-matter theories generalising [9] beyond $SU(2) \times SU(2)$ gauge groups, which in turn break $\mathcal{N} = 8$ supersymmetry and $SO(8)$ R-symmetry. The choice leading to the theory dual to M2-branes on a $\mathbb{C}_4/\mathbb{Z}_k$ singularity is the one with $U(N) \times U(N)$, which is the case that we will be focusing on. One could still consider the $SU(N) \times SU(N)$ case, which reduces to the \mathcal{A}_4 -theory for $N = 2$, but for which no M-theory dual interpretation has yet been found. Both these generically have explicit $\mathcal{N} = 6$ supersymmetry and $SU(4)$ R-symmetry. Different kinds of gauge groups like $U(M) \times U(N)$ and $O(M) \times Sp(N)$ have also been considered in the literature [32, 77]. A classification of all possibilities was made by [78, 79]. Since in this case one has bifundamental matter fields right from the beginning, they will automatically satisfy (2.15) and we might hope that we can obtain the GR fuzzy S^3 .

The authors of [80] expressed the original ABJM Lagrangian in component form. It employs $U(N) \times U(N)$ bifundamental matrix fields X^M , organised into the complex variables

$$Z^1 = X^1 + iX^2; \quad Z^2 = X^3 + iX^4; \quad W_1 = X^{5\dagger} + iX^{6\dagger}; \quad W_2 = X^{7\dagger} + iX^{8\dagger}, \quad (2.30)$$

that is $Z^\alpha, Z_\alpha^\dagger, W_{\dot{\alpha}}, W^{\dot{\alpha}\dagger}$, with $\alpha, \dot{\alpha} = 1, 2$. Then, in terms of $C^I = (Z^\alpha, W^{\dot{\alpha}\dagger})$ and C_I^\dagger , one gets the BPS fuzzy funnel equation (setting $W_{\dot{\alpha}} = 0$)

$$\partial_s Z^\alpha = -4a(Z^\beta Z_\beta^\dagger Z^\alpha - Z^\alpha Z_\beta^\dagger Z^\beta), \quad (2.31)$$

where a some normalisation constant. It was suggested [33–35] that this gives a fuzzy S^3 funnel, so we will now test whether one can obtain the GR fuzzy S^3 from it.

Since we have seen that for the \mathcal{A}_4 -theory we employed the identification (2.29), and we know how to obtain it as a particular case of the $SU(N) \times SU(N)$ ABJM theory for $N = 2$, we write the ansatz

$$\begin{aligned} Z^1 &= X_1^+ + iX_2^+, & Z_1^\dagger &= X_1^- - iX_2^- \\ Z^2 &= X_3^+ + iX_4^+, & Z_2^\dagger &= X_3^- - iX_4^-, \end{aligned} \quad (2.32)$$

which should certainly be correct for the $r = 1$ fuzzy sphere, but hopefully more.

Now, consider the fuzzy S^3 equivalent of (2.31), studied in [33–35], which can be obtained through the ansatz

$$Z^\alpha = \frac{1}{\sqrt{8as}} G^\alpha \quad (2.33)$$

and leads to

$$G^\alpha = G^\beta G_\beta^\dagger G^\alpha - G^\alpha G_\beta^\dagger G^\beta. \quad (2.34)$$

The real part for $\alpha = 1$ then gives

$$\text{Re}\{G^1\} = \text{Re}\{G^\beta G_\beta^\dagger G^1 - G^1 G_\beta^\dagger G^\beta\}. \quad (2.35)$$

But with the ansatz (2.32) for G^α , the right hand side becomes

$$\begin{aligned} \text{Re}\{G^\beta G_\beta^\dagger G^1 - G^1 G_\beta^\dagger G^\beta\} &= X_2^+ X_3^- X_4^+ - X_2^+ X_4^- X_3^+ + X_3^+ X_4^- X_2^+ - X_4^+ X_3^- X_2^+ \\ &\quad + X_3^+ X_3^- X_1^+ + X_4^+ X_4^- X_1^+ - X_1^+ X_3^- X_3^+ - X_1^+ X_4^- X_4^+. \end{aligned} \quad (2.36)$$

For $r = 1$, when the fuzzy three-sphere coordinates reduce to $\text{SO}(4)$ Γ -matrices, one has that $X_m^+ X_n^- = -X_n^+ X_m^-$ and $X_m^- X_n^+ = -X_n^- X_m^+$ for $m \neq n$, while zero for $m = n$. This in turn means that (2.36) reduces to just $4X_2^+ X_3^- X_4^+$. Compared to the left hand side of the first equation in (2.14) for $m = 1$, the difference is only a relative numerical factor of 4. Hence up to a constant rescaling there is a matching between (2.35) and (2.14), as expected.

Since we have just seen the matching for $r = 1$, it will serve as a test for the formalism for $r > 1$. However, we now have an r -dependent factor on the right hand side of (2.14), which means (2.36) should end up being proportional to

$$\frac{r+2}{2[(r+1)(r+3)+1]} \epsilon^{1mnp} X_m^+ X_n^- X_p^+ = \frac{r+2}{2[(r+1)(r+3)+1]} \left(X_2^+ X_3^- X_4^+ - X_2^+ X_4^- X_3^+ \right. \\ \left. + X_3^+ X_4^- X_2^+ - X_4^+ X_3^- X_2^+ + X_4^+ X_2^- X_3^+ - X_3^+ X_2^- X_4^+ \right). \quad (2.37)$$

The first line in (2.36) matches the first four terms of (2.37), but that is where the agreement stops. The best case scenario would be if the final result were proportional to (2.37), yet still give $4X_2^+ X_3^- X_4^+$ for the $r = 1$ case, *i.e.* if the following were true

$$X_3^+ X_3^- X_1^+ + X_4^+ X_4^- X_1^+ - X_1^+ X_3^- X_3^+ - X_1^+ X_4^- X_4^+ \\ = \frac{2}{3} [X_4^+ X_2^- X_3^+ - X_3^+ X_2^- X_4^+] - \frac{1}{3} [X_2^+ X_3^- X_4^+ - X_2^+ X_4^- X_3^+ + X_3^+ X_4^- X_2^+ - X_4^+ X_3^- X_2^+], \quad (2.38)$$

where the relative factor of 2 between the two brackets is such that the right hand side vanishes for $r = 1$, and then the $\frac{2}{3}$ and $\frac{1}{3}$ factors are fixed by the requirement that the final result is proportional to (2.37). But even if that were the case, one would end up with the same numerical factor on the right hand side of (2.36) for any value of $r > 1$, instead of the r -dependent factor required by (2.37).

Thus the GR fuzzy S^3 is not a solution to (2.34) for any $r > 1$, in agreement with the expectation that one would have due to the absence of $\text{SO}(4)$ R-symmetry for the above solution, which instead only sports an $\text{SU}(2) \times \text{U}(1)$. In the following sections we will show that for $r > 1$ the geometry of these solutions is instead a fuzzy S^2 .

3 Fuzzy two-sphere structure of M2-M5 ABJM solutions

As we have discussed in the previous section, the BPS equation for the ABJM scalars (2.31) was conjectured to give a fuzzy 3-funnel, with the proposed 3-sphere equation (2.34). Equivalently, the authors of [33] obtained a mass deformation of the ABJM-theory preserving maximal ($\mathcal{N} = 6$) supersymmetry, but breaking R-symmetry to $\text{SU}(2) \times \text{SU}(2) \times \text{U}(1) \subset \text{SU}(4)$. The mass-deformed theory has two sets of ground states written in terms of two sets of scalars R^α and $Q^{\dot{\alpha}}$, the equivalent of Z^α and $W_{\dot{\alpha}}$ for the undeformed ABJM theory.⁴ One set of ground-states corresponds to $Q^{\dot{\alpha}} = 0$ and R^α satisfying

$$-R^\alpha = \frac{2\pi}{\mu k} \left(R^\beta R_\beta^\dagger R^\alpha - R^\alpha R_\beta^\dagger R^\beta \right). \quad (3.1)$$

⁴We will give the ABJM and mass-deformed ABJM actions in due course. At the moment we wish to focus entirely on the matrix structure of the funnel/ground-state solutions.

The other set has $R^\alpha = 0$ with Q_α^\dagger satisfying the equation (2.34). Both sets were given the interpretation of a 3-sphere. Note that (3.1) is the same equation as (2.34). Hence, the above are directly related to the funnel solutions of the undeformed ABJM theory and there is no difference in looking at the proposed ‘fuzzy 3-funnel’ solution of ABJM or the proposed ‘fuzzy 3-sphere’ ground-states of the mass-deformed theory. As a result, the latter solutions cannot admit the GR 3-sphere interpretation either. In the following we will switch to the study of the ground-state solutions of [33] for concreteness and examine their symmetries to find that they are actually the ones of a fuzzy 2-sphere. We will later extrapolate the results to the funnel.

Notation: In this section, we will denote by k, l, m, n the matrix indices/indices of states in a vector space, while keeping $i, j = 1, \dots, 3$ as vector indices on the fuzzy S^2 . We will also use j for the SU(2) spin and Y_{lm} for S^2 spherical harmonics, following the standard notation. The distinction should be clear by the context.

3.1 Ground-state matrices and symmetries

The ground-state solutions to (2.34) were found in [33] and are given by the following set of matrices

$$\begin{aligned} (G^1)_{m,n} &= \sqrt{m-1} \delta_{m,n} \\ (G^2)_{m,n} &= \sqrt{(N-m)} \delta_{m+1,n} \\ (G_1^\dagger)_{m,n} &= \sqrt{m-1} \delta_{m,n} \\ (G_2^\dagger)_{m,n} &= \sqrt{(N-n)} \delta_{n+1,m} . \end{aligned} \quad (3.2)$$

Using the decomposition (2.30) of the fuzzy complex coordinates G^α into real coordinates X_p , these satisfy

$$\sum_{p=1}^4 X_p X^p \equiv G^\alpha G_\alpha^\dagger = N - 1 , \quad (3.3)$$

which at first glance would seem to indicate a fuzzy S^3 structure.

However, note that in the above $G^1 = G_1^\dagger$ for the ground-state solution. With the help of (2.30) this seems to suggest that $X_2 = 0$ which is instead indicative of a fuzzy S^2 structure. Nevertheless, given the fact that the $r = 1$ solution is in fact a fuzzy S^3 , one might still be open to the possibility that the symmetries and fluctuations of this solution could tell a different story.

In the following, and as is usual in the case of fuzzy sphere constructions, the matrices G^α will be used to construct both the symmetry operators (as bilinears in G, G^\dagger and acting on G^α themselves) and fuzzy coordinates (used to expand in terms of spherical harmonics on the fuzzy sphere).

3.1.1 GG^\dagger relations

We start by calculating the GG^\dagger bilinears

$$\begin{aligned} (G^1 G_1^\dagger)_{m,n} &= (m-1) \delta_{mn} \\ (G^2 G_2^\dagger)_{mn} &= (N-m) \delta_{mn} \\ (G^1 G_2^\dagger)_{mn} &= \sqrt{(m-1)(N-m+1)} \delta_{m,n+1} \\ (G^2 G_1^\dagger)_{mn} &= \sqrt{(N-m)m} \delta_{m+1,n} \end{aligned}$$

$$(G^\alpha G^\dagger_\alpha)_{mn} = (N - 1) \delta_{mn} \quad (3.4)$$

and examining their commutation relations. Defining $J_\beta^\alpha = G^\alpha G^\dagger_\beta$ we get

$$[J_\beta^\alpha, J_\nu^\mu] = \delta_\beta^\mu J_\nu^\alpha - \delta_\nu^\alpha J_\beta^\mu . \quad (3.5)$$

These are commutation relations of the generators of $U(2)$. The traceless combinations $\hat{J}_\beta^\alpha = J_\beta^\alpha - \frac{1}{2} J \delta_\beta^\alpha$ are generators of $SU(2)$ and can equivalently be given in terms of the usual angular momentum-type variables $J_i = (\tilde{\sigma}_i)_\beta^\alpha J_\alpha^\beta$,⁵ where $\tilde{\sigma}_i = \sigma_i^T$ are the transpose of the Pauli matrices. Note that more correctly, we should have written $J^\alpha_\beta = G^\alpha G^\dagger_\beta$ and

$$J_i = (\tilde{\sigma}_i)_\beta^\alpha J^\beta_\alpha = (\sigma_i)_\beta^\alpha J^\beta_\alpha , \quad (3.6)$$

but in the following we will stick to the notation J_β^α , as what kind of matrix multiplication we have will be made clear from the context. The commutation relations of J_i can be calculated as

$$[J_i, J_j] = 2i\epsilon_{ijk} J_k . \quad (3.7)$$

Note that the trace $J \equiv J^\alpha_\alpha = N - 1$ is a trivial $U(1) \simeq U(2)/SU(2)$ generator, commuting with everything.

3.1.2 $G^\dagger G$ relations

Next, we calculate the $G^\dagger G$ matrices

$$\begin{aligned} (G^\dagger_1 G^1)_{mn} &= (m - 1) \delta_{mn} \\ (G^\dagger_2 G^2)_{mn} &= (N - m + 1) \delta_{mn} - N \delta_{m1} \delta_{n1} \\ (G^\dagger_1 G^2)_{mn} &= \sqrt{(m - 1)(N - m)} \delta_{m+1, n} \\ (G^\dagger_2 G^1)_{mn} &= \sqrt{(m - 2)(N - m + 1)} \delta_{m, n+1} \\ (G^\dagger_\alpha G^\alpha)_{mn} &= N \delta_{mn} - N \delta_{m1} \delta_{n1} \end{aligned} \quad (3.8)$$

and define $\bar{J}_\alpha^\beta = G^\dagger_\alpha G^\beta$. The commutation relations then are

$$[\bar{J}_\beta^\alpha, \bar{J}_\nu^\mu] = \delta_\beta^\mu \bar{J}_\nu^\alpha - \delta_\nu^\alpha \bar{J}_\beta^\mu . \quad (3.9)$$

Similarly, we define traceless combinations $\hat{\bar{J}}_\beta^\alpha = \bar{J}_\beta^\alpha - \frac{1}{2} \bar{J} \delta_\beta^\alpha$ which are generators of $SU(2)$ and their angular momentum-type variable counterparts $\bar{J}_i = (\tilde{\sigma}_i)_\beta^\alpha \bar{J}_\alpha^\beta$ satisfy

$$[\bar{J}_i, \bar{J}_j] = 2i\epsilon_{ijk} \bar{J}_k . \quad (3.10)$$

Again, note that we should have written $\bar{J}_\alpha^\beta = G^\dagger_\alpha G^\beta$ which emphasises that for \bar{J} , the lower index is the first matrix index, and

$$\bar{J}_i = (\tilde{\sigma}_i)_\beta^\alpha \bar{J}_\alpha^\beta = (\sigma_i)_\beta^\alpha \bar{J}_\alpha^\beta , \quad (3.11)$$

which emphasises that as matrices, \bar{J}_i is defined with the Pauli matrices, whereas J_i was defined with their transpose. However, we will keep the notation \bar{J}_α^β , as which matrix contraction one has will once again be made clear from the context . The trace

$$(\bar{J})_{mn} = (\bar{J}_\alpha^\alpha)_{mn} = N \delta_{mn} - N \delta_{m1} \delta_{n1} , \quad (3.12)$$

⁵This observation was independently made by D. Rodríguez-Gómez.

which is a $U(1) \simeq U(2)/SU(2)$ generator, commutes with the $SU(2)$ generators \bar{J}_i , though as a matrix does not commute with the generators J_2^1 and J_1^2 of the first set of $SU(2)$ generators.

At this point, it seems that we have two $SU(2)$'s, *i.e.* $SO(4) \simeq SU(2) \times SU(2)$ as expected for a 3-sphere, but we will see that these are not independent.

3.2 Fuzzy S^2 harmonics from $U(N) \times U(\bar{N})$ with bifundamentals

The first important observation is that G^α and G_α^\dagger are bifundamental matrices of $U(N) \times U(\bar{N})$, mapping between two different vector spaces. However, one can construct combinations of the above that map each vector space back to itself, *e.g.* the form that we have just seen, GG^\dagger , being examples of fundamental matrices for the vector space corresponding to the group $U(N)$ and $G^\dagger G$ for the vector space corresponding to $U(\bar{N})$. We next analyse this structure.

3.2.1 The adjoint of $U(N)$

The matrices GG^\dagger are acting on an N dimensional vector space that we will call \mathbf{V}^+ . The space of linear maps from \mathbf{V}^+ back to itself, $End(\mathbf{V}_+)$, is the adjoint of the $U(N)$ factor in the $U(N) \times U(\bar{N})$ gauge group and GG^\dagger are examples of matrices belonging to it. The space \mathbf{V}^+ forms an irreducible representation of $SU(2)$ of spin $j = \frac{N-1}{2}$, denoted by V_N

$$\mathbf{V}^+ = V_N . \quad (3.13)$$

The set of all operators of the form $GG^\dagger, GG^\dagger GG^\dagger, \dots$ belong in $End(\mathbf{V}_+)$ and can be expressed in terms of a basis of ‘fuzzy spherical harmonics’ defined using the $SU(2)$ structure. Through the $SU(2)$ generators J_i we can form the fuzzy spherical harmonics as

$$\begin{aligned} Y^0 &= 1 \\ Y_i^1 &= J_i \\ Y_{(i_1 i_2)}^2 &= J_{(i_1} J_{i_2)} \\ Y_{(i_1 \dots i_l)}^l &= J_{(i_1} \dots J_{i_l)} . \end{aligned} \quad (3.14)$$

The brackets $(i_1 \dots i_l)$ denote symmetrisation. It is known that symmetrised elements with $l > 2j = N - 1$ can be reexpressed in terms of lower l elements. The complete space of $N \times N$ matrices can be expressed in terms of the fuzzy spherical harmonics with $0 \leq l \leq 2j = N - 1$. One indeed checks that

$$N^2 = \sum_{l=0}^{2j} (2l+1)^2 . \quad (3.15)$$

Then, a general matrix in the adjoint of $U(N)$ is expanded as

$$A = \sum_{l=0}^{N-1} a^{lm} Y_{lm}(J_i) , \quad (3.16)$$

where

$$Y_{lm}(J_i) = \sum_i f_{lm}^{(i_1 \dots i_l)} J_{i_1} \dots J_{i_l} . \quad (3.17)$$

The $Y_{lm}(J_i)$ become the usual spherical harmonics in the ‘classical’ limit, when $N \rightarrow \infty$ and the cut-off in the angular momentum is removed.

In conclusion, all the matrices of $U(N)$ can be organised into irreps of $SU(2)$ constructed out of J_i , which form the fuzzy spherical harmonics $Y_{lm}(J_i)$.

3.2.2 The adjoint of $U(\bar{N})$

Similarly to the $U(N)$ case, the matrices $G^\dagger G, G^\dagger G G^\dagger G, \dots$, are linear endomorphisms of \mathbf{V}^- . These matrices are in the adjoint of the $U(\bar{N})$ factor of the $U(N) \times U(\bar{N})$ gauge group, and will be organised into irreps of the $SU(2)$ constructed out of \bar{J}_i .

However, we now have an extra ingredient: We have already noticed in (3.12) that the $U(1)$ generator \bar{J} is nontrivial. We can express it as

$$\bar{J} = G_\alpha^\dagger G^\alpha = N - N \bar{E}_{11} . \quad (3.18)$$

This means that $End(\mathbf{V}_-)$ contains in addition to the identity matrix, the matrix \bar{E}_{11} which is invariant under $SU(2)$. If we label the basis states in \mathbf{V}^- as $|e_k^- \rangle$ with $k = 1, \dots, N$, then $\bar{E}_{11} = |e_1^- \rangle \langle e_1^-|$. This in turn means that \mathbf{V}^- is a reducible representation

$$\mathbf{V}^- = V_{N-1}^- \oplus V_1^- . \quad (3.19)$$

The first direct summand is the irrep of $SU(2)$ with dimension $N - 1$ while the second is the one-dimensional irrep. Indeed, one checks that the \bar{J}_i ’s annihilate the state $|e_1^- \rangle$, which is necessary for the identification with the one-dimensional irrep to make sense. This follows because $G^1 |e_1^- \rangle = G^2 |e_1^- \rangle = 0$. Equivalently one has

$$G^\alpha \bar{E}_{11} = 0 = \bar{E}_{11} G_\beta^\dagger . \quad (3.20)$$

As a result, the space $End(\mathbf{V}^-)$ decomposes as follows

$$End(\mathbf{V}^-) = End(V_{N-1}^-) \oplus End(V_1^-) \oplus Hom(V_{N-1}^-, V_1^-) \oplus Hom(V_1^-, V_{N-1}^-) . \quad (3.21)$$

The first summand has a decomposition in terms of another set of fuzzy spherical harmonics

$$Y_{lm}(\bar{J}_i) = \sum_i f_{lm}^{(i_1 \dots i_l)} \bar{J}_{i_1} \dots \bar{J}_{i_l} , \quad (3.22)$$

for l going from 0 to $N - 2$, since

$$(N - 1)^2 = \sum_{l=0}^{N-2} (2l + 1) . \quad (3.23)$$

However, in this case one is only getting the matrices in the $(N - 1)$ block, *i.e.* the $End(V_{N-1}^-)$. The second summand is just one matrix transforming in the trivial irrep. The remaining two $N - 1$ dimensional spaces of matrices cannot be expressed as products of \bar{J}_i . They are spanned by

$$\bar{E}_{1k} = |e_1^- \rangle \langle e_k^-| \equiv g_{1k}^-$$

$$\bar{E}_{k1} = |e_k^-\rangle\langle e_1^-| \equiv g_{k1}^{--}, \quad (3.24)$$

which are the equivalent of zero mode spherical harmonics for $Hom(V_{N-1}^-, V_1^-) \oplus Hom(V_1^-, V_{N-1}^-)$.

Indeed, usual spherical harmonics Y_{lm} are eigenfunctions of the symmetry operators \bar{L}^2 and L_3 , whereas $\bar{E}_{1k} = g_{1k}^{--}$ is a zero mode eigenfunction of the U(1) symmetry operator \bar{J} , *i.e.*

$$\bar{J}\bar{E}_{1k} = 0 \cdot \bar{E}_{1k} \quad (3.25)$$

just as on a circle, we have⁶

$$R\partial_x e^{\frac{2\pi i n x}{R}} = 0 \cdot e^{\frac{2\pi i n x}{R}} \Rightarrow n = 0. \quad (3.26)$$

Hence we have the ‘zero mode spherical harmonics’ g_{1k}^{--} and g_{k1}^{--}

$$\bar{J}g_{1k}^{--} = 0 \cdot g_{1k}^{--}; \quad g_{k1}^{--}\bar{J} = 0 \cdot g_{k1}^{--}. \quad (3.27)$$

These also transform in the $N - 1$ dimensional irrep of SU(2) under the adjoint action of \bar{J}_i .

Therefore, one can expand a general matrix in the adjoint of U(\bar{N}) as

$$\bar{A} = \bar{a}_0 \bar{E}_{11} + \sum_{l=0}^{N-2} \bar{a}_{lm} Y_{lm}(\bar{J}_i) + \sum_{k=2}^N b_k g_{1k}^{--} + \sum_{k=2}^N \bar{b}_k g_{k1}^{--} \quad (3.28)$$

and note that we could have replaced \bar{E}_{11} with the U(1) generator \bar{J} by redefining \bar{a}_0 and \bar{a}_{00} .

In the large N limit the $Y_{lm}(\bar{J}_i)$ become ordinary spherical harmonics of a scalar field on S^2 , just like $Y_{lm}(J_i)$. There are order N^2 of these modes, which is appropriate if one roughly think of the fuzzy S^2 as a space with linear dimensions discretised in N units. The mode \bar{a}_0 can be neglected at large N . The modes b_k and \bar{b}_k have order N degrees of freedom, and we will see that they will also become irrelevant at large N , as expected.

3.2.3 Physical fluctuations: bifundamental (N, \bar{N}) matrices

All the (N, \bar{N}) bifundamental scalar matrices are of the type $G, GG^\dagger G, GG^\dagger GG^\dagger G, \dots$ and form maps from \mathbf{V}^- to \mathbf{V}^+ , *i.e.* the space $Hom(\mathbf{V}^-, \mathbf{V}^+)$. Since we intend to perform a fluctuation analysis of the mass-deformed ABJM theory, in which the physical degrees of freedom are bifundamental, we will also need to find out how the symmetries act on these matrices.

It is easy to check that the matrices G^a satisfy

$$\begin{aligned} G^1 G_2^\dagger G^2 - G^2 G_2^\dagger G^1 &= G^1 \\ G^2 G_1^\dagger G^1 - G^1 G_1^\dagger G^2 &= G^2. \end{aligned} \quad (3.29)$$

Using the definitions $J_3 = J_1^1 - J_2^2$, $J_+ = J_1 + iJ_2 = J_2^1$, $J_- = J_1 - iJ_2 = J_1^2$ and likewise $\bar{J}_3 = \bar{J}_1^1 - \bar{J}_2^2$, $\bar{J}_+ = \bar{J}_2^1$, $\bar{J}_- = \bar{J}_1^2$, we find

$$\begin{aligned} J_+ G^2 - G^2 \bar{J}_+ &= G^1 \\ J_- G^1 - G^1 \bar{J}_- &= G^2. \end{aligned} \quad (3.30)$$

⁶ \bar{J} is the U(1) Lie algebra element, just like $R\partial_x$.

Notice that J_i by itself does not give a nice transformation of G^α . The G^α form a nice representation of an action constructed from both J_i and \bar{J}_i . This means that the geometry we will be constructing from bifundamental fluctuation modes has a single SU(2) symmetry, as opposed to two.

In fact, it is easy to see how we would have needed the above to be modified in order to obtain the full $SU(2) \times SU(2) \simeq SO(4)$ symmetry: The J_i and \bar{J}_i ought to act independently. For instance, in [81], the fuzzy S^3 was obtained from the BMN Plane Wave Matrix Model [82] through the intermediate step of a fuzzy S^2 . There, the matrices X_i were divided into $N_s \times N_t$ subsets $X_i^{(s,t)}$. Then the SU(2) generators in the j_s representation, $L_i^{(j_s)}$, acted separately on the two sides, giving an action expressed in terms of

$$L_i \circ X_i^{(s,t)} = L_i^{(j_s)} X_i^{(s,t)} - X_i^{(s,t)} L_i^{(j_t)}, \quad (3.31)$$

with an infinite set of s, t values for each side (in the large N limit), that can be used to create an extra coordinate.

Under this single SU(2) symmetry, Eq. (3.30), G^1 is the up-state of spin $\frac{1}{2}$, while G^2 is the down-state. This implies that we should also have

$$\begin{aligned} J_+ G^1 - G^1 \bar{J}_+ &= 0 \\ J_- G^2 - G^2 \bar{J}_- &= 0, \end{aligned} \quad (3.32)$$

which follow trivially by writing out the SU(2) generators. One also checks that

$$\begin{aligned} J_3 G^1 - G^1 \bar{J}_3 &= G^1 \\ J_3 G^2 - G^2 \bar{J}_3 &= -G^2. \end{aligned} \quad (3.33)$$

These in fact follow from (3.29). We can summarise all of the above relations by writing

$$J_i G^\alpha - G^\alpha \bar{J}_i = (\tilde{\sigma}_i)_\beta^\alpha G^\beta. \quad (3.34)$$

The G^1, G^2 transform like the (1,0) and (0,1) column vectors of the spin- $\frac{1}{2}$ rep. Note that the J 's and \bar{J} 's are matrices in the $U(N) \times U(\bar{N})$ Lie algebra.

By taking Hermitian conjugates in (3.34), we find that the G_α^\dagger transform as

$$G_\alpha^\dagger J_i - \bar{J}_i G_\alpha^\dagger = G_\beta^\dagger (\tilde{\sigma}_i)_\alpha^\beta. \quad (3.35)$$

To make the relation between the above and the $U(N) \times U(\bar{N})$ gauge group clear, it is useful to construct matrices acting on $\mathbf{V}^+ \oplus \mathbf{V}^-$, *i.e.* $2N \times 2N$ matrices. The G^α and G_α^\dagger matrices are bifundamental, so we have

$$\mathbf{G}^\alpha = \begin{pmatrix} 0 & G^\alpha \\ G_\alpha^\dagger & 0 \end{pmatrix}, \quad (3.36)$$

while the J_i, \bar{J}_i matrices are adjoint and

$$\mathbf{J}_i = \begin{pmatrix} J_i & 0 \\ 0 & \bar{J}_i \end{pmatrix}. \quad (3.37)$$

Hence, the G^α and G_a^\dagger transformation laws can be summarised as a single law acting in the $\mathbf{V}^+ \oplus \mathbf{V}^-$ space, by

$$\begin{aligned} [\mathbf{J}_i, \mathbf{G}^\alpha] &= \begin{pmatrix} 0 & J_i G^\alpha - G^\alpha \bar{J}_i \\ \bar{J}_i G_\alpha^\dagger - G_\alpha^\dagger J_i & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & (\tilde{\sigma}_i)_\beta^\alpha G^\beta \\ -G_\beta^\dagger (\tilde{\sigma}_i)_\alpha^\beta & 0 \end{pmatrix}. \end{aligned} \quad (3.38)$$

The matrices \mathbf{J}_i are elements of the $U(N) \times U(\bar{N})$ gauge Lie algebra. Their action on G^α is precisely the gauge symmetry action corresponding to the fact that the matrices R^α transform as bifundamentals. This embedding of a global symmetry into the gauge symmetry is an example of the mixing of symmetries in soliton physics and has interesting physical consequences [83]. In this case, the global $SU(2)$ rotations $\int d^3x R^\alpha D_0 R_\beta^\dagger$ do not leave the solution invariant. But combining the global symmetries with rotations generated by J_i gives symmetries which do leave the solution invariant. Hence the geometric $SU(2)$ does survive as an invariance of the solution and acts on the space of fluctuations.

3.2.4 Action of the full $SU(2) \times U(1)$

Writing the action of the full $SU(2) \times U(1)$ on the G^α , including the $U(1)$ trace \bar{J} , we obtain

$$J_\beta^\alpha G^\gamma - G^\gamma \bar{J}_\beta^\alpha = \delta_\beta^\gamma G^\alpha - \delta_\beta^\alpha G^\gamma. \quad (3.39)$$

This implies the relations⁷

$$\begin{aligned} G_\gamma^\dagger J_\beta^\alpha G^\gamma &= (N+1)\bar{J}_\beta^\alpha - N\delta_\beta^\alpha + N\delta_\beta^\alpha \bar{E}_{11} \\ G^\gamma \bar{J}_\beta^\alpha G_\gamma^\dagger &= (N-2)J_\beta^\alpha + (N-1)\delta_\beta^\alpha. \end{aligned} \quad (3.40)$$

In terms of the J_i 's these equations are simpler

$$\begin{aligned} G_\gamma^\dagger J_i G^\gamma &= (N+1)\bar{J}_i \\ G^\gamma \bar{J}_i G_\gamma^\dagger &= (N-2)J_i, \end{aligned} \quad (3.41)$$

while taking Hermitian conjugates of (3.39) we obtain the $U(2)$ transformation of G_α^\dagger ,

$$\bar{J}_\beta^\alpha G_\gamma^\dagger - G_\gamma^\dagger J_\beta^\alpha = -\delta_\gamma^\alpha G_\beta^\dagger + \delta_\beta^\alpha G_\gamma^\dagger. \quad (3.42)$$

The consequence of the above equations is that G^α has charge 1 under the $U(1)$ generator \bar{J} . This means that a global $U(1)$ symmetry action on R^α does not leave the solution invariant, but when combined with the action of \bar{J} from the gauge group does lead to an invariance.

3.3 $SU(2)$ harmonic decomposition of bifundamental matrices

As in the case of the $U(\bar{N})$ matrices, the bifundamental matrices of the form $G, GG^\dagger G, \dots$ giving physical fluctuating fields, are not enough to completely fill $Hom(\mathbf{V}^-, \mathbf{V}^+)$. Given the decomposition $\mathbf{V}^- = V_{N-1}^- \oplus V_1^-$, we decompose $Hom(\mathbf{V}^-, \mathbf{V}^+)$ as

$$Hom(\mathbf{V}^-, \mathbf{V}^+) = Hom(V_{N-1}^-, V_N^+) \oplus Hom(V_1^-, V_N^+). \quad (3.43)$$

⁷To obtain the first line one uses $\bar{E}_{11}\bar{J}_\beta^\alpha = 0$, since $\bar{E}_{11}G_\beta^\dagger = 0$.

The first summand has dimension $N(N-1)$. The second has dimension N . The N matrices in $Hom(V_1^-, V_N^+)$ form an irreducible representation of $SU(2)$ of dimension N .

Since the V_{N-1}^- and V_N^+ are irreps. of $SU(2)$ we can label the states with the eigenvalue of \bar{J}_3, J_3 respectively. Given the normalisation of the $SU(2)$ generators in (3.7), the conventionally defined spin is $\frac{J_3^{max}}{2}$. The matrices in $Hom(V_{N-1}^-, V_N^+)$ are of the form $|e_m^+\rangle\langle e_n^-|$, where $m = \frac{-N+1}{2}, \frac{-N+3}{2}, \dots, \frac{N-1}{2}$, $n = \frac{-N+2}{2}, \frac{-N+4}{2}, \dots, \frac{N-2}{2}$ denote the eigenvalues of $\frac{J_3}{2}$. These are spanned by matrices of the form $G(\bar{J}_{i_1})(\bar{J}_{i_2}) \cdots (\bar{J}_{i_l})$, *i.e.* the matrix G times matrices in $End(V_{N-1}^-)$. To understand the cutoff on the maximum value of l in the above ‘spherical harmonics’, note that acting on the lowest weight state $|e_{\frac{-N+2}{2}}^-\rangle$ we can have $G^1(\bar{J}_+)^{N-2}$ to obtain the highest weight state $|e_{\frac{N-1}{2}}^+\rangle$. Operators of the form $G^2(\bar{J}_+)^{N-1}$ can be rewritten to have G^1 first by using the relations (3.30).

The upshot is that operators in $Hom(V_{N-1}^-, V_N^+)$ transform in representations of spin $l + \frac{1}{2}$ for $l = 0, \dots, N-2$. The dimensions of these representations add up to

$$\sum_{l=0}^{N-2} (2l+2) = N(N-1). \quad (3.44)$$

This then gives the $SU(2)$ decomposition of $Hom(V_{N-1}^-, V_N^+)$ as follows

$$Hom(V_{N-1}^-, V_N^+) = \bigoplus_{l=0}^{N-2} V_{l+1/2}. \quad (3.45)$$

On the other hand, the matrices in $Hom(V_1^-, V_N^+)$ cannot be written in terms of the G ’s and G^\dagger ’s alone. Indeed, matrices $|e_k^+\rangle\langle e_1^-| \equiv \hat{E}_{k1} \in Hom(V_1^-, V_N^+)$ cannot be so expressed, because G^α acting on $|e_1^- \rangle$ gives zero. The index k runs over the N states in \mathbf{V}^+ . The operators \hat{E}_{k1} are eigenfunctions of the operator \bar{E}_{11}

$$\hat{E}_{k1} \bar{E}_{11} = \hat{E}_{k1}, \quad (3.46)$$

with unit charge. They span the $Hom(V_1^-, V_N^+)$ matrices.

Combining all of the above, the fluctuations r^α of the bifundamental fields R^α can be expanded in spherical harmonics as follows

$$r^\alpha = r_\beta^\alpha G^\beta + \sum_{k=1}^N t_k^\alpha \hat{E}_{k1}, \quad (3.47)$$

with

$$r_\beta^\alpha = \sum_{l=0}^{N-2} (r^{lm})_\beta^\alpha Y_{lm}(J_i). \quad (3.48)$$

We then decompose r_β^α into a trace and a traceless part and define the fluctuating fields

$$\begin{aligned} s_\beta^\alpha &= r_\beta^\alpha - \frac{1}{2} \delta_\beta^\alpha r_\gamma^\gamma \\ r &= r_\gamma^\gamma \\ T^\alpha &= t_k^\alpha \hat{E}_{k1}. \end{aligned} \quad (3.49)$$

Thus the complete expansion of the fluctuating field r^α is given simply in terms of

$$r^\alpha = r G^\alpha + s_\beta^\alpha G^\beta + T^\alpha. \quad (3.50)$$

We could also have used equivalently

$$r^\alpha = \sum_{l=0}^{N-2} (r^{lm})^\alpha_\beta G^\beta Y_{lm}(\bar{J}_i) + \sum_{k=1}^N t_k^\alpha \hat{E}_{k1} \quad (3.51)$$

using the spherical harmonics in \bar{J} in (3.28). This a signature for a \mathbb{Z}_2 symmetry of the fluctuation action that will become manifest as we progress in our analysis. We will choose, without loss of generality, to work with (3.50).

Until now we have focused on matrices in $Hom(\mathbf{V}^-, \mathbf{V}^+)$ but the case of $Hom(\mathbf{V}^+, \mathbf{V}^-)$ is similar. The matrices $G^\dagger, G^\dagger G G^\dagger, \dots$ will also form a representation of $SU(2)$ given by $\bar{J} \sim G^\dagger G$, times a G^\dagger matrix. And again matrices $\hat{F}_{1k} \equiv |e_1^-\rangle \langle e_k^+| \in Hom(V_N^+, V_1^-)$ cannot be so expressed, so one needs to add an extra $T_\alpha^\dagger = (t_k^\alpha)^* \hat{F}_{1k}$ fluctuation. In fact, the result for the complete fluctuating field can be obtained by taking a Hermitian conjugate of (3.50), yielding

$$r_\alpha^\dagger = G_\alpha^\dagger r + G_\beta^\dagger s_\alpha^\beta + T_\alpha^\dagger. \quad (3.52)$$

4 Action for matter fluctuations on the two-sphere

So far we have analysed the symmetries of the G^α matrices of (3.2) and found a single $SU(2)$. In this section we will begin computing the action for fluctuations around these ground-state solutions of the mass-deformed ABJM theory [33]. Our aim in the following sections will be to write the resultant action for the fluctuations in terms of a higher-dimensional worldvolume theory, which will make the existence of a fuzzy S^2 manifest.

The pure ABJM action is given by

$$\begin{aligned} S_{\text{ABJM}} = & \int d^3x \left[\frac{k}{4\pi} \epsilon^{\mu\nu\lambda} \text{Tr} \left(A_\mu^{(1)} \partial_\nu A_\lambda^{(1)} + \frac{2i}{3} A_\mu^{(1)} A_\nu^{(1)} A_\lambda^{(1)} - A_\mu^{(2)} \partial_\nu A_\lambda^{(2)} - \frac{2i}{3} A_\mu^{(2)} A_\nu^{(2)} A_\lambda^{(2)} \right) \right. \\ & - \text{Tr} \left(D_\mu C_I^\dagger D^\mu C^I \right) - i \text{Tr} \left(\psi^{I\dagger} \gamma^\mu D_\mu \psi_I \right) \\ & + \frac{4\pi^2}{3k^2} \text{Tr} \left(C^I C_I^\dagger C^J C_J^\dagger C^K C_K^\dagger + C_I^\dagger C^I C_J^\dagger C^J C_K^\dagger C^K \right. \\ & \left. + 4C^I C_J^\dagger C^K C_I^\dagger C^J C_K^\dagger - 6C^I C_J^\dagger C^J C_I^\dagger C^K C_K^\dagger \right) \\ & + \frac{2\pi i}{k} \text{Tr} \left(C_I^\dagger C^I \psi^{J\dagger} \psi_J - \psi^{\dagger J} C^I C_I^\dagger \psi_J - 2C_I^\dagger C^J \psi^{\dagger I} \psi_J + 2\psi^{\dagger J} C^I C_J^\dagger \psi_I \right. \\ & \left. + \epsilon^{IJKL} \psi_I C_J^\dagger \psi_K C_L^\dagger - \epsilon_{IJKL} \psi^{\dagger I} C^J \psi^{\dagger K} C^L \right) \Big], \quad (4.1) \end{aligned}$$

where on the first two lines we have the CS gauge fields and kinetic terms, on the 3rd and 4th lines $\int d^3x(-V_6)$, where V_6 is the scalar bosonic potential (sextic), and on the last two lines the fermionic interactions. We will treat the CS and scalar kinetic terms in the first two lines separately in the next section, as these will participate in a version of the Higgs mechanism. Focusing on the purely bosonic sector will also prove enough for our purposes, hence we will not study the fluctuations for the fermion fields in the rest of this paper, although doing so should be straightforward. Therefore, in this section we will exclusively discuss the scalar potential terms.

By splitting $C^I = (R^\alpha, Q^\alpha)$, the mass deformation changes the potential to

$$V = |M^\alpha|^2 + |N^\alpha|^2, \quad (4.2)$$

where

$$\begin{aligned} M^\alpha &= \mu Q^\alpha + \frac{2\pi}{k}(2Q^{[\alpha}Q_\beta^\dagger Q^{\beta]} + R^\beta R_\beta^\dagger Q^\alpha - Q^\alpha R_\beta^\dagger R^\beta + 2Q^\beta R_\beta^\dagger R^\alpha) \\ N^\alpha &= -\mu R^\alpha + \frac{2\pi}{k}(2R^{[\alpha}R_\beta^\dagger R^{\beta]} + Q^\beta Q_\beta^\dagger R^\alpha - R^\alpha Q_\beta^\dagger Q^\beta + 2R^\beta Q_\beta^\dagger Q^\alpha). \end{aligned} \quad (4.3)$$

In addition, the potential also involves a mass term μ for the fermions. At this point notice that the massive deformation couples R^α with Q^α ; that is the reason we have used the same index α for both, even though they had different kinds of indices in our treatment of funnels in the pure ABJM theory. This coupling thus breaks the $SO(6)$ invariance. Nevertheless, when writing down the full scalar potential the terms that couple R^α and Q^α vanish [33]. As a result, we will keep the different notation in their respective indices that we used in the previous section with R^α and $Q^{\dot{\alpha}}$.

4.1 Matrix fluctuation expansion

We now proceed to compute the action for quadratic fluctuations around the solution $R^\alpha = fG^\alpha$. Satisfying the classical equations of motion sets $f^2 = \frac{\mu k}{2\pi}$. A general fluctuation of the fields is given by⁸

$$\begin{aligned} R^\alpha &= fG^\alpha + r^\alpha, & R_\alpha^\dagger &= fG_\alpha^\dagger + r_\alpha^\dagger \\ Q^{\dot{\alpha}} &= q^{\dot{\alpha}}, & Q_{\dot{\alpha}}^\dagger &= q_{\dot{\alpha}}^\dagger \\ A_\mu &= A_\mu, & \psi^{\dagger I} &= \psi^{\dagger I}. \end{aligned} \quad (4.4)$$

This solution preserves an $SU(2) \times SU(2) \times U(1)$ subgroup of the R-symmetry group and $\mathcal{N} = 6$ supersymmetry [32, 33]. We will call these two $SU(2)$ factors ‘geometric’ and ‘transverse’ respectively, as the former will account for the geometric symmetry of the emergent fuzzy S^2 by rotating the scalars R^α entering the solution, while the transverse acts on the $Q^{\dot{\alpha}}$, which are zero on the solution. The vector index I of $SU(4)$ decomposes into the $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ of $SU(2) \times SU(2)$. This is clear since we have $C^I = (R^\alpha, Q^{\dot{\alpha}})$, as seen in Section 2.

In writing the fluctuation action, the symmetry operators that we will use along with the fields are

$$\begin{aligned} J_\beta^\alpha &= G^\alpha G_\beta^\dagger, & \hat{J}_\beta^\alpha &= J_\beta^\alpha - \frac{1}{2}\delta_\beta^\alpha J, & J &= J_\gamma^\gamma = N - 1, \\ \bar{J}_\beta^\alpha &= G_\beta^\dagger G^\alpha, & \hat{\bar{J}}_\beta^\alpha &= \bar{J}_\beta^\alpha - \frac{1}{2}\delta_\beta^\alpha \bar{J}, & \bar{J} &= \bar{J}_\gamma^\gamma = N - N\bar{E}_{11} \end{aligned} \quad (4.5)$$

and

$$\mathcal{L}_\beta^\alpha(q^{\dot{\gamma}}) = \hat{J}_\beta^\alpha q^{\dot{\gamma}} - q^{\dot{\gamma}} \hat{\bar{J}}_\beta^\alpha. \quad (4.6)$$

⁸For notational economy we use the same symbol for the gauge field and fermion fluctuations as for the classical fields. We hope that this will not cause confusion.

The \mathcal{L}_β^α give the action of the geometric $SU(2)$ on the field q^α , which will become a derivative (translation operator) in the classical limit. On the r_β^α and A_μ fluctuations, the same will be given by the adjoint action of

$$J_i = (\tilde{\sigma}_i)_\beta^\alpha J_\alpha^\beta, \quad J_\beta^\alpha = \frac{(N-1)}{2} \delta_\beta^\alpha + \frac{1}{2} J_i (\tilde{\sigma}_i)_\beta^\alpha. \quad (4.7)$$

Then $[J_i, \cdot]$ acts as a derivative (translation operator) on the 2-sphere in the classical limit, while at finite N it can be thought of as a ‘fuzzy derivative’ operator.

As we have already seen in (3.50) and (3.52), the scalar fluctuations r^α can be decomposed in terms of fields r, s_β^α and T^α . We will not consider T^α for the best part of this section, as we will see towards the end that it will decouple when we keep fluctuations up to quadratic order. Thus, we write

$$\begin{aligned} r^\alpha &= r G^\alpha + s_\beta^\alpha G^\beta \\ r_\alpha^\dagger &= G_\alpha^\dagger r + G_\beta^\dagger s_\alpha^\beta, \end{aligned} \quad (4.8)$$

where r and s_β^α have an expansion in the fuzzy spherical harmonics of $End(\mathbf{V}^+)$, given in terms of $1, GG^\dagger, \dots$. We have assumed that $r^\dagger = r$ and $s^\dagger = s$. Instead of s_β^α we will also find it more convenient to use the vector field s_i defined as

$$s_i = s_\alpha^\beta (\tilde{\sigma}_i)_\beta^\alpha, \quad s_\beta^\alpha = \frac{1}{2} s_i (\tilde{\sigma}_i)_\beta^\alpha. \quad (4.9)$$

This way we have traded the two complex fluctuations r^α with four real fields on S^2 . These will be like the fluctuations A_i of $X_i + A_i$ in usual matrix realisations of the fuzzy 2-sphere [23, 26]. From the s_i we will get a radial scalar $J_i s_i + s_i \bar{J}_i$ and a gauge field A_a on the sphere.

4.1.1 Fluctuations for the transverse scalars: sextic terms

The ABJM potential for the scalars is purely sextic. However, the mass-deformed theory of [33] or equivalently (as we will see shortly through the kinetic terms) the funnel solution, also have a quartic and a quadratic piece. We will evaluate these contributions separately, starting with the ABJM sextic potential $V_6 = \frac{4\pi^2}{3k^2} \hat{V}$, which is composed of 4 terms. We denote these by

$$\begin{aligned} \hat{V}_1 &= -\text{Tr}(C^I C_I^\dagger C^J C_J^\dagger C^K C_K^\dagger) \\ \hat{V}_2 &= -\text{Tr}(C_I^\dagger C^I C_J^\dagger C^J C_K^\dagger C^K) \\ \hat{V}_3 &= -4\text{Tr}(C^I C_J^\dagger C^K C_I^\dagger C^J C_K^\dagger) \\ \hat{V}_4 &= 6\text{Tr}(C^I C_J^\dagger C^J C_I^\dagger C^K C_K^\dagger). \end{aligned} \quad (4.10)$$

By plugging the transverse fluctuations ansatz into the above we find that the quadratic terms in

q^α give

$$\begin{aligned}
\hat{V}_1 &\rightarrow -3(N-1)^2 \text{Tr}(q^\beta q_\beta^\dagger) \\
\hat{V}_2 &\rightarrow -3N^2 \text{Tr}(q_\alpha^\dagger q^\alpha) + 3N^2 \text{Tr}(q_\alpha^\dagger q^\alpha \bar{E}_{11}) \\
\hat{V}_3 &\rightarrow 6\text{Tr}(q_\beta^\dagger J_\alpha^\gamma - \bar{J}_\alpha^\gamma q_\beta^\dagger)(J_\gamma^\alpha q^\beta - q^\beta \bar{J}_\gamma^\alpha) - 12N(N-1) \text{Tr}(q_\beta^\dagger q^\beta) \\
&\quad + 6N(N-1) \text{Tr}(q_\beta^\dagger q^\beta \bar{E}_{11}) \\
\hat{V}_4 &\rightarrow 18N(N-1) \text{Tr}(q^\alpha q_\alpha^\dagger) - 12N(N-1) \text{Tr}(q_\beta^\dagger q^\beta \bar{E}_{11}), \tag{4.11}
\end{aligned}$$

where we have made use of the definitions and identities in (4.5) as well as (3.20). Now, using the relation

$$6\text{Tr}(q_\beta^\dagger J_\alpha^\gamma - \bar{J}_\alpha^\gamma q_\beta^\dagger)(J_\gamma^\alpha q^\beta - q^\beta \bar{J}_\gamma^\alpha) = 6\text{Tr}\left(\mathcal{L}_\alpha^\gamma(q^\beta)^\dagger \mathcal{L}_\gamma^\alpha(q^\beta)\right) + 3\text{Tr}(q^\beta q_\beta^\dagger) + 3N(N-2) \text{Tr}(q_\beta^\dagger q^\beta \bar{E}_{11}) \tag{4.12}$$

one can easily see that the mass terms cancel and that the final answer for the transverse scalar fluctuations is

$$V_6^\perp = \frac{4\pi^2 f^4}{3k^2} 6\text{Tr}\left(\mathcal{L}_\alpha^\gamma(q^\beta)^\dagger \mathcal{L}_\gamma^\alpha(q^\beta)\right). \tag{4.13}$$

Thus we are getting the gradient (kinetic) terms of the q^α fields on the fuzzy S^2 . These 2 complex scalars are massless at this level, but we have an explicit mass term for the full C^I fields in the mass-deformed theory.

4.1.2 Fluctuations for the parallel scalars: sextic terms

Next we analyse the fluctuations r^α . The calculations are rather involved, so here we present only the final results. Some identities and intermediate steps are presented in Appendix A.

It is useful to organise the parallel fluctuations into ones separately involving only r^2 , s^2 and r - s . Restricting to r^2 terms coming from $r^\alpha \sim rG^\alpha$ one has

$$\begin{aligned}
\hat{V}_1 &\rightarrow -9(N-2) \text{Tr}([J_i, r][J_i, r]) - 3(N-1)(17N^2 - 18N - 15) \text{Tr}(r^2) \\
\hat{V}_2 &\rightarrow 12(N-1) \text{Tr}([J_i, r][J_i, r]) + N(N-1)(69N - 72) \text{Tr}(r^2) \\
\hat{V}_3 &\rightarrow -3(N-2) \text{Tr}([J_i, r][J_i, r]) - 12(N-1)(N^2 - 3N) \text{Tr}(r^2) \\
\hat{V}_4 &\rightarrow -3N \text{Tr}([J_i, r][J_i, r]) + 6N(N-1)(N-3) \text{Tr}(r^2), \tag{4.14}
\end{aligned}$$

where we have implemented the following relation

$$\text{Tr}\left(J_\beta^\alpha r J_\alpha^\beta r\right) = \frac{1}{2} \text{Tr}([J_i, r][J_i, r]) + N(N-1) \text{Tr}(r^2). \tag{4.15}$$

Therefore, the contribution from these terms is

$$V_6^{r^2} = -\frac{4\pi^2 f^4}{3k^2} [3(N-4) \text{Tr}([J_i, r][J_i, r]) - 45(N-1) \text{Tr}(r^2)]. \tag{4.16}$$

For the s^2 terms we obtain

$$\hat{V}_1 \rightarrow -\frac{3}{4}(N-1) \text{Tr}((s_i J_i s_j J_j + J_i s_i J_j s_j) - \frac{3}{2}(N-1) \text{Tr}(J_i s_i s_j J_j))$$

$$\begin{aligned}
& + \frac{3}{4} i\epsilon_{ijk} (N-1)^2 \text{Tr}(s_i J_j s_k) - \frac{3}{4} (N-1)^3 \text{Tr}(s_i s_i) \\
\hat{V}_2 \rightarrow & -\frac{3}{4} N(N-1)(N-4) \text{Tr}(s_i s_i) + \frac{3N}{4} (5N-4) i\epsilon_{ijk} \text{Tr}(s_i J_j s_k) \\
& - \frac{3}{4} N \text{Tr}(J_i s_i J_j s_j + s_i J_i s_j J_j) - \frac{3}{4} N \text{Tr}(s_i \square s_i) \\
\hat{V}_3 \rightarrow & -\frac{3}{2} (N-1)(2N^2 - 3N + 5) \text{Tr}(s_i s_i) - 6N \text{Tr}(s_i J_i J_j s_j) \\
& + \frac{3}{2} (3N^2 - 10N + 5) i\epsilon_{ijk} \text{Tr}(s_i J_j s_k) + \frac{3}{2} \text{Tr}(J_i s_i J_j s_j) \\
& + \frac{9}{4} \text{Tr}(s_i \square s_i) - \frac{3}{2} (2N-3) \text{Tr}(J_i s_i J_j s_j + s_i J_i s_j J_j) \\
& - \frac{9}{4} i\epsilon_{ijk} \text{Tr}((J_i s_j J_k)(s_l J_l + J_l s_l)) + \frac{3}{8} i\epsilon_{ijk} \text{Tr}(J_i s_j \square s_k) \\
\hat{V}_4 \rightarrow & \frac{3}{2} (N-1)^2 (3N-4) \text{Tr}(s_i s_i) + \frac{3}{2} (N-1) \text{Tr}(s_i \square s_i) \\
& + \frac{3}{2} (4N-3) \text{Tr}(J_i s_i J_j s_j + s_i J_i s_j J_j) - \frac{3}{2} (N-1)(7N-4) i\epsilon_{ijk} \text{Tr}(s_i J_j s_k) \\
& - \frac{3}{2} i\epsilon_{ijk} \text{Tr}((J_i s_j J_k)(J_l s_l + s_l J_l)) + 3(2N-1) \text{Tr}(s_i J_i J_j s_j) , \tag{4.17}
\end{aligned}$$

where we have defined the ‘fuzzy Laplacian’

$$\square s_k \equiv [J_p, [J_p, s_k]] . \tag{4.18}$$

Adding up the above contributions and using identities in (A.9) one arrives at

$$\begin{aligned}
V_6^{s^2} = & -\frac{4\pi^2 f^4}{3 k^2} \left[\frac{3}{4} (N-1) \text{Tr}(s_i s_i) - \frac{3}{4} (N+1) \text{Tr}([J_i, s_i]^2) + \frac{3}{4} (4N-1) i\epsilon_{ijk} \text{Tr}(s_i J_k s_k) \right. \\
& \left. - \frac{3}{4} N \text{Tr}(s_i \square s_i) + \frac{3}{8} i\epsilon_{ijk} \text{Tr}(J_i s_j \square s_k + s_i J_j \square s_k) \right] . \tag{4.19}
\end{aligned}$$

For the r - s fluctuations we get

$$\begin{aligned}
\hat{V}_1 \rightarrow & -\frac{15}{2} (N-1)^2 \text{Tr}(r(s \cdot J + J \cdot s)) \\
\hat{V}_2 \rightarrow & -\frac{3}{2} N(5N-4) \text{Tr}(r(s_i J_i + J_i s_i)) + 6N i\epsilon_{ijk} \text{Tr}(r(J_i s_j J_k)) \\
\hat{V}_3 \rightarrow & 3 \text{Tr}(r \square s_i J_i + r J_i \square s_i) - 6(5N^2 - 7N + 1) \text{Tr}(r(J_i s_i + s_i J_i)) \\
& + 12(N-1) i\epsilon_{ijk} \text{Tr}(r(J_i s_j J_k)) \\
\hat{V}_4 \rightarrow & 3 \text{Tr}((J_i r J_i)(s_j J_j + J_j s_j)) - 18(N-1) i\epsilon_{ijk} \text{Tr}(r(J_i s_j J_k)) \\
& + 21(N-1)(2N-1) \text{Tr}(r(s_i J_i + J_i s_i)) . \tag{4.20}
\end{aligned}$$

Collecting these terms and using identities in (A.10)

$$V_6^{r-s} = -\frac{4\pi^2 f^4}{3 k^2} \left[-\frac{9}{4} \text{Tr}(r(\square s_i J_i + J_i \square s_i)) + \frac{3}{4} \text{Tr}(r \square (J_i s_i + s_i J_i)) - \frac{9}{2} \text{Tr}(r(s_i J_i + J_i s_i)) \right] . \tag{4.21}$$

4.1.3 Quartic bosonic potential terms

At $Q^\alpha = 0$, the bosonic quartic term in the potential, linear in μ , for the massive deformation of [33] is

$$V_4 = \frac{8\pi\mu}{k} \text{Tr}(R^{[\alpha} R_\beta^\dagger R^{\beta]} R_{\alpha}^\dagger) . \tag{4.22}$$

Using that $J_\beta^\alpha J_\alpha^\beta = \frac{(N-1)}{2}$, we get for the r^2 terms

$$V_4^{r^2} = \frac{8\pi\mu f^2}{k} \left[\frac{1}{2} \text{Tr} ([J_i, r][J_i, r]) + 3(N-1) \text{Tr} (r^2) \right]. \quad (4.23)$$

Using the second identity (A.5) we get for the s^2 terms

$$V_4^{s^2} = \frac{8\pi\mu f^2}{k} \left[\frac{1}{8} \text{Tr} ([J_i, s_i]^2) + \frac{1}{8} \text{Tr} (s_i \square s_i) - \frac{1}{4} (N-1) \text{Tr} [s_i s_i] - \frac{1}{4} (2N-1) i \epsilon_{ijk} \text{Tr} (s_i J_j s_k) \right]. \quad (4.24)$$

The r - s terms are with the help of (A.4)

$$V_4^{r-s} = \frac{8\pi\mu f^2}{k} \left[-i \epsilon_{ijk} \text{Tr} (r J_i s_j J_k) + \frac{1}{2} \text{Tr} (r (s_i J_i + J_i s_i)) \right]. \quad (4.25)$$

There is *a priori* also a quartic term $V_4^{q^2}$ involving the q 's from (4.2)-(4.3), but after a short calculation one can check that it in fact vanishes.

4.1.4 Quadratic bosonic potential terms

From (4.3) the bosonic mass term is

$$V_2 = -\mu^2 \text{Tr} [R^\alpha R_\alpha^\dagger + Q^\dot{\alpha} Q_\dot{\alpha}^\dagger] \quad (4.26)$$

and can be easily evaluated on the solution. From it we obtain

$$\begin{aligned} V_2^{r-s-q} &= -\mu^2 \left[\frac{1}{4} (N-1) \text{Tr} (s_i s_i) + (N-1) \text{Tr} (r^2) + \frac{1}{2} \text{Tr} (r (s_i J_i + J_i s_i)) \right. \\ &\quad \left. - \frac{1}{4} i \epsilon_{ijk} \text{Tr} (s_i J_j s_k) + q^\dot{\alpha} q_\dot{\alpha}^\dagger \right]. \end{aligned} \quad (4.27)$$

4.1.5 The T^α fluctuations

Until now we have neglected the T^α terms. However, we have seen that the full expansion of the R^α scalars actually is

$$\begin{aligned} R^\alpha &= f G^\alpha + r G^\alpha + s_\beta^\alpha G^\beta + T^\alpha \\ R_\alpha^\dagger &= f G_\alpha^\dagger + G_\alpha^\dagger r + G_\beta^\dagger s_\alpha^\beta + T_\alpha^\dagger. \end{aligned} \quad (4.28)$$

Since these fluctuations are of order N , while the ones we have already consider of order N^2 , we assumed that they will not play a role in our large- N calculation. However, in order to show that they have been rightfully neglected let us look at some terms involving the T^α 's. A mass term $\text{Tr} R_\alpha^\dagger R^\alpha$ gives the new contribution

$$\begin{aligned} \mu^2 \text{Tr} (R_\alpha^\dagger R^\alpha) &\rightarrow \mu^2 \text{Tr} (T_\alpha^\dagger T^\alpha) \\ &= \mu^2 \sum_k t_k^\alpha (t_k^\alpha)^* \text{Tr} (\hat{E}_{k1} \hat{F}_{1k}) \\ &= \mu^2 \sum_k t_k^\alpha (t_k^\alpha)^*. \end{aligned} \quad (4.29)$$

Note that we do not get cross terms between T and the r, s at quadratic order in the fluctuations because

$$\begin{aligned} G^\alpha \hat{F}_{1K} &= 0 \\ \hat{E}_{K1} G_\alpha^\dagger &= 0, \end{aligned} \quad (4.30)$$

which follow from the previously established relations

$$\begin{aligned} G^\alpha |e_1^- \rangle &= 0 \\ \langle e_1^- | G_\alpha^\dagger &= 0. \end{aligned} \quad (4.31)$$

Following the above through for all possible contributions it is clear that the combinations that appear are

$$\begin{aligned} T^\alpha T_\beta^\dagger &= \sum_{k,l=1}^N t_k^\alpha (t_l^\beta)^* |e_k^+ \rangle \langle e_l^+ | \equiv K_\beta^\alpha \in \text{End}(\mathbf{V}^+) \\ T_\alpha^\dagger T^\beta &= \sum_k t_k^\alpha (t_k^\beta)^* |e_1^- \rangle \langle e_1^- | \sim \bar{E}_{11} \in \text{End}(V_1^-). \end{aligned} \quad (4.32)$$

However, because of equations (4.30) and (4.31) one finds that in any expression that involves G 's the $T_\alpha^\dagger T^\beta$ yields zero, much in the same fashion as similar terms also did in the r, s calculation. On the other hand $T^\alpha T_\beta^\dagger \equiv K_\beta^\alpha$ does not give zero and should admit an expansion in terms of fuzzy spherical harmonics. By evaluating all contributions to the scalar potential

$$V^T \rightarrow -\mu^2 [(N^2 - 8N + 6)\text{Tr}(K_\alpha^\alpha) + (N - 8)\text{Tr}(K_\beta^\alpha J_\alpha^\beta)] \quad (4.33)$$

and one explicitly finds no mixing between T and r, s at quadratic order in the fluctuating fields. Hence, the T^α fluctuation truly decouples from the rest of our discussion, as expected.

4.2 Fluctuations and bosonic field theory on the classical S^2

To summarise, the action of fluctuations on the fuzzy S^2 coming from the bosonic potential is

$$\begin{aligned} V &= V_6^{r^2} + V_6^{s^2} + V_6^\perp + V_6^{r-s} + V_2^{r-s-q} + V_4^{r-s} \\ &= -\frac{4\pi^2 f^4}{3 k^2} [3(N-4)\text{Tr}([J_i, r][J_i, r]) - 45(N-1)\text{Tr}(r^2)] \\ &\quad -\frac{4\pi^2 f^4}{3 k^2} \left[\frac{3}{4}(N-1)\text{Tr}(s_i s_i) - \frac{3}{4}(N+1)\text{Tr}([J_i, s_i]^2) + \frac{3}{4}(4N-1)i\epsilon_{ijk}\text{Tr}(s_i J_k s_k) \right. \\ &\quad \left. - \frac{3}{4}N\text{Tr}(s_i \square s_i) + \frac{3}{8}i\epsilon_{ijk}\text{Tr}(J_i s_j \square s_k + s_i J_j \square s_k) \right] + \frac{4\pi^2 f^4}{3k^2} 6\text{Tr}(\mathcal{L}_\alpha^\gamma (q^\beta)^\dagger \mathcal{L}_\gamma^\alpha (q^\beta)) \\ &\quad -\frac{4\pi^2 f^4}{3 k^2} \left[-\frac{9}{4}\text{Tr}(r(\square s_i J_i + J_i \square s_i)) + \frac{3}{4}\text{Tr}(r \square (J_i s_i + s_i J_i)) - \frac{9}{2}\text{Tr}(r(s_i J_i + J_i s_i)) \right] \\ &\quad -\mu^2 \left[\frac{1}{4}(N-1)\text{Tr}(s_i s_i) + (N-1)\text{Tr}(r^2) + \frac{1}{2}\text{Tr}(r(s_i J_i + J_i s_i)) - \frac{1}{4}i\epsilon_{ijk}\text{Tr}(s_i J_j s_k) \right. \\ &\quad \left. + q^{\dot{\alpha}} q_{\dot{\alpha}}^\dagger \right] + \frac{8\pi\mu f^2}{k} \left[-i\epsilon_{ijk}\text{Tr}(r J_i s_j J_k) + \frac{1}{2}\text{Tr}(r(s_i J_i + J_i s_i)) \right]. \end{aligned} \quad (4.34)$$

We now turn to the understanding of the classical (large N) limit, when the fuzzy S^2 becomes a commutative S^2 , in which the above expression simplifies considerably. This makes use of a correspondence between the fuzzy sphere matrix algebra and the algebra of functions on S^2 that is familiar from Matrix Theory and has seen a number of similar applications [23–26, 28–30].

4.2.1 Geometric decomposition of s_i

The fluctuations are fields on a fuzzy S^2 and at large N they become fields on S^2 . This S^2 is described as being embedded in \mathbb{R}^3 . We have already seen that the coordinates J_i have the property

$$\begin{aligned} J_i^2 &= (N^2 - 1) \\ [J_i, J_j] &= 2i\epsilon_{ijk}J_k . \end{aligned} \quad (4.35)$$

When we define $x_i = \frac{J_i}{\sqrt{N^2-1}}$ we get

$$\begin{aligned} x_i^2 &= 1 \\ [x_i, x_j] &= \frac{2i}{\sqrt{N^2-1}}\epsilon_{ijk}x_k . \end{aligned} \quad (4.36)$$

So these are the coordinates on the sphere that become commuting in the large N limit. Moreover, the appropriately normalised Tr becomes an integral over the unit S^2

$$\frac{1}{N}\text{Tr} \rightarrow \int d^2\sigma\sqrt{\hat{h}} , \quad (4.37)$$

where $\sigma^\alpha = (\theta, \phi)$. The adjoint action of J_i also has a well defined large N limit, which we can identify with the action of Killing vectors [23, 26]

$$[J_i, \cdot] = -2i\epsilon_{ijk}x_j\partial_k = -2iK_i = -2iK_i^a\partial_a . \quad (4.38)$$

The vector fields $K_i = \epsilon_{ijk}x_j\partial_k$ obey $x_iK_i = 0$. The index a above runs over (θ, ϕ) , and so we have an expansion in terms of partial derivatives of the angular coordinates on the sphere. In the following $e^{\theta\phi} = 1$, \hat{h}_{ab} is the metric on the unit sphere, which we use to raise and lower indices, and $\omega^{ab} = \epsilon^{ab}/\sqrt{\hat{h}}$ is the inverse of the symplectic form. For explicit expressions and identities we refer to Appendix A.2.

One can easily see that the ‘fuzzy Laplacian’ \square defined in (4.18) can be related to the geometric Laplacian $\hat{\square}$ on the sphere

$$\begin{aligned} \square = [J_i, [J_i, \cdot]] &= -4K_j^a\partial_a K_j^b\partial_b \\ &= -4\partial_a\partial^a - 4K_p^a(\partial_a K_p^b)\partial_b \\ &= -4\frac{1}{\sqrt{\hat{h}}}\partial_a(\sqrt{\hat{h}}\partial^a) \equiv -4\hat{\square} , \end{aligned} \quad (4.39)$$

with

$$\square x_i = -4\hat{\square}x_i = 8x_i . \quad (4.40)$$

We will then decompose the s_i as [23, 26]

$$s_i = K_i^a A_a + x_i\phi . \quad (4.41)$$

The s_i transform as vectors of the SO(3) rotational symmetry of the embedding \mathbb{R}^3 . The A_a transform as vectors of the SO(2) tangent space group of the embedded sphere. We are decomposing $3 = 2 + 1$ to get a vector and scalar of SO(2): The vector will give gauge fields on the S^2 and the scalar will correspond to radial motions.

4.2.2 Transverse scalars

Recall from (4.13) that we have

$$V_6^\perp = \frac{4\pi^2 f^4}{3k^2} 6\text{Tr} \left(\mathcal{L}_\alpha^\gamma (q^{\dot{\beta}})^\dagger \mathcal{L}_\gamma^\alpha (q^{\dot{\beta}}) \right), \quad (4.42)$$

which can be rewritten as

$$V_6^\perp = \mu^2 \text{Tr} [(q_\beta^\dagger J_i - \bar{J}_i q_\beta^\dagger)(J_i q^{\dot{\beta}} - q^{\dot{\beta}} \bar{J}_i)]. \quad (4.43)$$

The q^α are still bifundamentals but are essentially different from the bifundamental fields that we studied in Section 3.3, since they have a transverse spinor index α and no indices (scalars) on the S^2 . They can be decomposed in terms of spherical harmonics as

$$q^\alpha = \sum_{l=0}^{N-1} (Q_\alpha^\dagger)_{lm} Y_{lm}(J_i) G^\alpha = Q_\alpha^\dagger G^\alpha, \quad (4.44)$$

where in the last equality we have transformed to a representation in terms of commuting spinors on S^2 , Q_α^\dagger , decomposed in usual spherical harmonics, and a G^α .⁹

In terms of the scalar representation q^α , we can obtain an expression that takes the form of the usual massless scalar action. To do this, we must first extend the definition of the usual angular derivatives ∂_a which act on x_i , to act on a larger algebra including G^α . This will allow the definition of $\partial_a q^\alpha$. We start by defining the adjoint action of K_i on G^α as

$$-2iK_i^a \partial_a(G^\alpha) = -2iK_i(G^\alpha) \equiv N(x_i G^\alpha - G^\alpha \bar{x}_i) = [(\tilde{\sigma}_i)_\beta^\alpha - x_i \delta_\beta^\alpha] G^\beta. \quad (4.45)$$

The algebra generated by $x_i, G^\alpha, G_\alpha^\dagger$ is constrained by $x_i = \frac{1}{N}(\tilde{\sigma}_j)_\alpha^\beta G^\alpha G_\beta^\dagger$ and one can check that the consistency condition

$$K_i(x_j) = K_i \left(\frac{1}{N}(\tilde{\sigma}_j)_\alpha^\beta G^\alpha G_\beta^\dagger \right) = \epsilon_{ijk} x_k \quad (4.46)$$

is satisfied, as expected. In particular, using the projector

$$K_i^a h_{ab} K_j^b = \delta_{ij} - x_i x_j \quad (4.47)$$

we find

$$\partial_a(G^\alpha) = \frac{1}{-2i} h_{ab} K_i^b (\tilde{\sigma}_i)_\beta^\alpha G^\beta. \quad (4.48)$$

With these definitions, the operator whose square appears in (4.43) becomes in the large N limit

$$J_i q^{\dot{\alpha}} - q^{\dot{\alpha}} \bar{J}_i = -2iK_i(q^{\dot{\alpha}}) + x_i q^{\dot{\alpha}}. \quad (4.49)$$

The sextic term then gives

$$V_6^\perp \rightarrow 4\mu^2 N \int d^2\sigma \sqrt{\hat{h}} \left[h^{ab} \partial_a q_\alpha^\dagger \partial_b q^{\dot{\alpha}} + \frac{1}{4} q_\alpha^\dagger q^{\dot{\alpha}} \right]. \quad (4.50)$$

⁹We will see in Section 7 that G^α is the matrix-version of a ‘twistor-like’ coordinate for the classical Hopf fibration of S^3 over S^2 .

The quadratic part of the potential cancels the above mass term, since

$$V_2^\perp \rightarrow -\mu^2 N \int d^2\sigma \sqrt{\hat{h}} q_\alpha^\dagger q^{\dot{\alpha}}, \quad (4.51)$$

so that the full quadratic action for the scalars is just the usual massless term

$$V^\perp \rightarrow 4\mu^2 N \int d^2\sigma \sqrt{\hat{h}} h^{ab} \partial_a q_\alpha^\dagger \partial_b q^{\dot{\alpha}}. \quad (4.52)$$

Although the fields $q^{\dot{\alpha}}$ contain the elements G^α of an extended algebra, the Lagrangian above contains bilinears which can be expressed in terms of x_i . Hence the action obtained as an integral over angles on the 2-sphere is well-defined.

In order to have fields depending only on the usual Cartesian sphere coordinates x_i , one needs to go to the commuting spinor representation using (4.44). At large N

$$J_i Q_\alpha^{\dot{\alpha}} G^\alpha - Q_\alpha^{\dot{\alpha}} G^\alpha \bar{J}_i = [J_i, Q_\alpha^{\dot{\alpha}}] G^\alpha + (\tilde{\sigma}_i)_\gamma^\alpha Q_\alpha^{\dot{\alpha}} G^\gamma, \quad (4.53)$$

which is equivalent to (4.49) and implies

$$\begin{aligned} J_i Q_\alpha^{\dot{\alpha}} G^\alpha - Q_\alpha^{\dot{\alpha}} G^\alpha \bar{J}_i &= -2i K_i^a (\partial_a Q_\alpha^{\dot{\alpha}}) G^\alpha + (\tilde{\sigma}_i)_\gamma^\alpha Q_\alpha^{\dot{\alpha}} G^\gamma \\ &= -2i K_i^a ((\nabla_a)_\gamma^\alpha Q_\alpha^{\dot{\alpha}}) G^\gamma + x_i Q_\alpha^{\dot{\alpha}} G^\alpha, \end{aligned} \quad (4.54)$$

with the definition of ∇ given by

$$(\nabla_a)_\gamma^\alpha = \delta_\gamma^\alpha \partial_a + \frac{1}{-2i} h_{ab} K_j^b (\tilde{\sigma}_j)_\gamma^\alpha. \quad (4.55)$$

In these variables the sextic action for the transverse scalars then becomes

$$\begin{aligned} V_6^\perp &\rightarrow 4\mu^2 N \int d^2\sigma \sqrt{\hat{h}} \left[\hat{h}^{ab} ((\nabla_a)_\gamma^\alpha Q_\alpha^{\dot{\alpha}}) J_\beta^\gamma ((\nabla_b)_\mu^\beta Q_\alpha^\mu) + \frac{1}{4} Q_\alpha^{\dot{\alpha}} J_\beta^\alpha Q_\alpha^\beta \right] \\ &= 2\mu^2 N^2 \int d^2\sigma \sqrt{\hat{h}} \left[\hat{h}^{ab} ((\nabla_a)_\gamma^\alpha Q_\alpha^{\dot{\alpha}}) (\delta_\beta^\gamma + x_i (\tilde{\sigma}_i)_\beta^\gamma) ((\nabla_b)_\mu^\beta Q_\alpha^\mu) + \frac{1}{4} Q_\alpha^{\dot{\alpha}} (\delta_\beta^\alpha + x_i (\tilde{\sigma}_i)_\beta^\alpha) Q_\alpha^\beta \right]. \end{aligned} \quad (4.56)$$

The spinor transformation property of $Q_\alpha^{\dot{\alpha}}$ under the geometric $SU(2)$ is interesting and we will comment on it in Section 8. The fact that the large N action for $Q_\alpha^{\dot{\alpha}}$ can be written compactly, in this large k semiclassical analysis, in terms of $q^{\dot{\alpha}}$ (which contains the G^α) will be interpreted in Section 7 in terms of a reduction of the Hopf fibre S^1 of S^3 to a \mathbb{Z}_2 .

4.2.3 Parallel scalars: The sextic s^2 terms

From (4.19) we have for the sextic s^2 terms

$$\begin{aligned} V_6^{s^2} &= -\frac{4\pi^2}{3} \frac{f^4}{k^2} \left[\frac{3}{4} (N-1) \text{Tr}(s_i s_i) - \frac{3}{4} (N+1) \text{Tr}([J_i, s_i]^2) + \frac{3}{4} (4N-1) i\epsilon_{ijk} \text{Tr}(s_i J_k s_k) \right. \\ &\quad \left. - \frac{3}{4} N \text{Tr}(s_i \square s_i) + \frac{3}{8} i\epsilon_{ijk} \text{Tr}(J_i s_j \square s_k + s_i J_j \square s_k) \right]. \end{aligned} \quad (4.57)$$

The term $\text{Tr}(s_i s_i)$ contains mass terms for the gauge field and the radial scalar. However, there are also other hidden mass terms that could cancel the former. In fact it would be surprising if the gauge field had a mass. Note that $\text{Tr}(s_i s_i)$ gives a mass to the *diagonal* of the $U(N) \times U(\bar{N})$ gauge field, which needs to cancel if the story is to be similar to [7]. It was observed in [26] that there can be highly nontrivial cancellations of mass terms for gauge fields, some of which only become apparent after using the equations of motion.

Upon converting the terms using (4.41) we find the following in the large N limit: For the ijk -antisymmetric term

$$i\epsilon_{ijk}\text{Tr}(J_i s_j s_k) \rightarrow N \int d^2\sigma \sqrt{\hat{h}} (-\hat{h}^{ab} A_a A_b + \omega^{ab} \phi F_{ab} - 2\phi^2). \quad (4.58)$$

For $s_i s_i$ one easily gets

$$\text{Tr}(s_i s_i) \rightarrow N \int d^2\sigma \sqrt{\hat{h}} (\hat{h}^{ab} A_a A_b + \phi^2). \quad (4.59)$$

We also have an extra term of the type $i\epsilon_{ijk}(J_i s_j \square s_k + s_i J_j \square s_k)$ at leading order in N . However this will yield a zero result since

$$i\epsilon_{ijk}(J_i s_j \square s_k + s_i J_j \square s_k) \sim N i\epsilon_{ijk}(x_i K_j^a A_a + K_i^a A_a x_j) \hat{\square} s_k = 0, \quad (4.60)$$

by relabelling and since the K 's and x 's commute. We are left with the $s_i \square s_i$ and $[J_i, s_i]^2$ terms. One could evaluate these directly. However, it is simpler to use the following alternative method:

First compute $[J_i, s_j] - [J_j, s_i]$

$$\begin{aligned} \mathcal{F}_{ij} &\equiv [J_i, s_j] - [J_j, s_i] \\ &= 2i\epsilon_{ijk} K_k^a A_a - 2iK_i^a K_j^b F_{ab} + 4i\epsilon_{ijk} x_k \phi + 2i(x_i K_j(\phi) - x_j K_i(\phi)). \end{aligned} \quad (4.61)$$

Now observe that

$$\begin{aligned} \text{Tr}(\mathcal{F}_{ij} \mathcal{F}_{ij}) &= 2\text{Tr}[J_i, s_j][J_i, s_j] - 2\text{Tr}[J_i, s_j][J_j, s_i] \\ &= -2\text{Tr}s_j \square s_j - 2\text{Tr}[J_i, s_i]^2 + 2i\epsilon_{ijk} \text{Tr}s_j [J_k, s_i], \end{aligned} \quad (4.62)$$

where in the above we have also made use of the Jacobi identity. So we see that the combination $\text{Tr}s_j \square s_j + \text{Tr}[J_i, s_i]^2$ that we would like to evaluate can be directly related to $\text{Tr}\mathcal{F}_{ij}^2$. Now using (4.58) and (4.61) we can easily calculate $\text{Tr}(\mathcal{F}_{ij} \mathcal{F}_{ij})$

$$\text{Tr}(\mathcal{F}_{ij} \mathcal{F}_{ij}) \rightarrow N \int d^2\sigma \sqrt{\hat{h}} \left(-8A_a A^a - 4F_{ab} F^{ab} - 32\phi^2 - 8\partial_a \phi \partial^a \phi + 24\omega^{ab} F_{ab} \phi \right). \quad (4.63)$$

Combining all of the above and using (4.39), we find that the s^2 -type terms give, in the large N limit

$$V_6^{s^2} \rightarrow \frac{4\pi^2}{3} \frac{f^4}{k^2} N \int d^2\sigma \sqrt{\hat{h}} \left(3N\phi \hat{\square} \phi - \frac{3}{2} N F^{ab} F_{ab} + 9N\omega^{ab} \phi F_{ab} - \frac{45}{4} N \phi^2 - \frac{9}{4} N A^a A_a + \mathcal{O}(1) \right). \quad (4.64)$$

4.2.4 Parallel scalars: The sextic r^2 and r - s terms

We follow similar steps for these terms. Note that

$$\text{Tr}[J_i, r][J_i, r] \rightarrow -N \int d^2\sigma \sqrt{\hat{h}} (\partial^a \tilde{r} \partial_a \tilde{r}), \quad (4.65)$$

where $\tilde{r} = 2r$,

$$J_i s_i + s_i J_i \rightarrow 2\sqrt{N^2 - 1} \phi + \mathcal{O}(1) = 2N\phi + \mathcal{O}(1) \quad (4.66)$$

and

$$\begin{aligned} i\epsilon_{ijk} \text{Tr} [r(J_i s_j J_k)] &= \frac{1}{4} r [\square(s_i J_i) - (\square s_i) J_i] = \frac{1}{4} r [\square(J_i s_i) - J_i (\square s_i)] \\ &\rightarrow N^2 \int d^2\sigma \sqrt{\hat{h}} [-\tilde{r}\phi + \frac{1}{2}\omega^{ab} F_{ab} \tilde{r}]. \end{aligned} \quad (4.67)$$

Thus to leading order we obtain

$$V_6^{r^2, r-s} \rightarrow -\frac{4\pi^2}{3} \frac{f^4}{k^2} 12N^2 \int d^2\sigma \sqrt{\hat{h}} \left(\frac{1}{2} \tilde{r} \square \phi - \frac{3}{4} \tilde{r} \omega^{ab} F_{ab} - \frac{15}{8} \tilde{r} \phi - \frac{1}{4} \partial^a \tilde{r} \partial_a \tilde{r} - \frac{15}{16} \tilde{r}^2 \right) + \mathcal{O}(N). \quad (4.68)$$

Combining everything the sextic potential gives the following quadratic fluctuations at leading order

$$V_6^{r,s(\text{quad})} \rightarrow -6N^2 \frac{4\pi^2}{3} \frac{f^4}{k^2} \int d^2\sigma \sqrt{\hat{h}} \left(\frac{F_{ab}^2}{4} + \frac{[\partial_a(\tilde{r} + \phi)]^2}{2} - \frac{3}{2}(\tilde{r} + \phi)\omega^{ab} F_{ab} + \frac{3}{8} A^a A_a + \frac{15}{8}(\tilde{r} + \phi)^2 \right). \quad (4.69)$$

4.2.5 The quartic and quadratic terms

For the quartic terms we find

$$V_4 \rightarrow \frac{N^2}{4} \frac{8\pi\mu f^2}{k} \int d^2\sigma \sqrt{\hat{h}} \left(A^a A_a + 3(\tilde{r} + \phi)^2 - 2\omega^{ab} F_{ab}(\tilde{r} + \phi) \right), \quad (4.70)$$

where everything else is of order one.

The quadratic terms are

$$\begin{aligned} V_2 &\rightarrow -\mu^2 \frac{N^2}{4} \int d^2\sigma \sqrt{\hat{h}} (A^a A_a + (\tilde{r} + \phi)^2 + \frac{4}{N} q^{\dot{\alpha}} q_{\dot{\alpha}}^\dagger) \\ &= -\mu^2 \frac{N^2}{4} \int d^2\sigma \sqrt{\hat{h}} (A^a A_a + (\tilde{r} + \phi)^2 + 2Q_\alpha^\beta (\delta_\beta^\alpha + x_i (\tilde{\sigma}_i)_\beta^\alpha) Q_\beta^\alpha). \end{aligned} \quad (4.71)$$

4.2.6 Collecting all potential terms

We remind that the equations of motion require that we have $f^2 = \frac{k\mu}{2\pi}$. By taking this into account, the mass terms for the gauge fields cancel and we obtain the following simple final result

$$V^{\parallel}(r_\beta^\alpha) \rightarrow -2N^2 \mu^2 \int d^2\sigma \sqrt{\hat{h}} \left(\frac{1}{4} F_{ab}^2 + \frac{1}{2} (\partial_a(\tilde{r} + \phi))^2 + \frac{1}{2} (\tilde{r} + \phi)^2 - \frac{1}{2} \omega^{ab} F_{ab}(\tilde{r} + \phi) \right) \quad (4.72)$$

and

$$\begin{aligned} V^\perp(q^{\dot{\alpha}}) &\rightarrow 4\mu^2 N \int d^2\sigma \sqrt{\hat{h}} \hat{h}^{ab} \partial_a q_{\dot{\alpha}}^\dagger \partial_b q^{\dot{\alpha}} \\ &= 2\mu^2 N^2 \int d^2\sigma \sqrt{\hat{h}} \left[\hat{h}^{ab} ((\nabla_a)_\gamma Q_\alpha^{\dot{\alpha}}) (\delta_\beta^\gamma + x_i (\tilde{\sigma}_i)_\beta^\gamma) ((\nabla_b)_\mu Q_\alpha^\mu) \right]. \end{aligned} \quad (4.73)$$

4.2.7 The disappearance of the $\tilde{r} - \phi$ mode in the classical S^2

We notice that the above potential depends only on the $\tilde{r} + \phi$ combination. The fluctuation of the parallel scalars, expressed using the large N limit expansion of s_i , is

$$r^\alpha = rG^\alpha + s_\beta^\alpha G^\beta = rG^\alpha + \left(K_i^a A_a + \frac{J_i}{N} \phi \right) \frac{(\tilde{\sigma}_i)_\beta^\alpha}{2} G^\beta \quad (4.74)$$

and can be rewritten, using

$$\begin{aligned} \frac{J_i(\tilde{\sigma}_i)_\beta^\alpha}{2} G^\beta &= J_\beta^\alpha G^\beta - \frac{N-1}{2} G^\alpha \\ J_\beta^\alpha G^\beta = G^\alpha (G_\beta^\dagger G^\beta) &= G^\alpha N(1 - \bar{E}_{11}) \end{aligned} \quad (4.75)$$

and $G^\alpha \bar{E}_{11} = 0$, as

$$\begin{aligned} r^\alpha &= K_i^a A_a \frac{(\tilde{\sigma}_i)_\beta^\alpha}{2} G^\beta + rG^\alpha + \phi \frac{G^\alpha}{N} \left(\bar{J} - \frac{N-1}{2} \right) \\ &= K_i^a A_a \frac{(\tilde{\sigma}_i)_\beta^\alpha}{2} + rG^\alpha + \phi \frac{G^\alpha}{N} \left(\frac{N+1}{2} - N\bar{E}_{11} \right) \\ &\rightarrow K_i^a A_a \frac{(\tilde{\sigma}_i)_\beta^\alpha}{2} G^\beta + \frac{\tilde{r} + \phi}{2} G^\alpha. \end{aligned} \quad (4.76)$$

Thus at the classical level, there is no $\tilde{r} - \phi$ fluctuation at all, only $\tilde{r} + \phi$. This fact is not an accident: The disappearance of this mode is related to the fact that expanding around the ground-state solution triggers a form of the Higgs mechanism, which renders a linear combination of the CS gauge fields dynamical, as we will see next. In that context, the combination $\tilde{r} - \phi$ plays the role of a Goldstone boson.

A finite N version of the above calculation, keeping track of the commutators between $x_i = \frac{J_i}{\sqrt{N^2-1}}$ and ϕ , leads to

$$r^\alpha = K_i^a \mathcal{A}_a \frac{(\tilde{\sigma}_i)_\beta^\alpha}{2} G^\beta + \frac{\varphi}{2} G^\alpha, \quad (4.77)$$

where

$$\begin{aligned} \varphi &= \left(2r + \sqrt{\frac{N+1}{N-1}} \phi \right) = 2r + \phi \left(1 + \frac{1}{N} + \frac{1}{2N^2} + \dots \right) \\ \mathcal{A}_a &= A_a - i \frac{\partial_a \phi}{\sqrt{N^2-1}}. \end{aligned} \quad (4.78)$$

Note that the last line is a standard gauge transformation of A_a .

This indicates that our framework for the large N action can be extended to subleading orders in the $\frac{1}{N}$ expansion without a drastic change in field content. This is consistent with the philosophy of large N collective field theory [84], where classical field theories capture large N dynamics and $\frac{1}{N}$ interactions are described by ordinary field theoretic interaction terms. In the case at hand, the subleading corrections will be related to the geometry of the fuzzy S^2 . This is not to say that the extension to subleading orders is trivial. For example the fact that ϕ can be expressed as $\phi = s_i J_i \sim J_i s_i$ at large N is no longer true, since we have to take in to account $[J_i, s_i] \neq 0$.

4.3 The fuzzy funnel case

Up to now, we have focused on the theory around the fuzzy sphere solution of [33]. However, it is straightforward to see that the case of the fuzzy funnel solution of the undeformed ABJM theory is exactly the same as far as the action for fluctuations is concerned. Indeed, the Hamiltonian for the funnel solution with $Q^\alpha = 0$ and $R^\alpha \neq 0$ is given by [35]

$$H \propto \int dx_1 ds \operatorname{Tr} \left| \dot{R}^\alpha - \frac{2\pi}{k} R^{[\alpha} R_\beta^\dagger R^{\beta]} \right|^2 + \text{topological term} , \quad (4.79)$$

where s is the M2 worldvolume direction extending away from the M5-brane in the M2 \perp M5 funnel and a dot denotes differentiation with respect to that coordinate. The BPS condition is solved by

$$R^\alpha = f(s)G^\alpha \quad \text{where} \quad f(s) = \sqrt{\frac{k}{4\pi s}} . \quad (4.80)$$

The background field plus fluctuations will be

$$R^\alpha = f(s) \left(G^\alpha + rG^\alpha + s_\beta^\alpha G^\beta \right) \equiv f(s)R_0^\alpha , \quad (4.81)$$

with R_0^α the fluctuating fuzzy sphere field.¹⁰ We also have

$$\dot{R}^\alpha = -\frac{2\pi}{k} f(s)^3 \left(G^\alpha + rG^\alpha + s_\beta^\alpha G^\beta \right) \equiv -\frac{2\pi}{k} f(s)^3 R_0^\alpha . \quad (4.82)$$

Then the Hamiltonian picks up a common prefactor that we can ignore, since it will just normalise the overall coefficient

$$H \propto \frac{4\pi^2}{k^2} f(s)^6 \int dx_1 ds \operatorname{Tr} \left| R_0^\alpha - \frac{2\pi}{k} R_0^{[\alpha} R_0^\dagger R_0^{\beta]} \right|^2 + \text{topological term} \quad (4.83)$$

and after expanding the above the matrix form of every term can be exactly related back to the sextic, quartic and quadratic potential terms of [33].

Hence, there is no difference between the fuzzy funnel and the mass-deformed cases, except for the profile function being a function of s in the former, $f(s)$, while a function of the mass-deformation parameter μ in the latter, $f(\mu)$, and our calculation will go through essentially unchanged.

5 The CS-Higgs system and the Higgs mechanism

We now turn our attention to the fluctuations analysis of the CS-Higgs action in the first two lines of (4.1), that is

$$\int d^3x \left[\frac{k}{4\pi} \epsilon^{\mu\nu\rho} \operatorname{Tr} [A_\mu^{(1)} \partial_\nu A_\rho^{(1)} + \frac{2i}{3} A_\mu^{(1)} A_\nu^{(1)} A_\lambda^{(1)} - A_\mu^{(2)} \partial_\nu A_\rho^{(2)} - \frac{2i}{3} A_\mu^{(2)} A_\nu^{(2)} A_\lambda^{(2)}] - \operatorname{Tr} [D_\mu C_I^\dagger D^\mu C^I] \right] \quad (5.1)$$

¹⁰We could also have chosen to rescale the fluctuations r^α by $f(\mu)$ in the mass deformation calculation done up to now.

containing the gauge fields of the $U(N) \times U(\bar{N})$ group. The covariant derivative acts on the bifundamental fields as

$$D_\mu C^I = \partial_\mu C^I + iA_\mu^{(1)} C^I - iC^I A_\mu^{(2)}. \quad (5.2)$$

Hence we have for the fluctuations

$$-D_\mu C_1^\dagger D^\mu C^I = -(\partial_\mu r_\alpha^\dagger - ifG_\alpha^\dagger A_\mu^{(1)} + ifA_\mu^{(2)} G_\alpha^\dagger)(\partial_\mu r^\alpha + ifA_\mu^{(1)} G^\alpha - ifG^\alpha A_\mu^{(2)}). \quad (5.3)$$

In order to obtain a dynamical gauge field from the above CS expressions we will make use of a novel implementation of the Higgs mechanism, in accordance with [7]. In that reference, the Higgs mechanism was studied in the context of 3-algebra BLG theories.¹¹ The generalisation to the ABJM model is straightforward and was carried out in [86, 87]. The mechanism involves a linear combination involving the difference of the gauge fields becoming massive, as in the usual Higgs mechanism, due to the coupling between gauge fields and scalars. However, in the absence of a kinetic term said combination can be integrated out rendering the other linear combination dynamical. The surviving gauge field transforms in the adjoint of the diagonal subgroup of $A_\mu^{(1)}$ and $A_\mu^{(2)}$. The bifundamental scalars also get promoted to adjoint fields.

In the approach of [7] a Higgs vev was developed by a single scalar and was proportional to the identity matrix. In our case however, all the R^α 's get a background value, which is proportional to G^α . The parallel with [7] is obtained by remembering the observation made in Section 3, that in the large N limit both the fuzzy spherical harmonics $Y_{lm}(J_i)$, coming from $U(N)$, and the fuzzy spherical harmonics $Y_{lm}(\bar{J}_i)$, coming from $U(\bar{N})$, give rise to two $U(1)$ gauge fields. For these fields we will perform a Higgsing procedure analogous to [7], so that only the diagonal $U(1)$ gauge field will remain. In this limit, the Higgs vev will effectively act as a $U(1)$ vev.

At large N the operators J_i and \bar{J}_i become the classical sphere coordinates

$$\sum_i x_i^2 = \sum_i \bar{x}_i^2 = 1, \quad (5.4)$$

with $x_i = \frac{J_i}{\sqrt{N^2-1}}$ and $\bar{x}_i = \frac{\bar{J}_i}{\sqrt{N^2-2N}}$. Then to leading order in N

$$J_i \rightarrow Nx_i, \quad \bar{J}_i \rightarrow N\bar{x}_i, \quad (5.5)$$

but we will actually also have to consider $\frac{1}{N}$ corrections in the definitions of x_i, \bar{x}_i for our derivation of the large N action, as we will need to understand subleading orders in the fields for a subtle term. For the spherical harmonics

$$\begin{aligned} Y_{lm}(J_i) &\rightarrow Y_{lm}(x_i) = (N^2 - 1)^{\frac{l}{2}} \sum_i f_{lm}^{(i_1 \dots i_l)} x_{i_1} \dots x_{i_l} \\ Y_{lm}(\bar{J}_i) &\rightarrow Y_{lm}(\bar{x}_i) = (N^2 - 2N)^{\frac{l}{2}} \sum_i f_{lm}^{(i_1 \dots i_l)} \bar{x}_{i_1} \dots \bar{x}_{i_l}. \end{aligned} \quad (5.6)$$

We remind that the CS gauge fields $A_\mu^{(1)}$ are functions of the fuzzy sphere coordinates J_i , whereas

¹¹See also [85] for an exposition in terms of the bifundamental notation of [9].

$A_\mu^{(2)}$ are functions of \bar{J}_i . It will also be useful for us to define fields with J_i and \bar{J}_i interchanged

$$\begin{aligned} A_\mu^{(1)} &= \sum_{l=0}^{N-1} a_\mu^{lm} Y_{lm}(J_i) \quad , \quad \hat{A}_\mu^{(1)} = \sum_{l=0}^{N-1} a_\mu^{lm} Y_{lm}(\bar{J}_i) \\ A_\mu^{(2)} &= \sum_{l=0}^{N-2} \bar{a}_\mu^{lm} Y_{lm}(\bar{J}_i) \quad , \quad \hat{A}_\mu^{(2)} = \sum_{l=0}^{N-2} \bar{a}_\mu^{lm} Y_{lm}(J_i) \quad , \end{aligned} \quad (5.7)$$

We can view $A_\mu^{(1)} \rightarrow \hat{A}_\mu^{(1)}$ as a map from $End(\mathbf{V}^+)$ to $End(\mathbf{V}^-)$, which preserves the $SU(2)$ transformation properties. Likewise $A_\mu^{(2)} \rightarrow \hat{A}_\mu^{(2)}$ is a map from $End(\mathbf{V}^-)$ to $End(\mathbf{V}^+)$. In the classical limit, the four fields above are determined by the degrees of freedom $a_\mu^{lm}, \bar{a}_\mu^{lm}$ of two classical fields on S^2 . We will define such classical fields $\mathbf{A}^{(i)}$ in the course of the derivation.

5.1 Higgs mechanism

As a first step in the analysis, we will look at the Higgs mechanism for the gauge field components in the $\mu = 0, 1, 2$ membrane worldvolume directions. In the presence of the background scalar field $C^I = (fG^\alpha, 0)$, the CS action for the two $U(1)$ fields on the classical (large N) 2-sphere becomes a Yang-Mills action for a single $U(1)$ dynamical gauge field.

In more detail, the pure gauge ($A_\mu^{(i)}$) terms in (5.3) give

$$-f^2 \text{Tr} [JA^{(1)\mu} A_\mu^{(1)}] - f^2 \text{Tr} [\bar{J}A^{(2)\mu} A_\mu^{(2)}] + 2f^2 \text{Tr} [A^{(1)\mu} G^\alpha A_\mu^{(2)} G_\alpha^\dagger] . \quad (5.8)$$

The first trace is over \mathbf{V}^+ and equals

$$\begin{aligned} -f^2 \text{Tr} [JA^{(1)\mu} A_\mu^{(1)}] &= -f^2 (N-1) \text{Tr}_{\mathbf{V}^+} [A^{(1)\mu} A_\mu^{(1)}] \\ &\simeq -f^2 N \text{Tr}_{\mathbf{V}^+} [A^{(1)\mu} A_\mu^{(1)}] . \end{aligned} \quad (5.9)$$

The second trace is over \mathbf{V}^- , but can be mapped using (5.7) and can be written as

$$\begin{aligned} -f^2 \text{Tr} [\bar{J}A^{(2)\mu} A_\mu^{(2)}] &= -f^2 N \text{Tr}_{\mathbf{V}^-} [A^{(2)\mu} A_\mu^{(2)}] \\ &\simeq -f^2 N \text{Tr}_{\mathbf{V}^+} [\hat{A}^{(2)\mu} \hat{A}_\mu^{(2)}] . \end{aligned} \quad (5.10)$$

Using $\text{Tr}_{\mathbf{V}^+} \rightarrow N \int d^2\sigma \sqrt{\hat{h}}$ and $A^{(1)} \rightarrow \mathbf{A}^{(1)}$ along with $\hat{A}^{(2)} \rightarrow \mathbf{A}^{(2)}$ we have, always in the strict large N limit,¹²

$$-N^2 f^2 \int d^2\sigma \sqrt{\hat{h}} (\mathbf{A}^{(1)\mu} \mathbf{A}_\mu^{(1)} + \mathbf{A}^{(2)\mu} \mathbf{A}_\mu^{(2)}) . \quad (5.11)$$

For the third term in (5.8) we need to commute G^α past the fuzzy spherical harmonics. Using (3.34) one can show the following exact relation, which is valid for any finite value of N

$$G^\alpha Y_{lm}(\bar{J}_i) G_\alpha^\dagger = JY_{lm}(J_i) - lY_{lm}(J_i) . \quad (5.12)$$

¹²Note here that we could have instead expressed everything in terms of $\text{Tr}_{\mathbf{V}^-}$ and then used $\text{Tr}_{\mathbf{V}^-} \rightarrow N \int d^2\sigma \sqrt{\hat{h}}$, along with $A^{(2)} \rightarrow \mathbf{A}^{(1)}$, $\hat{A}^{(1)} \rightarrow \mathbf{A}^{(2)}$, to get the same large N action, as was the case for the matter fluctuations. This is a reflection of a \mathbb{Z}_2 symmetry inherent in the derivation of the classical sphere action.

This uses the fact that the expansion of the spherical harmonics involves symmetric and traceless combinations of the J_i 's in a crucial way.¹³ Being careful about applying the correct normalisation factors, the above becomes in terms of the coordinates on the sphere

$$\begin{aligned} G^\alpha Y_{lm}(\bar{x}_i) G_\alpha^\dagger &= \left(\frac{N^2 - 1}{N^2 - 2N} \right)^{\frac{1}{2}} \left((N - 1) Y_{lm}(x_i) - l Y_{lm}(x_i) \right) \\ &= (N - 1) Y_{lm}(x_i) - \frac{l(l + 1)}{2N} Y_{lm}(x_i) + \mathcal{O}\left(\frac{1}{N^2}\right). \end{aligned} \quad (5.13)$$

The reader might be puzzled about the fact that we have taken extra care to keep track of the subleading term in the above expression, given that we are interested in the large N limit. The reason for this will become apparent very soon, in Eq. (5.23). For the moment, note that

$$-l(l + 1) Y_{lm}(x_i) = \hat{\square} Y_{lm}(x_i). \quad (5.14)$$

As a result, one has that at large N the term in question becomes¹⁴

$$\begin{aligned} 2f^2 \text{Tr} [A^{(1)\mu} G^\alpha A_\mu^{(2)} G_\alpha^\dagger] &= 2f^2 N \text{Tr}_{\mathbf{V}^+} [A^{(1)} \hat{A}^{(2)\mu} + f^2 A_\mu^{(1)} \hat{\square} \hat{A}^{(2)\mu}] \\ &\rightarrow f^2 \int d^2\sigma \sqrt{\hat{h}} \left(2N^2 \mathbf{A}_\mu^{(1)} \mathbf{A}^{(2)\mu} + \mathbf{A}_\mu^{(1)} \hat{\square} \mathbf{A}^{(2)\mu} \right). \end{aligned} \quad (5.15)$$

The combined part of the scalar kinetic term involving only gauge fields gives

$$- \int d^3x \text{Tr} \left(D_\mu C_I^\dagger D^\mu C^I \right) \rightarrow f^2 N^2 \int d^3x d^2\sigma \sqrt{\hat{h}} \left(\frac{1}{N^2} \mathbf{A}_\mu^{(1)} \hat{\square} \mathbf{A}^{(2)\mu} - (\mathbf{A}_\mu^{(1)} - \mathbf{A}_\mu^{(2)})^2 \right). \quad (5.16)$$

We now form linear combinations of the two U(1) gauge fields on the classical S^2 ,

$$\begin{aligned} A_\mu &= \frac{1}{2} (\mathbf{A}_\mu^{(1)} + \mathbf{A}_\mu^{(2)}) \\ B_\mu &= \frac{1}{2} (\mathbf{A}_\mu^{(1)} - \mathbf{A}_\mu^{(2)}), \end{aligned} \quad (5.17)$$

so that (5.16) becomes

$$f^2 \int d^3x d^2\sigma \sqrt{\hat{h}} \left(A_\mu \hat{\square} A^\mu - B_\mu \hat{\square} B^\mu - 4N^2 B^\mu B_\mu \right). \quad (5.18)$$

Setting that aside, the Chern-Simons action for $A_\mu^{(1)}$ and $A_\mu^{(2)}$ gives in the classical limit

$$\begin{aligned} S_{\text{CS}} &= \int d^3x \frac{k}{4\pi} \epsilon^{\mu\nu\rho} \text{Tr} [A_\mu^{(1)} \partial_\nu A_\rho^{(1)} - A_\mu^{(2)} \partial_\nu A_\rho^{(2)}] \\ &\rightarrow N \frac{k}{2\pi} \int d^3x d^2\sigma \sqrt{\hat{h}} \left(\epsilon^{\mu\nu\rho} B_\mu F_{\nu\rho} \right), \end{aligned} \quad (5.19)$$

where $F_{\mu\nu}$ is the usual abelian field strength, defined as

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (5.20)$$

¹³We expand on the derivation of (5.12) in Appendix B.1.

¹⁴Once again, one could have chosen to express everything in terms of \mathbf{V}^- and \bar{x}_i instead to obtain the same action, as explained in Appendix B.2.

Thus the total action for A_μ and B_μ at leading N order is

$$\int d^3x d^2\sigma \sqrt{\hat{h}} \left(N \frac{k}{2\pi} \epsilon^{\mu\nu\rho} B_\mu F_{\nu\rho} - 4f^2 N^2 B_\mu B^\mu + f^2 A_\mu \hat{\square} A^\mu - f^2 B_\mu \hat{\square} B^\mu \right). \quad (5.21)$$

We now notice that B_μ is an auxiliary field, which can be eliminated through its equation of motion. The $B_\mu \hat{\square} B^\mu$ is subleading in N and can be dropped so as to get

$$B^\mu = \frac{1}{8f^2 N} \frac{k}{2\pi} \epsilon^{\mu\nu\lambda} F_{\nu\lambda}, \quad (5.22)$$

leading to the following expression, up to a total derivative

$$\int d^3x d^2\sigma \sqrt{\hat{h}} \left(-f^2 \partial^a A_\mu \partial_a A^\mu - \left(\frac{k}{2\pi} \right)^2 \frac{1}{8f^2} F^{\mu\nu} F_{\mu\nu} \right). \quad (5.23)$$

Note importantly that the first term came from the $\frac{1}{N}$ corrections in (5.13) and is a part of the gauge field strength on the sphere that otherwise would have been absent. The second term, coming from the manipulation of the CS part of the action, has become a Yang-Mills kinetic term. In 3d a Yang-Mills action contains one degree of freedom, whereas the Chern-Simons action contains none. The extra degree of freedom comes from the disappearance of a Goldstone boson as in [7]. As we have seen in Subsection 4.2.7 this is the scalar mode $\tilde{r} - \phi$.

5.2 Other contributions from the scalar kinetic term

Apart from (5.8), which contains only gauge fields, there are additional contributions coming from the scalar kinetic term (5.3)

$$\begin{aligned} -\text{Tr} \left(D_\mu C_I^\dagger D^\mu C^I \right) &\rightarrow \text{Tr} \left[-\partial_\mu q_\alpha^\dagger \partial^\mu q^\alpha - \frac{N-1}{4} \partial_\mu s_i \partial^\mu s_i - (\partial_\mu r) (\partial^\mu s_i) J_i + \frac{i}{4} \epsilon_{ijk} J_k \partial_\mu s_i \partial^\mu s_j \right] \\ &\quad -if \text{Tr} \left[\partial_\mu r_\alpha^\dagger (A_\mu^{(1)} G^\alpha - G^\alpha A_\mu^{(2)}) + \partial_\mu r^\alpha (A_\mu^{(2)} G_\alpha^\dagger - G_\alpha^\dagger A_\mu^{(1)}) \right]. \end{aligned} \quad (5.24)$$

After some algebra, the terms on the first line give in the large N classical limit

$$N \int d^3x d^2\sigma \sqrt{\hat{h}} \left[-\frac{N}{2} \partial^\mu Q_\alpha^\dagger \partial_\mu \bar{Q}_\beta^\alpha - \frac{N}{4} \partial_\mu (\phi + \tilde{r}) \partial^\mu (\phi + \tilde{r}) - \frac{N}{4} h^{ab} \partial_\mu A_a \partial^\mu A_b \right]. \quad (5.25)$$

The terms on the second line reduce to

$$\frac{if}{2} \text{Tr} \left(\partial_\mu s_i [J_i, A^{(1)\mu}] \right) \quad (5.26)$$

resulting in

$$f \int d^3x \text{Tr} \left(\partial_\mu A_a \partial^a (A^\mu + B^\mu) \right). \quad (5.27)$$

However, after substituting the leading result (5.22), one observes that the term coming from B_μ is also subleading in N and can be readily dropped from (5.27). This is a nice feature since, had it not done so, it would have led to a higher derivative interaction. Hence in the classical limit the second line yields

$$fN \int d^3x d^2\sigma \sqrt{\hat{h}} \left(\partial_\mu A^a \partial_a A^\mu \right). \quad (5.28)$$

6 Final action for fluctuations

We can now collect the above as well as all other bosonic terms for the action on the classical 2-sphere, to obtain the bosonic piece of the action for fluctuations

$$\begin{aligned}
S^B = \int d^3x d^2\sigma \sqrt{\hat{h}} & \left[- \left(\frac{k}{2\pi} \right)^2 \frac{1}{8f^2} F_{\mu\nu} F^{\mu\nu} - \frac{N^2 \mu^2}{2} F_{ab} F^{ab} - \frac{N^2}{4} \partial_\mu A^a \partial^\mu A_a + Nf \partial_\mu A_a \partial^a A_\mu \right. \\
& - f^2 \partial^a A_\mu \partial_a A^\mu + N^2 \mu^2 F_{ab} \omega^{ab} \Phi - \frac{N^2}{4} \partial_\mu \Phi \partial^\mu \Phi - N^2 \mu^2 \partial_a \Phi \partial^a \Phi - N^2 \mu^2 \Phi^2 \\
& \left. + 2N^2 \mu^2 \left((\nabla_a)^\alpha_\gamma Q_\alpha^\dot{\alpha} \right) (\delta_\beta^\gamma + x_i (\tilde{\sigma}_i)^\gamma_\beta) \left((\nabla^\beta)_\mu Q_\dot{\alpha}^\mu \right) - \frac{N^2}{2} \partial^\mu Q_\alpha^\dot{\alpha} (\delta_\beta^\alpha + x_i (\tilde{\sigma}_i)^\alpha_\beta) \partial_\mu Q_\dot{\alpha}^\beta \right], \tag{6.1}
\end{aligned}$$

where we have renamed $(\tilde{r} + \phi) = \Phi$. This action does not have the canonical form of a 5d theory but we will perform a set of rescalings for the fields, which will bring the right relative factors for the various terms.

In [35] it was observed that in order to convert the C^I kinetic term of the undeformed ABJM theory in (4.1)

$$S = - \int d^3x \text{Tr} (D_\mu C^I D^\mu C_I^\dagger) \tag{6.2}$$

to the physical form

$$S_{phys} = -T_2 \int d^3x (D_\mu X^I D^\mu X_I^\dagger), \tag{6.3}$$

where $T_2 = [l_p^3 (2\pi)^2]^{-1}$ is the membrane tension and X^I are spacetime coordinates with dimensions of length, one needs the redefinition $C^I = T_2^{1/2} X^I$. Note that the scalar fluctuations will also have dimensions of length in this language.

The classical solution $C^I = (R^\alpha, Q^\dot{\alpha}) = (fG^\alpha, 0)$ implies that the physical theory has a solution $X^I = (T_2^{-1/2} fG^\alpha, 0)$, hence $f \rightarrow T_2^{-1/2} f$. What this means in practice, is that all terms originating from the potential and scalar kinetic term will pick up a factor of T_2 apart from (5.28), which will pick up a $T_2^{1/2}$. The $F_{\mu\nu}^2$ and $(\partial_a A_\mu)^2$ contributions remain the same, since they came from terms which did not involve scalars. We can encode this into a ‘physical’ form of the action¹⁵

$$\begin{aligned}
S_{phys}^B = T_2 \int d^3x d^2\sigma \sqrt{\hat{h}} & \left[- \frac{k^2 T_2^{-1}}{32\pi^2 f^2} F_{\mu\nu} F^{\mu\nu} - \frac{N^2 \mu^2}{2} F_{ab} F^{ab} - \frac{N^2}{4} \partial_\mu A^a \partial^\mu A_a + \frac{Nf}{T_2^{1/2}} \partial_\mu A_a \partial^a A_\mu \right. \\
& - \frac{f^2}{T_2} \partial^a A_\mu \partial_a A^\mu + N^2 \mu^2 F_{ab} \omega^{ab} \Phi - \frac{N^2}{4} \partial_\mu \Phi \partial^\mu \Phi - N^2 \mu^2 \partial_a \Phi \partial^a \Phi \\
& - N^2 \mu^2 \Phi^2 + 2N^2 \mu^2 \left((\nabla_a)^\alpha_\gamma Q_\alpha^\dot{\alpha} \right) (\delta_\beta^\gamma + x_i (\tilde{\sigma}_i)^\gamma_\beta) \left((\nabla^\beta)_\mu Q_\dot{\alpha}^\mu \right) \\
& \left. - \frac{N^2}{2} \partial^\mu Q_\alpha^\dot{\alpha} (\delta_\beta^\alpha + x_i (\tilde{\sigma}_i)^\alpha_\beta) \partial_\mu Q_\dot{\alpha}^\beta \right] \tag{6.4}
\end{aligned}$$

¹⁵We keep the names for the various fields as before. We hope that this will not cause confusion.

and perform the following rescalings of the fields

$$A_\mu \rightarrow A_\mu (4\pi l_s) \frac{1}{T_2^{-1/2} f}, \quad A_a \rightarrow A_a (4\pi l_s) \frac{1}{N}, \quad \Phi \rightarrow \Phi (4\pi l_s) \frac{1}{N\mu}, \quad Q_\alpha^\beta \rightarrow Q_\alpha^\beta (4\pi l_s) \frac{1}{N\mu}. \quad (6.5)$$

We also rescale the metric on the sphere defining $h_{ab} = \mu^{-2} \hat{h}_{ab}$ so that $\sqrt{h} = \mu^{-2} \sqrt{\hat{h}}$, as well as the worldvolume coordinates $x^\mu \rightarrow \frac{1}{2} x^\mu$, leading to $\partial_\mu \rightarrow 2\partial_\mu$ and the scaling of the measure $d^3x \rightarrow 2^{-3} d^3x$. The action takes the form

$$\begin{aligned} S_{phys}^B = \frac{1}{g_s l_s} \int d^3x d^2\sigma \sqrt{h} & \left[-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{4} F_{ab} F^{ab} - \frac{1}{2} \partial^a A_\mu \partial_a A^\mu - \frac{1}{2} \partial_\mu A^a \partial^\mu A_a + \partial_\mu A_a \partial^a A^\mu \right. \\ & + \left((\nabla_a)_\gamma^\alpha Q_\alpha^\beta \right) (\delta_\beta^\gamma + x_i (\tilde{\sigma}_i)_\beta^\gamma) \left((\nabla^\alpha)_\mu^\beta Q_\alpha^\mu \right) - \partial_\mu Q_\alpha^\beta (\delta_\beta^\alpha + x_i (\tilde{\sigma}_i)_\beta^\alpha) \partial^\mu Q_\alpha^\beta \\ & \left. - \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \frac{1}{2} \partial_a \Phi \partial^a \Phi - \frac{\mu^2}{2} \Phi^2 + \frac{\mu}{2} \omega^{ab} F_{ab} \Phi \right]. \end{aligned} \quad (6.6)$$

Note that the rescalings that we have performed are precisely the ones needed to yield the correct N dependence for $(\partial_a A_\mu)^2$. This was the term that we got by keeping track of subleading corrections in (5.13). In fact, had it appeared at leading N order in that expression it would have dominated the whole action after the rescalings.

The above action resembles closely the canonical form of an abelian 5d Yang-Mills on a two-sphere, which is the low-energy limit of a D4-brane theory with $g_{YM}^2 = g_s l_s$. Using the index $A = \{\mu, a\}$ to include both the flat and angular variables, and the metric $g_{AB} = (\eta_{\mu\nu}, h_{ab})$ the action finally becomes

$$\begin{aligned} S_{phys}^B = \frac{1}{g_{YM}^2} \int d^3x d^2\sigma \sqrt{h} & \left[-\frac{1}{4} F_{AB} F^{AB} - \frac{1}{2} \partial_A \Phi \partial^A \Phi - \frac{\mu^2}{2} \Phi^2 - \partial_\mu Q_\alpha^\beta (\delta_\beta^\alpha + x_i (\tilde{\sigma}_i)_\beta^\alpha) \partial^\mu Q_\alpha^\beta \right. \\ & \left. + \left((\nabla_a)_\gamma^\alpha Q_\alpha^\beta \right) (\delta_\beta^\gamma + x_i (\tilde{\sigma}_i)_\beta^\gamma) \left((\nabla^\alpha)_\mu^\beta Q_\alpha^\mu \right) + \frac{\mu}{2} \omega^{ab} F_{ab} \Phi \right]. \end{aligned} \quad (6.7)$$

The size of the 2-sphere seen by the fluctuations is set by μ^{-1} .

Some comments are in order: The fact that the transverse scalars are naturally transformed to a representation in terms of commuting spinors is related to the issue of the exact interpretation of the final action in terms of branes and is a familiar feature of D-brane worldvolume theories on compact spaces. We elaborate on this in Section 8. We are also getting a mass term for Φ as well as an $F\Phi$ interaction term. This is a sign of the nontrivial geometry seen by the fluctuations.

7 M5 to D4 through Hopf

We have shown in the previous section that the analysis of fluctuations in the membrane theory leads to a $U(1)$ theory on $\mathbb{R}^{2,1} \times S^2$. This naturally suggests an interpretation as a D4-brane action. In this section, we will develop the D4-brane interpretation further. We start by looking at the energy of funnel solutions and show that they match the D4-brane interpretation just as well as the M5-brane interpretation [35]. We explain the reduction from M5 to D4 by using the Hopf fibration of S^3 over S^2 and the \mathbb{Z}_k quotient action along the S^1 fibre. We then show that the structure of the matrices of

[33] and in fact the field content of the ABJM theory can be anticipated by considering finite matrix constructions in fuzzy geometry inspired by the Hopf fibration. We extend these considerations to speculate on the problem of seeing the classical S^3 of M2-M5 systems in flat space from some appropriate large N membrane theory.

7.1 Physical radius and D4-brane energy for the funnel

In the case of the ground-state (zero energy) solutions for the massive deformation, where we have set up our calculation so far, we have seen that the form of the fluctuation action is a field theory on $\mathbb{R}^{2,1} \times S^2$. A simple calculation in favour of the S^2 D4-brane interpretation is also available in the energy of the funnel solution. Following the literature on brane polarisations, it is possible to define a physical radius at large N as

$$R_{\text{ph}}^2 = \frac{2}{N} \text{Tr} (X^I X_I^\dagger) = 8\pi^2 f^2 N l_p^3 = 2\pi N l_p^3 \frac{k}{s}, \quad (7.1)$$

where $f^2 = \frac{k}{4\pi s}$. The energy of the funnel can then be easily extracted from (4.79) and is given by

$$E = \frac{T_2^2}{2\pi} \int \frac{2\pi^2}{k} R_{\text{ph}}^3 dR_{\text{ph}} dx_1. \quad (7.2)$$

The authors of [35] interpreted the R_{ph}^3 as indicating a 5-brane wrapping an S^3/\mathbb{Z}_k in M-theory, with $T_5 = \frac{T_2^2}{2\pi}$ the M5-brane tension and $\frac{2\pi^2}{k}$ the volume of an S^3/\mathbb{Z}_k of unit radius ($V_{S^3} = 2\pi^2$ is divided in k units). However, the above formula is equally compatible with a 4-brane interpretation in Type IIA, which is in fact the one supported by the presence of the fuzzy S^2 obtained from our fluctuation analysis. Eq. (7.2) can be rewritten as

$$E = \frac{1}{g_s^2 l_s^6 (2\pi)^4} \int 4\pi \frac{R_{\text{ph}}^2}{4} \frac{R_{\text{ph}}}{k} dR_{\text{ph}} dx_1. \quad (7.3)$$

Now the ratio $\frac{R_{\text{ph}}}{k}$ is

$$\frac{R_{\text{ph}}}{k} = \sqrt{\frac{2\pi N l_p^3}{k s}} \quad (\text{funnel}). \quad (7.4)$$

For the ABJM theory to be weakly coupled one needs $\frac{N}{k} \rightarrow 0$, so that $k \rightarrow \infty$. Hence, with l_p, s fixed, the ratio $\frac{R_{\text{ph}}}{k} \rightarrow 0$. In this limit it is natural to go to a IIA picture, with

$$\frac{R_{\text{ph}}}{k} \equiv R_{11} = g_s l_s. \quad (7.5)$$

We can then write

$$\begin{aligned} E &= \frac{1}{g_s l_s^5 (2\pi)^4} \int 4\pi \left(\frac{R_{\text{ph}}}{2}\right)^2 dR_{\text{ph}} dx_1 \\ &= T_4 \int 4\pi \left(\frac{R_{\text{ph}}}{2}\right)^2 dR_{\text{ph}} dx_1, \end{aligned} \quad (7.6)$$

as is expected for a D4-brane in IIA involving an S^2 of radius $\frac{R_{\text{ph}}}{2}$ with the correct tension $T_4^{-1} = g_s l_s^5 (2\pi)^4$.

For the case of the mass-deformed theory we can similarly define the physical radius of the classical solution, which should correspond to the size of the brane in the D4 description. We have that $f(\mu)^2 = \frac{\mu k}{2\pi}$ and therefore

$$R_{\text{ph}}^2 = 4\pi N l_p^3 k \mu . \quad (7.7)$$

It is then also easy to obtain

$$R_{11} = \frac{R_{\text{ph}}}{k} = \sqrt{\frac{4\pi N l_p^3 \mu}{k}} \quad (\text{massive}) \quad (7.8)$$

and with this definition of the M-theory radius one can express (7.7) as

$$R_{\text{ph}}^2 = R_{11} 4\pi k N \mu l_s^2 = R_{\text{ph}} 2N \mu (2\pi l_s^2) . \quad (7.9)$$

Hence the radius of the two-sphere, evaluated according to the definition (7.1), is

$$\frac{R_{\text{ph}}}{2} = N \mu (2\pi l_s^2) . \quad (7.10)$$

Note that one would naturally expect R_{ph} to be the radius for the sphere that appears in the action for small fluctuations. However, we have explicitly seen that what actually appears in Eq.(6.7) is μ^{-1} . This is interesting and we will return to discuss it further in Section 8.

7.2 \mathbb{Z}_k reduction of S^3

We have just shown that there is a natural connection (7.5) between the radius of the proposed M-theory 3-sphere R_{ph} and the M-theory radius R_{11} . We now explain these relations by analysing the classical geometry of S^3/\mathbb{Z}_k . As a warm-up, we start by looking at the case of flat space solutions.

The moduli space of the $U(N) \times U(\bar{N})$ level k ABJM theory is the same as that for N M2-branes on $\mathbb{C}^4/\mathbb{Z}_k$ [14], namely $[(\mathbb{C}^4/\mathbb{Z}_k)^N]/S_N$. Here the action of \mathbb{Z}_k in the target space is

$$Z^i \rightarrow Z^i e^{\frac{2\pi i}{k}} \quad \text{with} \quad i = 1, \dots, 4 . \quad (7.11)$$

As argued above, in order to have a weakly coupled (perturbative) interpretation, we need to take $k \rightarrow \infty$. Let us first understand the case of $\mathbb{C}/\mathbb{Z}_{k \rightarrow \infty}$. For that example

$$Z \rightarrow Z e^{\frac{2\pi i}{k}} \simeq Z \left(1 + 2\pi i \frac{1}{k} + \dots \right) \simeq Z + 2\pi i \frac{Z}{k} . \quad (7.12)$$

Expanding around $Z = v + i0$, with $\frac{v}{k} \equiv r$ we have that

$$Z \rightarrow Z + 2\pi i r \quad (7.13)$$

is an invariance, or if $Z = X^1 + iX^2$ that X^2 is compactified with radius r .

We can do the same analysis for $\mathbb{C}^4/\mathbb{Z}_{k \rightarrow \infty}$, expanding around the background

$$Z^1 = v + i0 \quad \text{with} \quad Z^2 = Z^3 = Z^4 = 0 , \quad (7.14)$$

which means that if

$$Z^1 = X^1 + iX^2$$

$$Z^2 = X^3 + iX^4, \quad (7.15)$$

then X^2 is again compactified with radius r .

Now let us move on to the theory around the fuzzy sphere (or fuzzy funnel) solution, by imposing a 3-sphere constraint on the coordinates Z , for instance $|Z^1|^2 + |Z^2|^2 = R^2$. One obtains an S^3/\mathbb{Z}_k solution, as also considered in [35], since the action of \mathbb{Z}_k in (7.11) preserves this sphere constraint. We can then expand around the vacuum with $Z^1 = R + i0$, which means (exactly as above) that X^2 is compactified with radius $r = \frac{R}{k}$, and the fluctuations X^1, X^3, X^4 take us in the direction of an S^2 of radius $\frac{R}{2}$. Thus as $k \rightarrow \infty$, we have the spacetime result $S^3/\mathbb{Z}_k \rightarrow S^2_{R/2} \times S^1_{R/k}$, compatible with the worldvolume picture of the last section.

7.3 Hopf fibration and the classical limit of the fuzzy two-sphere

We next try to find a link between the classical spacetime and matrix (finite N) descriptions. The ground-state solution of [33] was $C^I = (R^\alpha, Q^\alpha)$ with $Q^\alpha = 0$ and

$$\begin{aligned} R^1 &= G^1 \\ R^2 &= G^2. \end{aligned} \quad (7.16)$$

In the classical interpretation of the fuzzy (matrix) coordinates C^I , we have

$$\begin{aligned} R^1 &= X^1 + iX^2 \\ R^2 &= X^3 + iX^4 \\ Q^1 &= X^5 + iX^6 \\ Q^2 &= X^7 + iX^8. \end{aligned} \quad (7.17)$$

We know that that the fuzzy coordinates satisfy $G^1 G_1^\dagger + G^2 G_2^\dagger = N - 1$ and $G^1 = G_1^\dagger$. The first one suggests a fuzzy 3-sphere, but the second is an extra constraint which reduces the geometry to a 2d one and we have indeed established that we have a fuzzy S^2 through the fluctuation analysis.

The construction of the fuzzy S^2 was obtained by

$$\begin{aligned} J_i &= (\tilde{\sigma}_i)_\beta^\alpha G^\beta G_\alpha^\dagger \\ x_i &= \frac{J_i}{\sqrt{N^2 - 1}} \Rightarrow \begin{cases} x_1 = \frac{J_1}{\sqrt{N^2 - 1}} = \frac{1}{\sqrt{N^2 - 1}} (G^1 G_2^\dagger + G^2 G_1^\dagger) \\ x_2 = \frac{J_2}{\sqrt{N^2 - 1}} = \frac{i}{\sqrt{N^2 - 1}} (G^1 G_2^\dagger - G^2 G_1^\dagger) \\ x_3 = \frac{J_3}{\sqrt{N^2 - 1}} = \frac{1}{\sqrt{N^2 - 1}} (G^1 G_1^\dagger - G^2 G_2^\dagger) \end{cases}. \end{aligned} \quad (7.18)$$

Following the standard Matrix Theory logic, the G^i should become coordinates in spacetime. On the other hand we are saying that bilinears in G^i are also spacetime coordinates. How can both be true? The answer is given by the classical Hopf map $S^3 \xrightarrow{\pi} S^2$.

The description of the Hopf map in classical geometry is precisely that we start with the Cartesian coordinates on the unit S^3 , X_1, X_2, X_3, X_4 , with

$$X_1^2 + X_2^2 + X_3^2 + X_4^2 = 1 \quad (7.19)$$

and then go to a set of ‘twistor-like’ variables $Z^1 = X_1 + iX_2$, $Z^2 = X_3 + iX_4$, which are related to the Cartesian coordinates on the unit S^2 by

$$x_i = (\tilde{\sigma}_i)_\beta^\alpha Z^\beta Z_\alpha^* , \quad (7.20)$$

to get

$$x_i x_i = (\tilde{\sigma}_i)_\beta^\alpha (\tilde{\sigma}_i)_\nu^\mu Z^\beta Z_\alpha^* Z^\nu Z_\mu^* = 1 . \quad (7.21)$$

The Z ’s obey $Z^\alpha Z_\alpha^* = 1$.

The various pieces of our geometric analysis are now falling into place: For an M5 wrapping the S^3 the relations between the radii, which we uncovered in a simple way from the spacetime picture, are in fact the ones prescribed by the full realisation of the three-sphere as the Hopf fibration $S^1 \hookrightarrow S^3 \xrightarrow{\pi} S^2$, with $R_{S^2} = \frac{R_{S^3}}{2}$ and the M-theory direction being identified with the S^1 fibre of radius $R_{S^1} = R_{S^3}$.¹⁶ The latter is a natural choice since this is the smallest circle that appears in the problem. For large R_{S^3} the orbifold $Z^i \sim e^{\frac{2\pi i}{k}} Z^i$ acts by shrinking the size of the fibre and hence the M-theory radius. In the limit of $k \rightarrow \infty$, R_{11} is very small and one is left with the S^2 base which supports a D4-brane in the IIA theory. This is consistent with the fact that our large N fluctuation action is an action for fields on S^2 .

Going in the other direction, one should start seeing the structure of the M-theory circle at finite k with the D4-brane becoming an M5-brane wrapping an $S^1/\mathbb{Z}_k \hookrightarrow S^3/\mathbb{Z}_k \xrightarrow{\pi} S^2$. However, as we have explicitly shown and we will discuss again in the next subsection, this is not something that can be captured by the large k semiclassical fluctuation analysis around the solution of [33]. Nevertheless, the calculation of physical quantities will be sensitive to the size of the S^1 , hence to k . Due to the high degree of supersymmetry, some quantities such as the energy (7.2) that we calculated on the classical solution at large k , give the correct finite- k form. Hence, they admit both the expected large- k Type IIA as well as a flat space ($k = 1$) M-theory interpretation. For general physical quantities one needs new finite- k methods to calculate properties of the the flat space M5-brane from the ABJM theory. In order to see the geometry of the extra circle more explicitly one would have to consider nonperturbative effects which we will come back to in Section 8.

7.4 Hindsight is 20/20 : ABJM structure from fuzzy Hopf

In retrospect all the key properties of $G^\alpha, G_\alpha^\dagger$ as summarised in Section 3 are determined by their interpretation as a fuzzy (matrix) realisation of the S^2 base of the Hopf fibration of S^3 . Indeed, suppose we want to lift the equations (7.20) from classical geometry to finite matrices. We need matrices for Z^α which we call G^α . Now the coordinates x_i transform in the spin-1 representation of $SU(2)$. If we want to build them from bilinears of the form $G^\dagger G$ we need G, G^\dagger to transform in the spin- $\frac{1}{2}$ representation. In the usual fuzzy 2-sphere, the x_i are operators mapping an irreducible N -dimensional $SU(2)$ representation V_N to itself. It is possible to do this in an $SU(2)$ -covariant fashion because the tensor product of spin-1 with V_N contains V_N . Since G^α are spin- $\frac{1}{2}$, and $\frac{1}{2} \otimes V_N = V_{N+1} \oplus V_{N-1}$ does not contain V_N , we need to work with reducible representations. The simplest thing to do would be to consider $V_N \oplus V_{N-1}$. The next simplest thing is to work with $V_N \oplus (V_{N-1} \oplus V_1)$. The latter possibility is chosen by the construction of [33] and allows a gauge group $U(N) \times U(N)$

¹⁶For a concise summary of the facts pertaining to the S^3 Hopf fibration see [88, 89].

which has a \mathbb{Z}_2 symmetry of exchange needed to preserve parity. To realise the $G^\alpha, G_\alpha^\dagger$ as solutions we need to set them equal to matter fields $R^\alpha, R_\alpha^\dagger$ which are in the bifundamental of the $U(N) \times U(N)$ gauge group.

So the rather curious property of the G matrices, notably the difference between the last of (3.4) and (3.8) which led directly to the fact that $\mathbf{V}^+ = V_N$ and $\mathbf{V}^- = V_{N-1} \oplus V_1$ follows from requiring a matrix realisation of the fuzzy S^2 base of the Hopf fibration. These in turn lead to the $SU(2)$ decompositions of $End(\mathbf{V}^+), End(\mathbf{V}^-), Hom(\mathbf{V}^+, \mathbf{V}^-), Hom(\mathbf{V}^-, \mathbf{V}^+)$, for the fluctuations and the Yang-Mills action. This novel realisation of fuzzy S^2 is thus intimately tied to the Hopf fibration. The usual fuzzy S^2 has also been discussed in terms of the Hopf fibration, where the realisation of the $SU(2)$ generators in terms of bilinears in Heisenberg algebra oscillators yields an infinite dimensional space which admits various projections to finite N constructions [90]. In that case the x_i are not bilinears in finite matrices.

Further light on the geometry can be shed by considering the expectation values of the operators x_i, G, G^\dagger . The situation for the x_i is familiar [21, 91]. If we choose the state of maximum spin, we see $\langle x_3 \rangle = 1$. If we choose states with lower spins we find $\langle x_1^2 + x_2^2 \rangle = 1 - \langle x_3^2 \rangle$. This gives a description of the fuzzy S^2 as a sequence of fuzzy circles fibering a discretised axis from N-pole to S-pole. The same manipulation can be done with the x_i in our construction to reveal the picture of a fuzzy S^2 at finite N .

Such an interpretation is also desirable for the G, G^\dagger , given the usual role of D-brane transverse coordinates as matrices. Here we need to consider $\langle -|G|+ \rangle, \langle +|G^\dagger|- \rangle$ to extract, at finite N , numbers which can be compared for example with $Z^1 = Z_1^*$. The x_i, G, G^\dagger are operators in $\mathbf{V}^+ \oplus \mathbf{V}^-$ which is isomorphic, as a vector space, to $\mathbf{V}_N \otimes V_2$. The endomorphisms of \mathbf{V}_N correspond to the fuzzy sphere. The N states of \mathbf{V}_N generalise the notion of points on S^2 to noncommutative geometry. The 2-dimensional space V_2 is invariant under the $SU(2)$. It is acted on by G, G^\dagger which have charge $+1, -1$ under the $U(1)$ (corresponding to (J, \bar{J})) acting on the fibre of the Hopf fibration. So in a sense the S^1 fibre has been reduced to a noncommutative geometry consisting of two points. It is interesting to note that this emerges naturally from our large- N fluctuation analysis: The field content of the theory we derived on S^2 can be organised using an extension of the algebra of the Cartesian coordinates x_i by G, G^\dagger subject to the relation $x_i = \frac{1}{N}(\tilde{\sigma}_i)^\beta_\alpha G^\alpha G^\dagger_\beta$. This was seen in Subsection 4.2.2 where the action for the transverse scalars in terms of bosonic spinors $Q_\alpha^\dot{\alpha}$ on the sphere was packaged elegantly using $q^{\dot{\alpha}} = Q_\alpha^\dot{\alpha} G^\alpha$ into (4.52).

We emphasise that the S^2 we constructed in matrix geometry from the ground-state solution really describes the noncommutative version of the base of the Hopf fibration in a finite N setting. In the large N limit it approaches the S^2 base of the standard Hopf fibration of S^3 . The fluctuation analysis does not have enough modes to describe the full space of functions on S^3 , even if we drop the requirement of $SO(4)$ covariance and allow for the possibility of an $SU(2) \times U(1)$ description. We elaborate on this below.

For $k \rightarrow \infty$, where the fluctuation action around the classical solutions of [33] is valid, the S^1/\mathbb{Z}_k fibre (parametrised by a coordinate y) becomes a very small circle. As we explained above, the only remnant of the circle in the matrix construction is the multiplicity associated with having states $|+\rangle, |-\rangle$ in \mathbf{V}^+ and \mathbf{V}^- . A classical description of the S^3 metric as a Hopf fibration contains a coordinate y transverse to the S^2 . However, the matrix fluctuations of the solution are mapped to

functions on S^2 and lead to a field theory on S^2 . This means we do not see the coordinate y as a worldvolume coordinate for the emergent D4-brane. We do not see it as a transverse coordinate either, which is as expected since in M-theory this is an M5-brane wrapping the y direction. Indeed, we have shown in Section 6 that only the mode $\phi + \tilde{r}$ (and not $\phi - \tilde{r} \equiv y$) appears in the action for fluctuations.

In sum, the action we have derived does not contain y neither as a transverse scalar nor as a worldvolume coordinate and it can be consistently interpreted according to Subsection 7.3 as the double-dimensional reduction of the M5-brane action along the \mathbb{Z}_k quotient of the S^3 Hopf fibre (with length $\frac{R}{k}$). This is a IIA reduction of M-theory on the same small circle.

7.5 Multi-membrane actions and S^3 at large N

One might ask what kind of multi-membrane action would contain a fuzzy S^3 solution and fluctuations which are appropriate for an $SU(2) \times U(1)$ covariant description of the S^3 . A dimensional reduction on the S^1 fibre is compatible with the symmetry. In such a description, there are fields on S^2 with extra index n , as expected for a simple S^1 reduction

$$\phi(x, y) = \sum_n \phi_n(x) e^{in2\pi y/R} . \quad (7.22)$$

The blow-up of a fuzzy S^2 into a fuzzy S^3 was analysed for instance in [81]. In that analysis, the fibre direction y is Fourier transformed by replacing the ∂_y with a semi-integer $-iq$. The vacuum solution in that case has fuzzy sphere solutions $L_i^{(j_s)}$ of various sizes j_s that are therefore labelled by an extra semi-integer index j_s , identified with q above. Then all the multiplicities N_s are taken to be equal, $N_s = N, \forall s$, with a \mathbb{Z}_N identification of the N blocks, where $N \rightarrow \infty$. In that fashion one obtains the fuzzy 3-sphere spherical harmonics from the fuzzy 2-sphere spherical harmonics with an extra index q . In our case, the absence of the extra index q signifies that the theory is dimensionally reduced to the $q = 0$ sector, or that we have a very small circle (as we argued, in the $k \rightarrow \infty$ limit). The extra fluctuations $\hat{E}_{k1}, \hat{F}_{1k}$ from Subsection 3.3 are uncharged under the $U(1)$ action of (J, \bar{J}) , so they do not provide the tower of charges corresponding to the S^1 .

As we have argued, the structure of the ABJM theory, with fields acting on $V_N \otimes V_2$ can be given a noncommutative geometry interpretation in terms of a discrete bundle over a fuzzy S^2 . We may speculate that a description capable of seeing more of the geometry of the Hopf fibration would involve a vector space $V_N \otimes V_K$, where V_K is a K -dimensional vector space. This would give a K -point approximation to the fibre and would suggest a gauge group $U(N)^{\otimes K}$, with $K \rightarrow \infty$ giving the classical geometry of the S^3 . This could presumably draw on an embedding of [81, 89] in an ABJM type membrane action. Another way to construct a theory that contains a classical S^3 , described in an $SU(2) \times U(1)$ covariant manner, might be to realise the standard Holstein-Primakoff construction with Heisenberg oscillators [90]. If we are to set fields in the theory to a, a^\dagger in a solution, we need a $U(\infty)$ theory. This could be viewed as a $K \rightarrow \infty$ version of a construction with $V_N \otimes V_K$. The question we would like to pose is whether an appropriate field theory with $U(\infty)$ gauge group and containing a realisation of S^3 as a Hopf fibration correctly describes the large N limit of membranes. A related question is to understand the relation of such a construction to the large N limit of ABJM. These questions may well lead to a new perspective on the solutions to the BLG fundamental identity

in terms of infinite dimensional algebras [92, 93].

8 Discussion and future avenues

There are a number of issues that emerge from our analysis and would need to be better understood:

- **The number of spherical branes**

Since we have derived a $U(1)$ theory in $\mathbb{R}^{2,1} \times S^2$ it is clear that the solutions of [33] describe a single D4-brane. We may obtain an extension to multiple D4-branes by considering reducible representations of the algebra satisfied by the G^α matrices. Such reducible representations are easy to construct (see appendix C). Some formal similarities between the current fuzzy S^2 and the $SO(4)$ -covariant fuzzy S^3 constructions become apparent in the proof. If one had knowledge of how to go from spherical (S^2) D4-branes to spherical (S^3) M5-branes by blowing up the Hopf fibre over S^2 , this would provide a starting point for studying *multiple* fivebranes in M-theory.

- **Regimes and parameters**

As we discussed, we need k large so that a semiclassical approach around classical solutions gives a reliable account of the physics, while large N is needed in order to see a classical geometry. Moreover, for the validity of the 2+1 dimensional Chern-Simons perturbation theory, we need its 't Hooft coupling $\lambda = \frac{N}{k}$ to be fixed and small. Finally, we also need $l_p \rightarrow 0$ for the validity of the low energy ABJM action (decoupling).

We have only analysed quadratic fluctuations, but the parameter $f^{-2} = \frac{2\pi}{\mu k}$ controls the terms in the action of higher order in the fields (*i.e.* cubic, quartic, *etc.*). Since k is large with μ fixed, the parameter is small. On the other hand, fuzzy and higher derivative corrections are of order $\frac{1}{\mu N}$, which is also small, as $\frac{N}{k}$ is fixed.

Note then that the limits considered here are further constrained with respect to the limits discussed in [14]. The issue of what limits can be relaxed or modified while keeping some of the features our analysis is one that deserves further study.

- **Dual brane descriptions : D2, D4, M5**

We have given strong evidence in Section 7 that the final action Eq.(6.7), derived from the M2-brane worldvolume, is that of a D4-brane on $\mathbb{R}^{2,1} \times S^2$. A beautiful theme in the study of D-brane intersections is the reconstruction of the same physics from the lower dimensional branes as from the higher dimensional ones *e.g.* [17, 26, 30]. It will be an elegant consistency check of the ABJM proposal to recover the same result directly from a 4-brane probe in an appropriate spacetime background. Given the subtleties related to the counting of vacua in the mass-deformed ABJM theory [33], solving this problem would also shed some light on important issues of the spacetime interpretation. Some hints about the nature of the spacetime geometry probed by the dual D4-brane can be inferred from the fact that preserving some supersymmetry in the 2+1 noncompact directions requires performing a Maldacena-Núñez-type twisted compactification of the higher codimension worldvolume theory [94]. In that case, as well as in the case of 2+1 dimensional Chern-Simons system studied in [95], the radial evolution

of the topologically nontrivial compact sphere was crucial to the consistency of the twisting in the spacetime background solution. However, the example at hand seems to point towards the D4-brane wrapping a topologically trivial but dynamically stable 2-cycle due to external flux, in the spirit of [29, 30]. This expectation is supported by considering the spacetime description of the M2-M5 system, for which the fully backreacted geometry sourced by membranes polarised into fivebranes in the presence of flux was found in [96–98]. Using the M-theory interpretation of the ABJM model, the geometry describing the D4-brane configuration should be obtainable by performing the \mathbb{Z}_k quotient of the former and taking $k \rightarrow \infty$. Our analysis suggests that the action for fluctuations around these D4-brane solutions should see a KK-spectrum with a scale set by the background flux (μ^{-1}) and independent of the size of the brane (R_{ph}). A similar feature is also encountered in the study of fluctuation actions around giant graviton solutions in $\text{AdS}_5 \times S^5$ [99].

Whatever the geometric description, which we will not try to tackle in this paper, the implementation of the twisting requires the flat space 5d Lorentz invariance to first get broken to $\text{SO}(1, 4) \rightarrow \text{SO}(1, 2) \times \text{U}(1)_{34}$. After replacing the 3-4 plane with a sphere there is only a local $\text{U}(1)_{34}$ invariance left. The theory is then twisted with a $\text{U}(1)_R$ Cartan subgroup of the R-symmetry group and the $Q_\alpha^{\dot{\alpha}}$'s of Eq. (6.7) should be spinors under the twisted local rotation group, $\text{U}(1)_T$ [29, 30]. The full details of the twisting are closely connected to the description of fermions, which we did not address here, so we will also leave a full investigation as an open question for future work. A related intriguing question, is whether the same action we have derived from M2, which has structures expected in a Type IIA picture, can be obtained directly from a multiple D2-action.

Alternatively one may consider working from the beginning in a dual picture of 5-branes in M-theory. The starting point would be the M2-M5 system, as described by the M5-worldvolume. At any given fixed radius, there is an S^3 in the M5-worldvolume and an \mathbb{R}^4 transverse to both the M2's and the M5. The transverse \mathbb{R}^4 contains an S^3 surrounding the M2-M5 system. Consider both the worldvolume S^3 and the transverse S^3 as Hopf fibrations. When considering small fluctuations of worldvolume fields around a configuration where the transverse S^3 is of vanishing size and the longitudinal S^3 is large, the quotient acts effectively on the longitudinal S^3 to give a reduction to IIA. By considering fluctuations of the M5-action in this quotient spacetime for $k \rightarrow \infty$, it should be possible to recover our large N action.

Another interesting question is to recover the $\frac{1}{N}$ corrections from these different pictures. We have sketched how these could emerge from the Matrix description in Subsection 4.2.7. From the D4-brane point of view they are expected to arise as corrections in a noncommutative $\text{U}(1)$ theory, where the noncommutativity parameter goes like $\frac{1}{N}$. Seeing $\frac{1}{N}$ corrections from the M5 perspective will likely require an improvement of our current knowledge of 5-brane actions.

- **Fermions, Supersymmetry**

We have focused on the bosonic part of the fluctuation action, while leaving a detailed study of the fermionic terms for the future. As we have already mentioned, doing so would help in elucidating the twisting and the amount of preserved supersymmetry, hence the nature of the D4 compactification. Some of the fermions would be expected to behave as scalars or vectors

under the twisted local rotation group, $U(1)_T$, in order for the fields to appropriately furnish supersymmetry multiplets in 3d.

- **Nonperturbative effects and the M-theory circle**

One way to see the structure of the M-theory direction would be to look at momentum modes along the circle. These are D0-branes in Type IIA. In the standard connection between M5-branes and D4-branes, they translate into 4d-instanton solutions embedded in the D4-brane worldvolume theory. Hence an avenue to seek the physics of the hidden direction would be to study instanton sectors in the $U(1)$ theory we have obtained. This could be done following [100] who investigated instanton effects in the undeformed ABJM theory, while also showing that such effects will start contributing at four-derivative order in an l_p expansion. Similar conclusions in the context of scattering were reached in [101].

9 Summary and conclusions

In this paper we analysed the equations, symmetries and fluctuations for the classical ‘M2-M5’ fuzzy sphere solutions of the $U(N) \times U(N)$ mass-deformed ABJM theory of [33], and equivalently the fuzzy funnel solution of undeformed ABJM. We found them to correspond to a fuzzy S^2 , as opposed to the fuzzy S^3 previously conjectured.

As a warm-up, we showed that a newly found set of defining equations for the Guralnik-Ramgoolam $SO(4)$ -covariant fuzzy S^3 are not compatible with the BPS equations in ABJM, except for the case of $SU(2) \times SU(2)$ gauge group, that is the BLG \mathcal{A}_4 -theory. We then explicitly displayed the symmetries of the solution to find that the physical fluctuations are only invariant under a single $SU(2)$, indicating a fuzzy S^2 , as opposed to an $SU(2) \times SU(2) \simeq SO(4)$ expected for a fuzzy S^3 . The fluctuations were also found to admit an expansion in terms of fuzzy S^2 spherical harmonics (plus some zero modes that decoupled in the large- N limit).

A detailed calculation gave the action for bosonic fluctuations in the large- N limit, Eq. (6.7), in which the fuzzy S^2 matrix algebra approaches the algebra of functions on the commutative (‘classical’) S^2 , and the result was expressed as a $U(1)$ Yang-Mills theory on $\mathbb{R}^{2,1} \times S^2$. This crucially involved a novel version of the Higgs mechanism for the three dimensional Chern-Simons-matter action. The answer is compatible with an interpretation in terms of a D4-brane wrapping the S^2 in Type IIA.

We then explained how the above is in agreement with a spacetime picture in which an M5-brane wraps an S^3/\mathbb{Z}_k , with the three-sphere given by the Hopf fibration of an S^1/\mathbb{Z}_k over S^2 , and identified S^1/\mathbb{Z}_k with the M-theory circle. In the $k \rightarrow \infty$ limit one has a (double) dimensional reduction and a D4-brane wrapping the S^2 in Type IIA string theory, as opposed to an M5-brane wrapping the S^3/\mathbb{Z}_k in M-theory. We also discussed how the Hopf structure persists in the finite- N matrix description, in a new realisation of the fuzzy S^2 base manifold.

In conclusion, the problem of finding a perturbative formulation of multiple membrane theory that would obtain the full classical geometry of S^3 at large N remains open. As seen from the above, one needs to avoid the large- k \mathbb{Z}_k projection, and that seems impossible in ABJM theory. Having a perturbative multiple membrane action at large N in flat space (as opposed to $\mathbb{C}^4/\mathbb{Z}_k$) would seem to be needed for that goal.

Acknowledgements

We would like to thank James Bedford, David Berenstein, David Berman, Robert de Mello Koch, Aki Hashimoto, Koji Hashimoto, Katsushi Ito, Shiraz Minwalla, Sunil Mukhi and Dan Thompson for discussions and comments. HN would like to thank the Queen Mary Physics Department for the opportunity to visit which resulted in the beginning of this work while CP for hospitality during the course of the project. SR is supported by an STFC grant ST/G000565/1. HN's research has been done with partial support from MEXT's program "Promotion of Environmental Improvement for Independence of Young Researchers" under the Special Coordination Funds for Promoting Science and Technology, and also with partial support from MEXT KAKENHI grant nr. 20740128.

A Some useful formulae for the parallel scalars

There are several useful identities that can be used to obtain the action for the matrix fluctuations in Section 4.1, involving the change of basis from traceless symmetric part of $V \otimes \bar{V}$ of $SU(2)$ to spin-1, where V is the fundamental

$$\begin{aligned}
s_i &= s_\alpha^\beta (\tilde{\sigma}_i)_\beta^\alpha \\
s_\beta^\alpha &= \frac{1}{2} s_i (\tilde{\sigma}_i)_\beta^\alpha \\
J_\beta^\alpha &= \frac{(N-1)}{2} \delta_\beta^\alpha + \frac{1}{2} J_i (\tilde{\sigma}_i)_\beta^\alpha \\
J_i &= (\tilde{\sigma}_i)_\beta^\alpha J_\alpha^\beta .
\end{aligned} \tag{A.1}$$

A.1 Identities for the $s_i s_i$ and r - s fluctuations

In order to avoid the proliferation of indices in the calculation, one can also define shorthand notation for several combinations of fields that appear frequently. We have collected a number of such definitions and identities in the following.

A.1.1 $s_i s_i$ fluctuations

Inner products

$$\begin{aligned}
s_\beta^\alpha s_\alpha^\beta &= \frac{1}{2} s^2 \\
J_\beta^\alpha J_\alpha^\beta &= N(N-1) \\
J_i J_i &= (N^2 - 1) \\
s_\beta^\alpha J_\alpha^\beta &= \frac{1}{2} s_i J_i = \frac{1}{2} (s \cdot J) \\
J_\beta^\alpha s_\alpha^\beta &= \frac{1}{2} J_i s_i = \frac{1}{2} (J \cdot s) \\
\text{Tr}(s \wedge J \wedge s) &= i \epsilon_{ijk} \text{Tr}(s_i J_j s_k) .
\end{aligned} \tag{A.2}$$

General quadratic products

$$J_\mu^\alpha s_\beta^\mu = \frac{(N-1)}{2} s_\beta^\alpha + \frac{(J \cdot s)}{4} \delta_\beta^\alpha - \frac{i \epsilon_{ijk}}{4} J_i s_j (\tilde{\sigma}_k)_\beta^\alpha$$

$$s_\mu^\alpha J_\beta^\mu = \frac{(N-1)}{2} s_\beta^\alpha + \frac{(s \cdot J)}{4} \delta_\beta^\alpha - \frac{i\epsilon_{ijk}}{4} s_i J_j (\tilde{\sigma}_k)_\beta^\alpha. \quad (\text{A.3})$$

Cubic products

$$\begin{aligned} J_\mu^\alpha s_\beta^\mu J_\alpha^\beta &= \frac{1}{4}(N-1)(J \cdot s + s \cdot J) - \frac{1}{4} J \wedge s \wedge J \\ s_\mu^\alpha J_\beta^\mu s_\alpha^\beta &= \frac{1}{4}(N-1)s^2 - \frac{1}{4} s \wedge J \wedge s \\ J_\mu^\alpha s_\beta^\mu s_\alpha^\beta &= \frac{1}{4}(N-1)s^2 - \frac{1}{4} J \wedge s \wedge s \\ s_\beta^\alpha J_\gamma^\beta J_\alpha^\gamma &= \frac{1}{2} N(s \cdot J) \\ J_\beta^\alpha J_\gamma^\beta s_\alpha^\gamma &= \frac{1}{2} N(J \cdot s). \end{aligned} \quad (\text{A.4})$$

Quartic traced products

$$\begin{aligned} \text{Tr}(J_\mu^\beta s_\alpha^\mu J_\nu^\alpha s_\beta^\nu) &= -\frac{1}{4}(N-1)\text{Tr}(s^2) + \frac{1}{8}\text{Tr}((J \cdot s)^2 + (s \cdot J)^2) \\ &\quad - \frac{1}{4}(N-1)\text{Tr}(s \wedge J \wedge s) - \frac{1}{16}\text{Tr}([J_i, s_j][J_i, s_j]) \\ \text{Tr}(s_\mu^\alpha s_\beta^\mu J_\nu^\beta J_\alpha^\nu) &= \frac{1}{4}N(N-1)\text{Tr}(s^2) - \frac{1}{4}N\text{Tr}(s \wedge J \wedge s). \end{aligned} \quad (\text{A.5})$$

Other useful formulae

$$\begin{aligned} (\tilde{\sigma}_i)_\beta^\alpha (\tilde{\sigma}_i)_\nu^\mu &= 2\delta_\nu^\alpha \delta_\beta^\mu - \delta_\beta^\alpha \delta_\nu^\mu \\ G^\alpha \bar{J}_i \bar{J}_j G_\alpha^\dagger &= (N-3)J_i J_j + (N-1)\delta_{ij} + i\epsilon_{ijk} J_k. \end{aligned} \quad (\text{A.6})$$

We further define

$$\square s_k \equiv [J_p, [J_p, s_k]]. \quad (\text{A.7})$$

Note that

$$\begin{aligned} \text{Tr}(s_i \square s_i) &= -\text{Tr}[J_k, s_i][J_k, s_i] \\ J_i s_j J_i &= (N^2 - 1)s_j - \frac{1}{2} \square s_j \end{aligned} \quad (\text{A.8})$$

and

$$\begin{aligned} \frac{1}{2}\text{Tr}(s \wedge J \wedge \square s) &= 2(N^2 - 1)\text{Tr}(s \wedge J \wedge s) - \text{Tr}(J \wedge s \wedge J)(J \cdot s + s \cdot J) \\ &\quad - 2\text{Tr}(J \cdot s)^2 + 2(N^2 - 1)s^2 - \text{Tr}(s \cdot \square s) \\ \frac{1}{2}\text{Tr}(J \wedge s \wedge \square s) &= 2(N^2 - 1)\text{Tr}(s \wedge J \wedge s) - \text{Tr}(J \wedge s \wedge J)(J \cdot s + s \cdot J) \\ &\quad - 2\text{Tr}(s \cdot J)^2 + 2(N^2 - 1)s^2 - \text{Tr}(s \cdot \square s) \\ \text{Tr}(s \wedge J \wedge \square s) - \text{Tr}(J \wedge s \wedge \square s) &= 4\text{Tr}((s \cdot J)^2 - (J \cdot s)^2). \end{aligned} \quad (\text{A.9})$$

A.1.2 r - s fluctuations

We also give the following useful definitions for the r - s fluctuations

$$J \cdot r \cdot J \equiv J_i r J_i$$

$$\begin{aligned}
J \wedge s \wedge J &\equiv i\epsilon_{ijk}J_i s_j J_k \\
J_i J_j J_i &= (N^2 - 5)J_j \\
\frac{1}{2}\square(s.J) &= \frac{1}{2}\square s.J + 2(J \wedge s \wedge J) \\
\frac{1}{2}\square(J.s) &= \frac{1}{2}J.\square s + 2(J \wedge s \wedge J) \\
\frac{1}{2}(\square(s.J) + \square(J.s)) &= \frac{1}{2}(\square s.J + J.\square s) + 4(J \wedge s \wedge J) \\
\frac{1}{2}\text{Tr}(rJ.\square s) &= (N^2 - 1)\text{Tr}(r(J.s)) - \text{Tr}(J.r.J)(J.s) - 2\text{Tr}(r(J \wedge s \wedge J)) \\
\frac{1}{2}\text{Tr}(r\square s.J) &= (N^2 - 1)\text{Tr}(r(s.J)) - \text{Tr}(J.r.J)(s.J) - 2\text{Tr}(r(J \wedge s \wedge J)) \\
\frac{1}{2}\text{Tr}(rs.\square J + r\square J.s) &= 2\text{Tr}(r(J.s + s.J)) \\
\text{Tr}A\square B &= \text{Tr}(\square A)B.
\end{aligned} \tag{A.10}$$

In the last line we can have $A = r, B = J.s + s.J$ for example. One also has

$$\begin{aligned}
J_\gamma^\beta s_\mu^\alpha J_\beta^\mu J_\alpha^\gamma &= -\frac{1}{16}(\square(s.J) + (\square s).J) + \frac{1}{4}(N-1)(2N+1)(s.J) - \frac{1}{4}(N-1)(J.s) \\
J_\gamma^\beta J_\mu^\alpha s_\beta^\mu J_\alpha^\gamma &= -\frac{1}{16}(\square(J.s) + J.(\square s)) + \frac{1}{4}(N-1)(2N+1)(J.s) - \frac{1}{4}(N-1)(s.J) \\
J_\gamma^\beta s_\mu^\alpha J_\beta^\mu J_\alpha^\gamma + J_\gamma^\beta J_\mu^\alpha s_\beta^\mu J_\alpha^\gamma &= -\frac{1}{16}(\square(s.J + J.s) + (\square s).J + J.(\square s)) + \frac{1}{2}N(N-1)(J.s + s.J).
\end{aligned} \tag{A.11}$$

A.2 Expressions and identities for s_i decomposition

In obtaining the action on the classical sphere in Section 4.2 we made use of a set of Killing vectors K_i^a . The explicit formulae for the latter are given by

$$\begin{aligned}
K_1^\theta &= -\sin\phi & K_1^\phi &= -\cot\theta \cos\phi \\
K_2^\theta &= \cos\phi & K_2^\phi &= -\cot\theta \sin\phi \\
K_3^\theta &= 0 & K_3^\phi &= 1,
\end{aligned} \tag{A.12}$$

as given in [26]. The relations between Cartesian and spherical coordinates is

$$\begin{aligned}
x_1 &= \sin\theta \cos\phi \\
x_2 &= \sin\theta \sin\phi \\
x_3 &= \cos\theta.
\end{aligned} \tag{A.13}$$

Other formulae that we have made use of include

$$\begin{aligned}
K_i^a K_i^b &= h^{ab} \\
x_i K_i^a &= 0 \\
K_i^a K_i^b \partial_a x_j \partial_b x_j &= 2 \\
\epsilon_{ijk} x_i K_j^a K_k^b &= \frac{\epsilon^{ab}}{\sin\theta} = \omega^{ab} \\
K_i^a K_j^b h_{ab} &= \delta_{ij} - x_i x_j \\
K_i^a (\partial_a K_i^b) &= \frac{1}{\sqrt{h}} (\partial^b \sqrt{h}) \\
K_i^a (\partial_a K_i^b) \partial_b &= K_i(K_i) = \epsilon_{ijk} x_j \epsilon_{ilm} \partial_k(x_l) \partial_m = -2x_i \partial_i.
\end{aligned} \tag{A.14}$$

B Technical aspects of Higgsing

B.1 Derivation of Eq.(5.12)

It is easy to experimentally verify that commuting a number of \bar{J}_i 's past G^α , while then contracting with G_α^\dagger gives

$$\begin{aligned}
G^\alpha \bar{J}_i G_\alpha^\dagger &= J J_i - J_i \\
G^\alpha \bar{J}_i \bar{J}_j G_\alpha^\dagger &= J J_i J_j - 2 J_i J_j + \delta_{ij} \\
G^\alpha \bar{J}_i \bar{J}_j \bar{J}_k G_\alpha^\dagger &= J J_i J_j J_k - 3 J_i J_j J_k \\
&\quad + J_i \delta_{jk} + J_j \delta_{ik} + J_k \delta_{ij} \\
&\quad - i \epsilon_{ijs} J_k J_s - i \epsilon_{jks} J_i J_m - i \epsilon_{kis} J_j J_s \\
G^\alpha \bar{J}_i \bar{J}_j \bar{J}_k \bar{J}_n G_\alpha^\dagger &= J J_i J_j J_k J_n - 4 J_i J_j J_k J_n \\
&\quad + J_i J_j \delta_{kn} + \text{permutations} \\
&\quad - i \epsilon_{ijs} J_k J_n J_s + \text{permutations} \\
&\quad \vdots
\end{aligned} \tag{B.1}$$

and so on. Higher powers of \bar{J}_i will follow a similar pattern involving δ 's and ϵ 's for subleading terms, as these are the only invariant tensors of SU(2). However, we are mainly interested in commuting the whole set of fuzzy spherical harmonics. We remind that the latter are defined as

$$Y_{lm}(\bar{J}_i) = \sum_i f_{lm}^{(i_1 \dots i_l)} \bar{J}_{i_1} \dots \bar{J}_{i_l}, \tag{B.2}$$

with $f_{lm}^{(i_1 \dots i_l)}$ a symmetric traceless tensor in i . We then have

$$\begin{aligned}
f_{lm}^i G^\alpha \bar{J}_i G_\alpha^\dagger &= f_{lm}^i (J J_i - J_i) \\
f_{lm}^{(ij)} G^\alpha \bar{J}_i \bar{J}_j G_\alpha^\dagger &= f_{lm}^{(ij)} (J J_i J_j - 2 J_i J_j) \\
f_{lm}^{(ijk)} G^\alpha \bar{J}_i \bar{J}_j \bar{J}_k G_\alpha^\dagger &= f_{lm}^{(ijk)} (J J_i J_j J_k - 3 J_i J_j J_k) \\
f_{lm}^{(ijkn)} G^\alpha \bar{J}_i \bar{J}_j \bar{J}_k \bar{J}_n G_\alpha^\dagger &= f_{lm}^{(ijkn)} (J J_i J_j J_k J_n - 4 J_i J_j J_k J_n) \\
&\quad \vdots \\
f_{lm}^{(i_1 \dots i_l)} G^\alpha \bar{J}_{i_1} \dots \bar{J}_{i_l} G_\alpha^\dagger &= f_{lm}^{(i_1 \dots i_l)} (J J_{i_1} \dots J_{i_l} - l J_{i_1} \dots J_{i_l}),
\end{aligned} \tag{B.3}$$

that is only the first two terms survive in each expression, as the \bar{f}_{lm} 's project out all the remaining contributions. Summing over the above, this leads to the exact relation

$$G^\alpha Y_{lm}(\bar{J}_i) G_\alpha^\dagger = J Y_{lm}(J_i) - l Y_{lm}(J_i). \tag{B.4}$$

B.2 Higgsing from trace over \mathbf{V}^-

We could have chosen to commute $Y_{lm}(J_i)$ past the G^α 's in the third term of (5.8). This would have led to the same final answer for the action for fluctuations, but involving a trace over \mathbf{V}^- in the intermediate steps.¹⁷ In order to see this one needs to make use of the following exact expression, analogous to (B.4),

$$G_\alpha^\dagger Y_{lm}(J_i) G^\alpha = \bar{J} Y_{lm}(\bar{J}_i) + l Y_{lm}(\bar{J}_i), \tag{B.5}$$

¹⁷More precisely this restricts to a trace over V_{N-1}^- of \mathbf{V}^- .

which can then get converted to classical sphere variables using the normalisations (5.6)

$$\begin{aligned}
G_\alpha^\dagger Y_{lm}(x_i) G^\alpha &= \left(\frac{N^2 - 2N}{N^2 - 1} \right)^{\frac{1}{2}} (\bar{J} Y_{lm}(\bar{x}_i) + l Y_{lm}(\bar{x}_i)) \\
&= N Y_{lm}(\bar{x}_i) - \frac{l(l+1)}{2N} Y_{lm}(\bar{x}_i) + \mathcal{O}\left(\frac{1}{N^2}\right) \\
&= N Y_{lm}(\bar{x}_i) + \frac{1}{2N} \hat{\square} Y_{lm}(\bar{x}_i) + \mathcal{O}\left(\frac{1}{N^2}\right). \tag{B.6}
\end{aligned}$$

After plugging the above back into (5.8) we end up with the same subleading term in (5.15). This is very satisfying, since we expect to obtain a single answer for the classical action regardless of whether we use x_i or \bar{x}_i , although it was not obvious that this would happen at the outset of the calculation. The result reflects the \mathbb{Z}_2 symmetry inherent in the derivation of the classical action. Note however that beyond the strict large N approximation the difference between x_i and \bar{x}_i (and consequently also the difference between $A^{(i)}$ and $\hat{A}^{(i)}$) could become manifest. To probe that regime one ought to first get a better understanding of the finite N definitions of the fields.

C Towards multiple M5-branes

In the case of the usual Myers effect [22] (multiple D0-branes expanding into spherical D2-branes), the equations of motion for N D0-branes give

$$[\Phi_i, \Phi_j] = i f \epsilon_{ijk} \Phi_k, \tag{C.1}$$

where Φ_i are the transverse scalars on the D0 worldvolume with $i = 1, 2, 3$ and the rest of them set to zero. These can be solved by $\Phi_i = X_i$ an $N \times N$ irrep of $SU(2)$, resulting into a single spherical D2-brane, or by $\Phi_i = X_i^{(1)} \oplus X_i^{(2)}$, where $X_i^{(1)}$ is an irrep of size N_1 and $X_i^{(2)}$ is an irrep of size N_2 , with $N = N_1 + N_2$, resulting into two concentric spherical D2-branes with radii that depend on N_1, N_2 . This is the case because the sum of two irreps of $SU(2)$ is a reducible representation

$$[X_i^{(1)} \oplus X_i^{(2)}, X_j^{(1)} \oplus X_j^{(2)}] = [X_i^{(1)}, X_j^{(1)}] \oplus [X_i^{(2)}, X_j^{(2)}] = i \epsilon_{ijk} (X_k^{(1)} \oplus X_k^{(2)}). \tag{C.2}$$

A similar property arises in our case, for the equation of motion

$$R^\alpha = R^\alpha R_\beta^\dagger R^\beta - R^\beta R_\beta^\dagger R^\alpha \tag{C.3}$$

and solutions defined on the space $\mathbf{V}^+ \oplus \mathbf{V}^-$, where $\mathbf{V}^+ = V_N^+$ and $\mathbf{V}^- = V_{N-1}^- \oplus V_1^-$. The single 4-brane solutions we have been describing in this paper, with a fuzzy S^2 , can be written as

$$\begin{aligned}
G^\alpha &= \mathcal{P}_{V_N^+} G^\alpha \mathcal{P}_{V_{N-1}^-} & G_\alpha^\dagger &= \mathcal{P}_{V_{N-1}^-} G_\alpha^\dagger \mathcal{P}_{V_N^+} \\
G^\beta G_\beta^\dagger &= (N-1) \mathcal{P}_{V_N^+} & G_\beta^\dagger G^\beta &= N \mathcal{P}_{V_{N-1}^-}, \tag{C.4}
\end{aligned}$$

where \mathcal{P} 's are projectors $\mathcal{P}^2 = \mathcal{P}$ on each subspace of the total space, giving

$$G^\alpha G_\beta^\dagger G^\beta = N \mathcal{P}_{V_N^+} G^\alpha \mathcal{P}_{V_{N-1}^-}, \quad G^\beta G_\beta^\dagger G^\alpha = (N-1) \mathcal{P}_{V_N^+} G^\alpha \mathcal{P}_{V_{N-1}^-}. \tag{C.5}$$

Then the multi-4-brane solutions are defined on a space

$$\mathbf{V}^+ = V_{N_1}^+ \oplus V_{N_2}^+$$

$$\mathbf{V}^- = (V_{N_1-1}^- \oplus V_1^-) \oplus (V_{N_2-1}^- \oplus V_1^-) \quad (\text{C.6})$$

by the ansatz

$$\begin{aligned} G^\alpha &= G_1^\alpha \oplus G_2^\alpha \\ G^\dagger_\alpha &= G_{1\alpha}^\dagger \oplus G_{2\alpha}^\dagger, \end{aligned} \quad (\text{C.7})$$

where as before

$$\begin{aligned} G_i^\alpha &= \mathcal{P}_{V_{N_i}^+} G_i^\alpha \mathcal{P}_{V_{N_i-1}^-} \\ G_{i\alpha}^\dagger &= \mathcal{P}_{V_{N_i-1}^-} G_{i\alpha}^\dagger \mathcal{P}_{V_{N_i}^+}. \end{aligned} \quad (\text{C.8})$$

The equations of motion are trivially satisfied since

$$\begin{aligned} G^\alpha G^\beta G^\dagger_\beta &= (N_1 - 1) G_1^\alpha \mathcal{P}_{V_{N_1}^+} \oplus (N_2 - 1) G_2^\alpha \mathcal{P}_{V_{N_2}^+} \\ &= (N_1 - 1) \mathcal{P}_{V_{N_1-1}^-} G_1^\alpha \mathcal{P}_{V_{N_1}^+} \oplus (N_2 - 1) \mathcal{P}_{V_{N_2-1}^-} G_2^\alpha \mathcal{P}_{V_{N_2}^+} \\ G^\dagger_\beta G^\beta G^\alpha &= N_1 \mathcal{P}_{V_{N_1-1}^-} G_2^\alpha \mathcal{P}_{V_{N_1}^+} \oplus N_2 \mathcal{P}_{V_{N_2-1}^-} G_2^\alpha \mathcal{P}_{V_{N_2}^+}. \end{aligned} \quad (\text{C.9})$$

One could also think of more general block-diagonal combinations with $N_1 + N_2 + N_3 + \dots = N$, corresponding to concentric spherical D4-branes, the radius of which depends on the dimension of each representation for a fixed s , as per (7.1) in the funnel case. These are also all the zero energy solutions for fixed μ in the mass-deformed case and the collection of partitions of N parametrise the set of classical vacua of the mass-deformed theory, as discussed in [33].¹⁸

Of particular interest are the possibilities with m copies of $N_m \times N_m$ equally sized blocks, where $mN_m = N$, since in that case the branes are coincident and there is a worldvolume gauge symmetry enhancement to $U(m)$. Clarifying how to go to spherical (S^3) M5-branes by blowing up the Hopf fibre over S^2 , perhaps along the lines of [89], in combination with the above could provide a starting point for studying multiple fivebranes in M-theory.

References

- [1] J. Bagger and N. Lambert, “Modeling multiple M2’s,” *Phys. Rev.* **D75** (2007) 045020, [arXiv:hep-th/0611108](#).
- [2] J. Bagger and N. Lambert, “Gauge Symmetry and Supersymmetry of Multiple M2-Branes,” *Phys. Rev.* **D77** (2008) 065008, [arXiv:0711.0955 \[hep-th\]](#).
- [3] J. Bagger and N. Lambert, “Comments On Multiple M2-branes,” *JHEP* **02** (2008) 105, [arXiv:0712.3738 \[hep-th\]](#).
- [4] A. Gustavsson, “Algebraic structures on parallel M2-branes,” *Nucl. Phys.* **B811** (2009) 66–76, [arXiv:0709.1260 \[hep-th\]](#).

¹⁸There is a puzzle related to this counting involving possible partitions corresponding to solutions with both $R^\alpha \neq 0, Q^\alpha \neq 0$.

- [5] J. H. Schwarz, “Superconformal Chern-Simons theories,” *JHEP* **11** (2004) 078, [arXiv:hep-th/0411077](#).
- [6] A. Basu and J. A. Harvey, “The M2-M5 brane system and a generalized Nahm’s equation,” *Nucl. Phys.* **B713** (2005) 136–150, [arXiv:hep-th/0412310](#).
- [7] S. Mukhi and C. Papageorgakis, “M2 to D2,” *JHEP* **05** (2008) 085, [arXiv:0803.3218 \[hep-th\]](#).
- [8] M. A. Bandres, A. E. Lipstein, and J. H. Schwarz, “N = 8 Superconformal Chern–Simons Theories,” *JHEP* **05** (2008) 025, [arXiv:0803.3242 \[hep-th\]](#).
- [9] M. Van Raamsdonk, “Comments on the Bagger-Lambert theory and multiple M2- branes,” *JHEP* **05** (2008) 105, [arXiv:0803.3803 \[hep-th\]](#).
- [10] N. Lambert and D. Tong, “Membranes on an Orbifold,” *Phys. Rev. Lett.* **101** (2008) 041602, [arXiv:0804.1114 \[hep-th\]](#).
- [11] J. Distler, S. Mukhi, C. Papageorgakis, and M. Van Raamsdonk, “M2-branes on M-folds,” *JHEP* **05** (2008) 038, [arXiv:0804.1256 \[hep-th\]](#).
- [12] D. Gaiotto and E. Witten, “Janus Configurations, Chern-Simons Couplings, And The Theta-Angle in N=4 Super Yang-Mills Theory,” [arXiv:0804.2907 \[hep-th\]](#).
- [13] K. Hosomichi, K.-M. Lee, S. Lee, S. Lee, and J. Park, “N=4 Superconformal Chern-Simons Theories with Hyper and Twisted Hyper Multiplets,” *JHEP* **07** (2008) 091, [arXiv:0805.3662 \[hep-th\]](#).
- [14] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, “N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals,” *JHEP* **10** (2008) 091, [arXiv:0806.1218 \[hep-th\]](#).
- [15] P. S. Howe, N. D. Lambert, and P. C. West, “The self-dual string soliton,” *Nucl. Phys.* **B515** (1998) 203–216, [arXiv:hep-th/9709014](#).
- [16] N. R. Constable, R. C. Myers, and O. Tafjord, “The noncommutative bion core,” *Phys. Rev.* **D61** (2000) 106009, [arXiv:hep-th/9911136](#).
- [17] N. R. Constable, R. C. Myers, and O. Tafjord, “Non-Abelian brane intersections,” *JHEP* **06** (2001) 023, [arXiv:hep-th/0102080](#).
- [18] P. L. H. Cook, R. de Mello Koch, and J. Murugan, “Non-Abelian Bionic brane intersections,” *Phys. Rev.* **D68** (2003) 126007, [arXiv:hep-th/0306250](#).
- [19] J. Hoppe, “*Quantum Theory of a Massless Relativistic Surface and a Two-Dimensional Bound State Problem*”. PhD thesis, Massachusetts Institute of Technology, Dept. of Physics, 1982. <http://hdl.handle.net/1721.1/15717>.
- [20] J. Hoppe, “Diffeomorphism Groups, Quantization and SU(infinity),” *Int. J. Mod. Phys.* **A4** (1989) 5235.

- [21] J. Madore, “The fuzzy sphere,” *Class. Quant. Grav.* **9** (1992) 69–88.
- [22] R. C. Myers, “Dielectric-branes,” *JHEP* **12** (1999) 022, [arXiv:hep-th/9910053](#).
- [23] S. Iso, Y. Kimura, K. Tanaka, and K. Wakatsuki, “Noncommutative gauge theory on fuzzy sphere from matrix model,” *Nucl. Phys.* **B604** (2001) 121–147, [arXiv:hep-th/0101102](#).
- [24] K. Dasgupta, M. M. Sheikh-Jabbari, and M. Van Raamsdonk, “Matrix perturbation theory for M-theory on a PP-wave,” *JHEP* **05** (2002) 056, [arXiv:hep-th/0205185](#).
- [25] K. Dasgupta, M. M. Sheikh-Jabbari, and M. Van Raamsdonk, “Protected multiplets of M-theory on a plane wave,” *JHEP* **09** (2002) 021, [arXiv:hep-th/0207050](#).
- [26] C. Papageorgakis, S. Ramgoolam, and N. Toumbas, “Noncommutative geometry, quantum effects and DBI-scaling in the collapse of D0-D2 bound states,” *JHEP* **01** (2006) 030, [arXiv:hep-th/0510144](#).
- [27] J. Polchinski and M. J. Strassler, “The string dual of a confining four-dimensional gauge theory,” [arXiv:hep-th/0003136](#).
- [28] N. Dorey and A. Sinkovics, “ $N = 1^*$ vacua, fuzzy spheres and integrable systems,” *JHEP* **07** (2002) 032, [arXiv:hep-th/0205151](#).
- [29] R. P. Andrews and N. Dorey, “Spherical deconstruction,” *Phys. Lett.* **B631** (2005) 74–82, [arXiv:hep-th/0505107](#).
- [30] R. P. Andrews and N. Dorey, “Deconstruction of the Maldacena-Nunez compactification,” *Nucl. Phys.* **B751** (2006) 304–341, [arXiv:hep-th/0601098](#).
- [31] I. Bena, “The M-theory dual of a 3 dimensional theory with reduced supersymmetry,” *Phys. Rev.* **D62** (2000) 126006, [arXiv:hep-th/0004142](#).
- [32] K. Hosomichi, K.-M. Lee, S. Lee, S. Lee, and J. Park, “ $N=5,6$ Superconformal Chern-Simons Theories and M2-branes on Orbifolds,” *JHEP* **09** (2008) 002, [arXiv:0806.4977 \[hep-th\]](#).
- [33] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk, and H. Verlinde, “A Massive Study of M2-brane Proposals,” *JHEP* **09** (2008) 113, [arXiv:0807.1074 \[hep-th\]](#).
- [34] S. Terashima, “On M5-branes in $N=6$ Membrane Action,” *JHEP* **08** (2008) 080, [arXiv:0807.0197 \[hep-th\]](#).
- [35] K. Hanaki and H. Lin, “M2-M5 Systems in $N=6$ Chern-Simons Theory,” *JHEP* **09** (2008) 067, [arXiv:0807.2074 \[hep-th\]](#).
- [36] T. Fujimori, K. Iwasaki, Y. Kobayashi, and S. Sasaki, “Time-dependent and Non-BPS Solutions in $N=6$ Superconformal Chern-Simons Theory,” *JHEP* **12** (2008) 023, [arXiv:0809.4778 \[hep-th\]](#).
- [37] M. Arai, C. Montonen, and S. Sasaki, “Vortices, Q-balls and Domain Walls on Dielectric M2-branes,” [arXiv:0812.4437 \[hep-th\]](#).

- [38] A. M. Low, “N=6 Membrane Worldvolume Superalgebra,” [arXiv:0903.0988 \[hep-th\]](#).
- [39] Z. Guralnik and S. Ramgoolam, “On the polarization of unstable D0-branes into non-commutative odd spheres,” *JHEP* **02** (2001) 032, [arXiv:hep-th/0101001](#).
- [40] S. Ramgoolam, “On spherical harmonics for fuzzy spheres in diverse dimensions,” *Nucl. Phys.* **B610** (2001) 461–488, [arXiv:hep-th/0105006](#).
- [41] S. Ramgoolam, “Higher dimensional geometries related to fuzzy odd- dimensional spheres,” *JHEP* **10** (2002) 064, [arXiv:hep-th/0207111](#).
- [42] T. Azuma and M. Bagnoud, “Curved-space classical solutions of a massive supermatrix model,” *Nucl. Phys.* **B651** (2003) 71–86, [arXiv:hep-th/0209057](#).
- [43] B. Janssen, Y. Lozano, and D. Rodriguez-Gomez, “A microscopical description of giant gravitons. II: The AdS(5) x S**5 background,” *Nucl. Phys.* **B669** (2003) 363–378, [arXiv:hep-th/0303183](#).
- [44] B. P. Dolan and D. O’Connor, “A fuzzy three sphere and fuzzy tori,” *JHEP* **10** (2003) 060, [arXiv:hep-th/0306231](#).
- [45] V. P. Nair and S. Randjbar-Daemi, “Quantum Hall effect on S(3), edge states and fuzzy S(3)/Z(2),” *Nucl. Phys.* **B679** (2004) 447–463, [arXiv:hep-th/0309212](#).
- [46] H. Nastase, “On fuzzy spheres and (M)atrix actions,” [arXiv:hep-th/0410137](#).
- [47] M. M. Sheikh-Jabbari, “Tiny graviton matrix theory: DLCQ of IIB plane-wave string theory, a conjecture,” *JHEP* **09** (2004) 017, [arXiv:hep-th/0406214](#).
- [48] B. Janssen, Y. Lozano, and D. Rodriguez-Gomez, “Giant gravitons and fuzzy CP(2),” *Nucl. Phys.* **B712** (2005) 371–391, [arXiv:hep-th/0411181](#).
- [49] Y. Lozano and D. Rodriguez-Gomez, “Fuzzy 5-spheres and pp-wave Matrix actions,” *JHEP* **08** (2005) 044, [arXiv:hep-th/0505073](#).
- [50] C. Papageorgakis and S. Ramgoolam, “On time-dependent collapsing branes and fuzzy odd-dimensional spheres,” *Int. J. Mod. Phys.* **A21** (2006) 6055–6086, [arXiv:hep-th/0603239](#).
- [51] Y. Lozano and D. Rodriguez-Gomez, “Type II pp-wave matrix models from point-like gravitons,” *JHEP* **08** (2006) 022, [arXiv:hep-th/0606057](#).
- [52] J. M. Maldacena, M. M. Sheikh-Jabbari, and M. Van Raamsdonk, “Transverse fivebranes in matrix theory,” *JHEP* **01** (2003) 038, [arXiv:hep-th/0211139](#).
- [53] D. S. Berman, “M-theory branes and their interactions,” *Phys. Rept.* **456** (2008) 89–126, [arXiv:0710.1707 \[hep-th\]](#).
- [54] N. B. Copland, “Aspects of M-Theory Brane Interactions and String Theory Symmetries,” [arXiv:0707.1317 \[hep-th\]](#).

- [55] D. S. Berman, L. C. Tadrowski, and D. C. Thompson, “Aspects of Multiple Membranes,” *Nucl. Phys.* **B802** (2008) 106–120, [arXiv:0803.3611 \[hep-th\]](#).
- [56] P.-M. Ho, Y. Imamura, and Y. Matsuo, “M2 to D2 revisited,” *JHEP* **07** (2008) 003, [arXiv:0805.1202 \[hep-th\]](#).
- [57] P.-M. Ho, Y. Imamura, Y. Matsuo, and S. Shiba, “M5-brane in three-form flux and multiple M2-branes,” *JHEP* **08** (2008) 014, [arXiv:0805.2898 \[hep-th\]](#).
- [58] P.-M. Ho and Y. Matsuo, “M5 from M2,” *JHEP* **06** (2008) 105, [arXiv:0804.3629 \[hep-th\]](#).
- [59] I. A. Bandos and P. K. Townsend, “Light-cone M5 and multiple M2-branes,” *Class. Quant. Grav.* **25** (2008) 245003, [arXiv:0806.4777 \[hep-th\]](#).
- [60] C.-S. Chu and D. J. Smith, “Towards the Quantum Geometry of the M5-brane in a Constant C -Field from Multiple Membranes,” [arXiv:0901.1847 \[hep-th\]](#).
- [61] G. Bonelli, A. Tanzini, and M. Zabzine, “Topological branes, p -algebras and generalized Nahm equations,” *Phys. Lett.* **B672** (2009) 390–395, [arXiv:0807.5113 \[hep-th\]](#).
- [62] C. Krishnan and C. Maccaferri, “Membranes on Calibrations,” *JHEP* **07** (2008) 005, [arXiv:0805.3125 \[hep-th\]](#).
- [63] C. Krishnan and C. Maccaferri, “Membranes, Strings and Integrability,” [arXiv:0810.3825 \[hep-th\]](#).
- [64] G. Papadopoulos, “M2-branes, 3-Lie Algebras and Plucker relations,” *JHEP* **05** (2008) 054, [arXiv:0804.2662 \[hep-th\]](#).
- [65] J. P. Gauntlett and J. B. Gutowski, “Constraining Maximally Supersymmetric Membrane Actions,” [arXiv:0804.3078 \[hep-th\]](#).
- [66] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni, and H. Verlinde, “ $N=8$ superconformal gauge theories and M2 branes,” *JHEP* **01** (2009) 078, [arXiv:0805.1087 \[hep-th\]](#).
- [67] J. Gomis, G. Milanese, and J. G. Russo, “Bagger-Lambert Theory for General Lie Algebras,” *JHEP* **06** (2008) 075, [arXiv:0805.1012 \[hep-th\]](#).
- [68] M. A. Bandres, A. E. Lipstein, and J. H. Schwarz, “Ghost-Free Superconformal Action for Multiple M2-Branes,” *JHEP* **07** (2008) 117, [arXiv:0806.0054 \[hep-th\]](#).
- [69] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk, and H. Verlinde, “Supersymmetric Yang-Mills Theory From Lorentzian Three- Algebras,” *JHEP* **08** (2008) 094, [arXiv:0806.0738 \[hep-th\]](#).
- [70] B. Ezhuthachan, S. Mukhi, and C. Papageorgakis, “D2 to D2,” *JHEP* **07** (2008) 041, [arXiv:0806.1639 \[hep-th\]](#).
- [71] H. Verlinde, “D2 or M2? A Note on Membrane Scattering,” [arXiv:0807.2121 \[hep-th\]](#).

- [72] S. Cecotti and A. Sen, “Coulomb Branch of the Lorentzian Three Algebra Theory,” [arXiv:0806.1990 \[hep-th\]](#).
- [73] E. Antonyan and A. A. Tseytlin, “On 3d N=8 Lorentzian BLG theory as a scaling limit of 3d superconformal N=6 ABJM theory,” *Phys. Rev.* **D79** (2009) 046002, [arXiv:0811.1540 \[hep-th\]](#).
- [74] Y. Honma, S. Iso, Y. Sumitomo, and S. Zhang, “Scaling limit of N=6 superconformal Chern-Simons theories and Lorentzian Bagger-Lambert theories,” *Phys. Rev.* **D78** (2008) 105011, [arXiv:0806.3498 \[hep-th\]](#).
- [75] Y. Honma, S. Iso, Y. Sumitomo, H. Umetsu, and S. Zhang, “Generalized Conformal Symmetry and Recovery of SO(8) in Multiple M2 and D2 Branes,” [arXiv:0807.3825 \[hep-th\]](#).
- [76] J. Kluson, “Remark About Scaling Limit of ABJ Theory,” [arXiv:0902.4122 \[hep-th\]](#).
- [77] O. Aharony, O. Bergman, and D. L. Jafferis, “Fractional M2-branes,” *JHEP* **11** (2008) 043, [arXiv:0807.4924 \[hep-th\]](#).
- [78] J. Bagger and N. Lambert, “Three-Algebras and N=6 Chern-Simons Gauge Theories,” *Phys. Rev.* **D79** (2009) 025002, [arXiv:0807.0163 \[hep-th\]](#).
- [79] M. Schnabl and Y. Tachikawa, “Classification of N=6 superconformal theories of ABJM type,” [arXiv:0807.1102 \[hep-th\]](#).
- [80] M. Benna, I. Klebanov, T. Klose, and M. Smedback, “Superconformal Chern-Simons Theories and AdS4/CFT3 Correspondence,” *JHEP* **09** (2008) 072, [arXiv:0806.1519 \[hep-th\]](#).
- [81] T. Ishii, G. Ishiki, S. Shimasaki, and A. Tsuchiya, “N=4 Super Yang-Mills from the Plane Wave Matrix Model,” *Phys. Rev.* **D78** (2008) 106001, [arXiv:0807.2352 \[hep-th\]](#).
- [82] D. E. Berenstein, J. M. Maldacena, and H. S. Nastase, “Strings in flat space and pp waves from N = 4 super Yang Mills,” *JHEP* **04** (2002) 013, [arXiv:hep-th/0202021](#).
- [83] R. Jackiw and C. Rebbi, “Spin from Isospin in a Gauge Theory,” *Phys. Rev. Lett.* **36** (1976) 1116.
- [84] S. R. Das and A. Jevicki, “String Field Theory and Physical Interpretation of d = 1 Strings,” *Mod. Phys. Lett. A* **5** (1990) 1639.
- [85] B. Ezhuthachan, S. Mukhi, and C. Papageorgakis, “The Power of the Higgs Mechanism: Higher-Derivative BLG Theories,” [arXiv:0903.0003 \[hep-th\]](#).
- [86] Y. Pang and T. Wang, “From N M2’s to N D2’s,” *Phys. Rev.* **D78** (2008) 125007, [arXiv:0807.1444 \[hep-th\]](#).
- [87] T. Li, Y. Liu, and D. Xie, “Multiple D2-Brane Action from M2-Branes,” [arXiv:0807.1183 \[hep-th\]](#).

- [88] M. Nakahara, “Geometry, topology and physics,”. Bristol, UK: Hilger (1990) 505 p. (Graduate student series in physics).
- [89] T. Ishii, G. Ishiki, S. Shimasaki, and A. Tsuchiya, “Fiber Bundles and Matrix Models,” *Phys. Rev.* **D77** (2008) 126015, [arXiv:0802.2782 \[hep-th\]](#).
- [90] A. P. Balachandran, S. Kurkcuoglu, and S. Vaidya, “Lectures on fuzzy and fuzzy SUSY physics,” [arXiv:hep-th/0511114](#).
- [91] D. N. Kabat and W. Taylor, “Spherical membranes in matrix theory,” *Adv. Theor. Math. Phys.* **2** (1998) 181–206, [arXiv:hep-th/9711078](#).
- [92] C.-S. Chu, P.-M. Ho, Y. Matsuo, and S. Shiba, “Truncated Nambu-Poisson Bracket and Entropy Formula for Multiple Membranes,” *JHEP* **08** (2008) 076, [arXiv:0807.0812 \[hep-th\]](#).
- [93] I. A. Bandos and P. K. Townsend, “SDiff Gauge Theory and the M2 Condensate,” [arXiv:0808.1583 \[hep-th\]](#).
- [94] J. M. Maldacena and C. Nunez, “Towards the large N limit of pure N = 1 super Yang Mills,” *Phys. Rev. Lett.* **86** (2001) 588–591, [arXiv:hep-th/0008001](#).
- [95] J. M. Maldacena and H. S. Nastase, “The supergravity dual of a theory with dynamical supersymmetry breaking,” *JHEP* **09** (2001) 024, [arXiv:hep-th/0105049](#).
- [96] C. N. Pope and N. P. Warner, “A dielectric flow solution with maximal supersymmetry,” *JHEP* **04** (2004) 011, [arXiv:hep-th/0304132](#).
- [97] I. Bena and N. P. Warner, “A harmonic family of dielectric flow solutions with maximal supersymmetry,” *JHEP* **12** (2004) 021, [arXiv:hep-th/0406145](#).
- [98] H. Lin, O. Lunin, and J. M. Maldacena, “Bubbling AdS space and 1/2 BPS geometries,” *JHEP* **10** (2004) 025, [arXiv:hep-th/0409174](#).
- [99] S. R. Das, A. Jevicki, and S. D. Mathur, “Vibration modes of giant gravitons,” *Phys. Rev.* **D63** (2001) 024013, [arXiv:hep-th/0009019](#).
- [100] K. Hosomichi *et al.*, “A Nonperturbative Test of M2-Brane Theory,” *JHEP* **11** (2008) 058, [arXiv:0809.1771 \[hep-th\]](#).
- [101] T. Hirayama and D. Tomino, “Membrane scattering from Bagger-Lambert theory,” [arXiv:0812.1866 \[hep-th\]](#).