

1 Towards Ecosystem-Based Management: identifying operational food-web  
2 indicators for marine ecosystems

3 Jamie C. Tam<sup>1\*</sup>, Jason S. Link<sup>1</sup>, Axel G. Rossberg<sup>2,3</sup>, Stuart I. Rogers<sup>3</sup>

4<sup>1</sup> National Oceanographic and Atmospheric Administration, National Marine Fisheries  
5 Service, 166 Water St., Woods Hole, MA, USA. 02543.

6<sup>2</sup> Queen Mary University of London, School of Biological Sciences, Mile End Road, London  
7 E1 4NS

8<sup>3</sup> Centre for Environment Fisheries and Aquaculture Science. Pakefield Road, Lowestoft,  
9 Suffolk, NR33 0HT, UK.

10 \*Corresponding author [jamie.tam@noaa.gov](mailto:jamie.tam@noaa.gov)

11

12 **Keywords:** Ecosystem-Based Management, food-web indicators, operational ecological  
13 indicators, selection criteria

## 14 Abstract

15 Modern approaches to Ecosystem-Based Management and sustainable use of marine  
16 resources must account for the myriad impacts (interspecies, human and environmental  
17 pressures) affecting marine ecosystems. The network of feeding interactions between co-  
18 existing species and populations (food webs) are an important aspect of all marine  
19 ecosystems and biodiversity. Here we describe and discuss a quantitative process to evaluate  
20 the selection of operational food-web indicators for use in evaluating marine ecosystem  
21 status. This process brought together experts in food-web ecology, marine ecology, and  
22 resource management, to identify available indicators that can be used to inform marine  
23 management. Standard evaluation criteria (availability of data, quality of data, conceptual  
24 basis, communicability, relevancy to management) were implemented to identify and  
25 evaluate practical food-web indicators ready for operational use and indicators that hold  
26 reasonable promise for future use in policy and management. It was recognized that structure  
27 and functioning of food webs were the major attributes for which indicators were required  
28 and that resilience of food webs was a key aspect of ecosystem behavior and environmental  
29 status. Over 60 potential food-web indicators were evaluated and the final selection of  
30 operational food-web indicators includes: the primary production required to sustain a  
31 fishery, the productivity of seabirds (or similar charismatic megafauna), zooplankton  
32 indicators based on community biomass, size structure and productivity, integrated trophic  
33 indicators (including mean trophic level, mean size, etc.), and the biomass of trophic guilds.  
34 It was emphasised that more efforts should be made to determine suitable reference points in  
35 terms of threshold identification for achieving Good Environmental Status, as well as a  
36 greater level of integration in the development of indicators for international use.

### 371. Introduction

38Balancing the long-term maintenance of both biological diversity and human well-being is  
39key to sustainable resource management (e.g. Garcia et al., 2015, 2012; Link, 2010; Ostrom  
40et al., 1999; Pretty, 2003; Rockstrom et al., 2009). As such, ecosystem approaches to resource  
41management that address ecological and human interactions are an essential tool for  
42conservation. While there are number of differing definitions for Ecosystem-Based  
43Management (EBM), there is agreement about the need to move towards a more holistic  
44environmental management approach that recognizes the full array of interactions within an  
45ecosystem (Christensen et al., 1996; Link, 2010, 2005; McLeod et al., 2005). Currently,  
46activities stemming from EBM are used to support a number of management actions in  
47multiple ecosystems. In terrestrial habitats, EBM has been applied to management a number  
48of times (e.g. Caldwell, 1970; Slocombe, 1998, 1993) and localized EBM efforts for shallow  
49coastal habitats have also been undertaken (Kershner et al., 2011; Tallis et al., 2010).  
50Globally, a push for EBM in marine ecosystems has been made to balance the trade-offs  
51inherent in managing these complex ecosystems (Link, 2010). For example, EBM is central  
52to NOAA's Integrated Ecosystem Assessments (IEAs: Levin et al., 2009), Fisheries and  
53Oceans Canada has implemented aspects of EBM in the Canada Oceans Act (Curran et al.,  
542012), there has been a strong shift towards EBM in Australian fisheries driven by a number  
55of policy directions and initiatives (Smith et al., 2007), the European Union's Marine  
56Strategy Framework Directive (MSFD) has developed an overarching plan to reach and  
57maintain Good Environmental Status (GES;Rogers et al., 2010) and EBM is the recognized  
58mechanism to implement the Convention on the Conservation of Antarctic Living Marine  
59Resources (Constable, 2011; Constable et al., 2000). Thus there is a diverse and widespread  
60effort to continue to better manage marine ecosystems by taking into account all pressures,  
61responses and dynamics simultaneously.

62 Many aspects of ecosystem dynamics are reflected in food webs, the networks formed  
63by the trophic interactions between species in ecological communities. Historically, food  
64web studies developed from simply recording biological data through to a phase where  
65patterns in the data were identified and catalogued. Much work has since focused on  
66interpreting data and patterns, using either phenomenological or mechanistic models in food  
67webs (Rossberg, 2012). Among representations of food webs in the literature are simple  
68directed graphs (topological webs), flow diagrams (energy budgets), representations  
69aggregated by size or trophic level, and complex dynamic models (Link et al., 2005; Piroddi  
70et al., 2015). Depending on the representation, different structural and dynamic properties of  
71food webs emerge from the data. The relationships between these emergent patterns are the  
72subjects of much ongoing research (de Ruiter et al., 2005; Link et al., 2015; Rossberg, 2013).

73 Ecological indicators are important to EBM because they serve as proxies for several  
74ecological processes (e.g. growth dynamics, energy flow) and are representations of  
75ecosystem state (e.g. biodiversity, resilience). In particular, food-web indicators have become  
76increasingly important as they represent ecosystem services about which policy makers and  
77stakeholders are concerned. The global uses of these indicators to better inform management  
78of living marine resources has continued to increase over time (Coll et al., 2008; Fay et al.,  
792013; Jackson et al., 2001; Large et al., 2013; Large et al., 2015a; Levin et al., 2009). By  
80addressing much of the inherent complexity of marine ecosystems, food-web indicators are  
81one of the primary interfaces between policy and science. A critical step in the policy process  
82is to agree on food-web indicators that are compelling, intuitive, understandable and  
83defensible to all stakeholders, but also capture key food-web states and processes that  
84underlie critical and complex ecosystem dynamics. Important instances of such indicators are  
85those addressing emergent properties of food webs, which can be predicted without  
86understanding in detail the intricate processes operating in these complex systems. This

87predictability is reflected in the existence of simplified models or representations of food  
88webs addressing specific emergent properties (ICES, 2013a; Rossberg, 2013). Examples are  
89representations of food webs as food chains passing energy and biomass from lower to higher  
90trophic levels, representations in form of dynamically interacting aggregated groups of  
91species, representations as graphs with arrows (feeding interaction) linking nodes (species),  
92where a small number of top predators are supported by increasing numbers of species at  
93lower-trophic levels (de Ruiter et al., 2005) or complementarily, representation of the  
94distribution of community biomass over body sizes (Kerr and Dickie, 2001). It is important to  
95take into account these properties in selecting food-web indicators to develop pragmatic  
96indicators applicable to describe ecosystems at regional or larger scales.

97       For operational use, primary requirements are that food-web (or for that matter, any)  
98indicators be sensitive, have a basis in theory and be measurable (Dale and Beyeler, 2001;  
99Kershner et al., 2011; Link, 2010; Rice and Rochet, 2005a). Those indicators that are well  
100studied and linked with emergent properties can address cumulative impacts, integrated  
101dynamic responses, detect indirect and unintended consequences and can help to evaluate  
102trade-offs in managing ecosystems. Globally, a set of best-practices is coalescing around  
103indicator selection: a plethora of indicator selection criteria have been developed that identify  
104key facets of indicators (Fulton et al., 2005; Garcia et al., 2000; Greenstreet and Rogers,  
1052006; Greenstreet et al., 2011; ICES, 2013a, 2013b; Institute for European Environmental  
106Policy (IEEP), 2005; Link, 2005; Methratta and Link, 2006a; Piet and Jennings, 2005; Rice  
107and Rochet, 2005b; Rochet and Rice, 2005; Shin and Shannon, 2009; Shin et al., 2010a,  
1082010b)

109       While there have been some efforts to develop operational ecological indicators to  
110evaluate ecosystem status (ICES, 2015), the task of selecting specific food-web indicators has  
111been difficult for a number of reasons. Food-web ecology is a relatively new area of research

112(compared to more established community ecology and population ecology) with rapidly  
113emerging information and methods (Link et al., 2015; Longo et al., 2015; Thompson et al.,  
1142012). In light of new methodologies, historical data is often unsuitable to calculate the  
115necessary metrics to use potential food-web indicators for evaluating ecosystem status. Like  
116many other types of ecological indicators, selection of a specific set of food-web indicators  
117can imply that some aspects of marine food webs are valued more than others. Therefore, a  
118well-balanced selection process for indicators is required that encompasses all currently  
119known properties of marine food webs with the necessary data to be confidently used by both  
120management and stakeholders.

121 This study aims to provide a list of operational food-web indicators that can be used to  
122quantify the emergent properties of food webs in marine ecosystems. The context for this  
123work was the EU's MSFD need to delineate GES with regards to food webs (Descriptor 4;  
124ICES, 2014; Rogers et al., 2010), but was conducted cognizant of broader potential  
125applications to assess ocean status. Here, we develop a strategy using the best available  
126knowledge from scientific experts and a quantitative methodology for evaluating food-web  
127indicators for implementation in EBM. We also discuss the future development of these  
128indicators for practical use as reference points in management.

## 1292. **Methods**

130To address ongoing global requirements (Europe, North America and elsewhere), three  
131objectives related to food-web indicators were explored:

- 132 • To determine a defined process for selecting and developing food-web  
133 indicators.
- 134 • To develop a short list of suggested food-web indicators related to  
135 management contexts (EBM) in Europe and globally.

136 • To establish future direction for operationalizing food-web indicators.

137 This approach led to a two-part set of efforts to a) identify and evaluate operational food-web  
138 indicators that can currently be used and b) identify food-web indicators that hold promise in  
139 the future for management, but that require further development. This guidance would allow  
140 for increased clarity in selecting food-web indicators coherently within and across regions  
141 and lead to more defined response and pressure targets for control rules in EBM. As part of  
142 this broader effort, this project was developed as part of the ICES workshop to develop food-  
143 web indicators for operational use in EBM (ICES, 2014). The workshop brought together  
144 international experts in food webs, marine ecology and management to identify appropriate  
145 food-web indicators for current use.

#### 146 2.1. *Food-web indicators*

147 An initial set of 40 food-web indicators were selected from a list of over 60 candidate  
148 indicators presented by the workshop experts. Presentations covered all marine functional  
149 groups and all attributes of food webs that were considered necessary for a comprehensive  
150 evaluation. Duplicate and technically inappropriate indicators were eliminated from the pool  
151 of candidate indicators. The remaining 40 food-web indicators were grouped depending on  
152 three main food-web attributes which they addressed: functional indicators linked to energy  
153 flow, functional indicators linked to ecosystem resilience and structural indicators linked to  
154 diversity and ‘canary’ species (for more detailed descriptions see Appendix A).

#### 155 2.2. *Selection criteria*

156 A list of 5 criteria and 13 sub-criteria (Table 1) was initially synthesized from a set of criteria  
157 determined by previous working groups of experts examining ecological indicators (ICES,  
158 2015; Kershner et al. 2011). These criteria were adapted to broadly examine the functionality

159of the food-web indicators that could be operational within the global context (useful for  
160several countries and regions).

161 Each indicator was evaluated against the selection criteria and scored as 0, 1 or 2, where 0  
162= not met, 1 = partly met, and 2 = fully met. A Delphi method was used whereby sets of  
163indicators were scored by small groups based on consensus, following a discussion  
164establishing common understanding of the indicators themselves and how to apply the criteria  
165to the indicators. Each of the 13 sub-criteria was scored equally and no weighting was  
166applied. Scores were presented as percentages of the total score available (maximum score by  
167the number of categories; i.e.  $2 \times 13 = 26$ ). Indicators were ranked within the agreed  
168attributes of food webs (Functioning – energy flows, Resilience - ability to recover from  
169perturbation, Structure - species organization). Particular issues or concerns with individual  
170scores were highlighted for subsequent discussions. These were then examined so that all  
171scores were adjusted through consensus-based discussions. This process was used to quantify  
172the usefulness of indicators and to aid in the final selection.

### 1732.3. *Wider consideration for selecting food-web indicators*

174In addition to the specific criteria for each food-web indicator, a broader set of features was  
175considered through consensus of the experts involved when evaluating the final  
176recommended suite of indicators. The indicators were categorized into two groups, one set  
177that may be currently implemented and one that holds promise for future development. Key  
178considerations were:

179**Relative ranks** within the major food-web indicator attributes informed the choice of  
180indicators, but were not adhered to in a strictly quantitative manner.



181 **Coverage of all attributes** of food webs. To the extent practicable, all three main categories  
182 of food-web indicator attributes were represented.

183 **Coverage of all functional groups** found within a food web. Recognizing that much  
184 indicator development has occurred for upper trophic level contexts, we ensured that lower  
185 trophic level taxa were not omitted, even though as a group they may have scored lower than  
186 more commonly or routinely monitored upper trophic levels.

187 **Major indicator classes (structure, function and resilience)** were as well represented as  
188 possible to ensure that important facets of food webs were included.

189 **Current operability** was effectively based on an *ad hoc* review (or weighting) of operability  
190 issues related to data availability, management relevance and existence of baselines, targets  
191 or related reference points, which although were selection criteria, were deemed critical  
192 enough to warrant additional consideration.

193 **Links to other indicator uses** were considered to ensure that we emphasized food-web  
194 indicators that are unique to describing food webs. Other indicator uses include biodiversity,  
195 fisheries, eutrophication and sea floor integrity.

### 1963. Results

197 Within each attribute, indicators tended to cluster into groups with similar underlying  
198 ecological theory. When selecting priority indicators for further development it was therefore  
199 considered necessary to review the full list of indicators and ensure that those that clustered  
200 together, but with lower scores, were also taken into consideration to maintain a diversity of  
201 indicator formulations.

202 The rank scores were obtained from the unweighted sum of all 13 evaluation sub-criteria  
203 (Table 2a, b). When the evaluation was re-run separately using only the first six sub-criteria

204 in Table 1 (linked to practical aspects of indicator measurement), and the next seven criteria  
205 (linked to aspects of indicator implementation), there was relatively little difference in the  
206 final overall outcome. This suggests that the rank scores were robust to variability in criteria  
207 selection and were minimally influenced by single criteria evaluations.

### 208 3.1. Energy flow indicators

209 A relatively large number of indicators were identified which had clear links to functional  
210 aspects of food webs (Table 2a). Production or biomass ratios for various parts of the food  
211 web detect gross structural changes in the energy flow through a food web which may have  
212 been caused by, for example, harvesting of key species, or disruption of distributional overlap  
213 between predators and prey through climatic factors.

214 Total Mortality  $Z$  (Fishing mortality + natural mortality or production to biomass  
215 ratio), is commonly used in the ecosystem modelling community (Ecopath with Ecosim:  
216 Christensen and Pauly, 2008; Pauly et al., 2000). Despite the relatively high score this was  
217 not the most easily interpretable indicator of food web functioning. This was evident in the  
218 low score for the communication criteria (Table 2b). Ecosystem exploitation was considered  
219 useful to describe the harvesting pattern of exploited ecosystems. It is an indicator of the  
220 pressure of the fisheries on the food web.

221 Primary Production Required (PPR) to sustain a fishery has a solid conceptual basis  
222 (Pauly and Christensen, 1995). However, the difficulty of explaining the concept to the lay  
223 public contributed to a moderate score for this indicator. Moreover, this indicator does  
224 require estimates of transfer efficiency (TE), which is generally assumed to be 10-15%  
225 between trophic levels. Note that indicators of transfer efficiency themselves were not  
226 selected as indicators for use immediately due to lack of data to systematically estimate TE.  
227 Monitoring intermediate marine productivity and chlorophyll  $a$  fronts by satellite using

228remote observation was considered effective to estimate indicators of energy-flow in food  
229webs.

230 Four fairly similar indicators based on trophic level were evaluated (the mean trophic  
231level of the catch, the mean trophic index of the fish community, the mean trophic level of  
232the community and the trophic balance index). Each has a slightly different formulation, but  
233all require good quality and regularly updated data on dietary relationships, time series of  
234survey catch or landings from broad regional seas to avoid local population or fleet effects,  
235and accurate, agreed upon and regularly updated assessments of the trophic levels of the  
236ingested food. Similarly the Trophic Balance Index, describing the fishing pattern of local  
237métiers, can be useful in the context of assessing food web effects of fisheries harvesting, but  
238has limited application for other pressures.

239 Low scores allocated to indicators such as the disturbance index, loss in production  
240index, mean transfer efficiency and Finn Cycling Index were due to uncertainty over the  
241quality of the technical assessment (data needs and rigor) and the likely ease of  
242implementation. However, some of the indicators may warrant further investigation.

### 2433.2. *Resilience indicators*

244It was interesting to note that the six indicators that had a link to resilience of the food web  
245were generally scored lower than many other indicators (Table 2b). This may be because they  
246are more conceptually complex. It was considered that the top three in this category, the  
247Mean number of trophic links per species, Ecological Network Analysis derived indicators,  
248and the Gini-Simpson dietary diversity index, all held promise as food-web indicators, but the  
249group of experts felt that these would not be recommended as suitable for implementation in  
250the short-term. The conceptual and technical difficulty of measuring food-web resilience and  
251ability to recover from perturbation partly explains the low scores allocated to the assessment

252 criteria in the area of cost-effectiveness of data gathering, although they all have strong  
253 support in the literature.

254 The indicators for this attribute that scored poorly (Herbivory: Detritivory Ratio,  
255 Ecological Network Indices, System Omnivory Indices) will take more time to develop. The  
256 complexity of their formulation also suggests that, even if further developed, they may be  
257 difficult to explain in a management context. More importantly, these indicators need regular  
258 diet time series data encompassing the entire food web, which have not been made widely  
259 available even to support applied multispecies fishery assessments.

### 260 3.3. *Structural indicators*

261 Several indicators in this category obtained relatively high scores, suggesting that managers  
262 may want to use these indicators to help interpret patterns observed particularly at higher  
263 trophic levels. Another important consideration is the role of aggregated sets of structural  
264 indicators, such as those related to phytoplankton, zooplankton, forage fish, scavengers and  
265 birds, which together have important implications for food-web resilience (e.g. low or high  
266 biodiversity) as well as structure of the individual components. Many structural indicators are  
267 describing the same ecosystem components in multiple ways (Table 2a, b) and due to the  
268 multi-faceted uses of these indicators (in addition to characterizing food webs) the data are  
269 likely to be collected and available.

270 Higher-scoring indicators were those which informed trends in absolute biomass,  
271 production, or ratios of both, for a number of guild-level ecosystem components, especially  
272 higher predators. For those structural indicators that aggregate across multiple components, it  
273 was generally thought preferable to have indicators comprising absolute values rather than  
274 ratios, as these data would be necessary anyway to interpret ratio metrics. Some of these  
275 abundance-related indicators may be given a higher priority if they are also useful for

276informing an aspect of food web resilience. For example, both the Gini-Simpson diversity  
277indices for small and large fish and the Species Richness Index were thought to be potentially  
278useful for assessing food web resilience.

#### 2793.4. *Suggested food-web indicators*

280The following indicators are the refined set of food-web indicators recommended for current  
281use based on the selection criteria (Table 1) and accounting for the wider considerations in  
282the selection process (Table 2a,b):

#### 283 **Guild level biomass (and production)**

284Guild-level biomasses and production address structural attributes of food webs, and can also  
285serve as proxies for functioning. It was noted that the typical use of this type of indicator has  
286been for fishes, but if feasible this indicator should include multiple guilds across all trophic  
287levels, such as primary producers, zooplankton, benthos, and charismatic megafauna, beyond  
288just fish or upper trophic levels. The guilds should be determined as appropriate for the taxa in  
289a given regional sea.

#### 290 **Primary Production Required to sustain a fishery (PPR)**

291This addresses the functioning attribute of food webs and is a measure of the ecological  
292footprint of a fishery. However, this metric can (and often does) integrate a wide range of  
293removals from the food web. Derivatives of this food-web indicator could, where feasible, be  
294contrasted to measures of primary production to ensure it is directly appraised against field  
295data. Satellite imagery makes estimates of primary production widely available (given the  
296usual caveats of remotely sensed data), and typical landings and associated data are also  
297widely available, making PPR more integrative and feasible than is often perceived.

298        **Seabird (charismatic megafauna) productivity**

299 The breeding success of seabirds addresses the structural and functional attribute of a food  
300 web and can also serve as a proxy for resilience. Although particular to seabirds, especially  
301 breeding success/chicks per pair, it was recognized that seabirds may not be prominent or  
302 important in all regional seas. Similar productivity indicator could be calculated for marine  
303 mammal taxa (i.e. pup production rates).

304        **Zooplankton size biomass index**

305 This indicator addresses both structural and functional attributes of food webs. Although  
306 indicators associated with this taxonomic group were often ranked lower, they represent an  
307 important part of the food web - the link between primary production at lower trophic level  
308 and upper trophic level consumption and growth.

309        **Integrated trophic indicators (mean TL, mean size)**

310 Trophic indicators address both structural and resilience attributes of food webs. It was  
311 critical to include an explicitly integrative measure that provided some view of the overall  
312 system and did not focus on only certain facets of it. There are many possible indicators in  
313 this category from which to choose, such as mean trophic level, mean, or proportion at size of  
314 the community (depending upon abundance) and trophic data availability in a given regional  
315 sea.

316 3.5.    *Indicators for development*

317 Food-web indicators that were recommended for future development were Ecological  
318 Network Analysis indicators, the Gini-Simpson dietary diversity index and condition  
319 indicators. These indicators lacked the available data to be considered currently useful for  
320 management, but all were determined to be representative of multiple aspects of the food-web

321(integrated food-web perspective). Some indicators that were suggested to be currently  
322operational (marine trophic level indicators, primary producers and zooplankton indicators)  
323were also thought to require more development to fully meet their potential and range as  
324indicators for food-web and other indicator uses.

#### 3254. Discussion

326 The five food-web indicators recommended from this process cover important facets of  
327food webs, particularly addressing structural, functional and resilient features of marine food  
328webs (Jennings and Collingridge, 2015; Polis and Strong, 1996; Thompson et al., 2012). It is  
329likely that multiple indicators are needed to track the multiple features that comprise food  
330webs and delineation of GES (Large et al., 2015a, 2015b; Mallory et al., 2010; Rice and  
331Rochet, 2005a) of which these five candidates are suitable options. All of the five food-web  
332indicators proposed here are generally applicable in terms of capturing the main facets of  
333food-web dynamics (ICES, 2014; Methratta and Link, 2006b; Shannon et al., 2009) and  
334readily link to known behaviors of food webs. Many of these indicators are broad enough in  
335context to be applied across many marine ecosystems (coastal, temperate, arctic, tropical,  
336etc.; Andrews et al., 2013; Coll and Libralato, 2012; Fulton et al., 2005; Hayes et al., 2015;  
337Parsons et al., 2008; Zador et al., 2014).

338 Yet even the five proposed indicators may not all have widely and consistently  
339monitored data available to sufficiently calculate the metrics. Although important to track  
340lower-trophic level dynamics and linkages to upper-trophic level taxa, the zooplankton  
341indicator may not have widely collected data nor be as easily interpreted, given the high  
342seasonality of these taxa (Pershing et al., 2005; Stige et al., 2014; Vargas et al., 2006). The  
343integrated trophic indicators hold equal promise, but similarly may not always have measures  
344of trophic level or equivalent (TL; Gaichas et al., 2012; Hornborg et al., 2013; Pranovi et al.,  
3452012; Rossberg et al., 2006). Justifiable assumptions regarding TL, using common databases

346 on trophic ecology of taxa (e.g. fishbase; Froese and Pauly, 2013; Froese, 1992), may provide  
347 a means to more readily calculate these indicators in the absence of local trophic data. Size-  
348 based integrated indicators are an type of indicator that are less demanding on data and has  
349 been found to show clearer responses in food webs (Engelhard et al., 2015; Fung et al., 2013;  
350 Greenstreet et al., 2011; Shephard et al., 2011) The salient point is that there are well-studied  
351 extant indicators able to track and delineate environmental status in marine food webs (Houle  
352 et al., 2012). These were explored in the MSFD GES context (ICES, 2015, 2013b, 2008;  
353 Shephard et al., 2014), but are generally applicable for marine conservation considerations.

354     Regardless of the specific indicator set chosen, a replicable, transparent, defensible and  
355 clear process for selection is required (Dale and Beyeler, 2001; Link, 2010; Shin et al.,  
356 2010a). The process demonstrated here is broadly applicable in a wide array of conservation  
357 situations and it is as important as the outcomes. It is essentially a multi-criteria decision  
358 analysis (Mendoza and Martins, 2006), whereupon the selection of indicators is agreed-to  
359 before use in tracking ecosystem status. The criteria for indicator assessment used here are  
360 sufficiently robust to be applied in a range of situations, with one of the five main criteria  
361 specifically evaluating how useful a given indicator is to management. These criteria are  
362 converging in the marine management context, but can be readily used in other forms of  
363 natural resource management (e.g. terrestrial, estuarine). Due to the well-documented  
364 quantitative and qualitative evaluation in the selection process, there is a high level of  
365 confidence in the choice of the final set of indicators. This process allows for regular updates  
366 and inclusion of novel information (Curtin and Prellezo, 2010; Kershner et al., 2011) while  
367 maintaining a record of how selections are made. This process is general enough to be used  
368 regardless of the type of ecosystem and conservation issue being considered, as long as the  
369 criteria are agreed upon *a priori* (Espinosa-Romero et al., 2011; Martin et al., 2009).



370 Although similar selection processes have a wide history of use in conservation (Mendoza  
371 and Martins, 2006), it could be even more widely and rigorously applied.

372 Based on the evaluation process, the food-web indicators selected in this study can offer  
373 some guidance towards possible management actions. Guild-level biomass reflects measures  
374 of biodiversity and structural relationships within ecosystems (Garrison and Link, 2000;  
375 Rosenfeld, 2002). It can be an integrative indicator to evaluate the status of a particular guild  
376 group in relation to another. For instance, lower numbers of forage fish will have direct (and  
377 indirect) impacts on larger predators and seabirds (Cury et al., 2011; Garrison and Link,  
378 2000) or could indicate low levels of primary productivity (Jennings and Collingridge, 2015;  
379 Polivina et al., 2001). Either way, management responses to maintain forage fish could be  
380 identified based on the information that guild-level indicators provide (Heath et al., 2014).  
381 Similarly, integrated trophic indicators can address multiple aspects of structure, function and  
382 resilience in ecosystems, where lower mean size or trophic level indicate impacts on large,  
383 predatory animals (Methratta and Link, 2006b; Pauly and Watson, 2005; Rosenfeld, 2002).  
384 Specific fisheries management actions with respect to changes in these indicators over time  
385 could include adjusting harvest control rules for particularly overexploited guilds, but could  
386 also include concentrating fishing efforts on lower age or size groups (Anderson et al., 2008).  
387 Both higher-trophic (seabird and charismatic megafauna productivity) and lower-trophic  
388 indicators (PPR and zooplankton index) are reflective of bottom-up processes viewed from  
389 opposing ends of the food web (Cury et al., 2011; Einoder, 2009; Hilting et al., 2013). PPR is  
390 an integrative indicator that represents the amount of primary productivity to sustain a  
391 fishery, and offers a means to compare energy requirements across different fisheries  
392 (Chassot et al., 2010; Gascuel et al., 2005). Seabird productivity is an indicator of food  
393 availability (forage fish) and can also be sensitive to contaminants and environmental  
394 pollutants (Mallory et al., 2010). Direct management actions to influence these indicators

395 could be either top-down control rules aimed at relieving fishing pressure on lower-trophic  
396 species or bottom-up policies directed to improve water quality or habitat, which may also  
397 include improved management at land-sea interfaces (Furness and Camphuysen, 1997;  
398 Kendall et al., 2010; King and Baker, 2010; Mallory et al., 2010; Teichert et al., 2015).  
399 Specific management actions will be dependent on regional circumstances and the responses  
400 of the indicators to local pressures, but by using common indicators it will be possible to  
401 compare ecosystem status between regions and to help management at all levels (from  
402 regional to national to international) and to make effective decisions to improve the world's  
403 oceans.

404       This proposed set of candidate indicators is a start towards operationalizing the  
405 delineation of marine ecosystem status, but may require a few further steps before becoming  
406 fully operational. Food-web indicators may be interesting scientifically and relevant for  
407 management, but if they cannot inform management actions directly they certainly have less  
408 utility. Establishing decision criteria that trigger management actions for EBM requires an  
409 understanding of how pressure variables influence indicators, as well as the level of a  
410 particular pressure at which significant changes in ecosystem structure or function appear  
411 (Blanchard et al., 2010; Coll et al., 2010; Groffman et al., 2006; Link, 2010, 2002a; Samhuri  
412 et al., 2010). Such thresholds have been explored with a wide range of analytical methods,  
413 such as cumulative sums (CUSUM; Hinkley, 1970), sequential t-test (STARS; Rodinov,  
414 2004), empirical fluctuation processes (Zeileis and Kleiber, 2005), and significant zero  
415 crossings of piecewise regression models (Chaudhuri and Marron, 1999; Samhuri et al.,  
416 2012, 2010; Sonderegger et al., 2008; Toms and Lesperance, 2003; Toms and Villard, 2015)  
417 or generalized additive models (Large et al., 2013), all to identify the level of pressure that  
418 results in a significant indicator response (Andersen et al., 2009). These univariate  
419 relationships are useful for establishing decision criteria (Fay et al., 2013; Large et al., 2013;

420 Samhuri et al., 2010), however, they do not fully account for multiple pressures that likely  
421 interact and occur concurrently. An assessment of ecosystem status based on suites of  
422 indicators will be more powerful. Using multiple indicators to evaluate ecosystems will help  
423 to avoid the possibility of misinterpretation which can occur when indicators are evaluated in  
424 isolation (Coll and Libralato, 2012; Longo et al., 2015; Rice and Rochet, 2005b). Multivariate  
425 approaches exist to detect thresholds, including translating indicator response into a surface  
426 dependent on multiple pressures (i.e., fishing and environmental pressure; Frederiksen et al.,  
427 2007; Large et al., 2015a; Scott et al., 2006), multivariate ordination methods (Baker and  
428 King, 2010; King and Baker, 2010) and extensions of regression tree and gradient forest  
429 analyses (Baker and Hollowed, 2014; Ellis et al., 2008; Large et al., 2015b; Liaw and Wiener,  
430 2002; Pitcher et al., 2012; Prasad et al., 2006). Understanding how multiple pressure  
431 variables concurrently influence ecosystem status, as evinced by thresholds in indicators, will  
432 help to further operationalize these indicators as reference points for management.

433       Both the EU MSFD and US IEA efforts have a similar framework that includes  
434 indicators as a critical part of the management decision-making process (Andrews et al.,  
435 2013; Levin et al., 2009; Rogers et al., 2010; Shephard et al., 2014). Currently, many of the  
436 efforts from both the MSFD and IEA frameworks assess ecosystems by using a suite of  
437 indicators. Despite the frequent absence of thresholds of indicators to establish reference  
438 points, there is still sufficient information and examples of using such indicators to inform  
439 marine ecosystem management advice. Even qualitative and directional features of indicators  
440 can and have been used operationally (Andrews et al., 2013; Espinosa-Romero et al., 2011;  
441 Foley et al., 2015; Greenstreet et al., 2012; Large et al., 2015a, 2015b; Link et al., 2015;  
442 Longo et al., 2015; Samhuri et al., 2012, 2010; Zador et al., 2014). Thus, the monitoring,  
443 tracking and presentation of the food-web indicators proposed here can help to operationally  
444 delineate GES.

445           When assessing the status of marine ecosystems, it is important to adequately  
446characterize the food web (Branch et al., 2010; Link, 2002b; Thompson et al., 2012).  
447Certainly there are other aspects of marine ecosystem status, a fact which is explicitly  
448acknowledged in the MSFD. Yet, too often the development of marine indicators neglect to  
449consider food webs (Hayes et al., 2015). Understanding food webs in ecosystems is  
450paramount because they are able to unify ecological sub-disciplines (behavior, dispersal,  
451physiology, thermodynamics etc.) and to examine interactions among guilds (Polis and  
452Strong, 1996; Rossberg, 2013; Thompson et al., 2012). Food webs are able to integrate  
453species-based and functional-based approaches to examine biomass distributions and  
454energetic flows within systems. Another key aspect of ecosystems that is encompassed by  
455food webs is resilience. A resilient system reacts only weakly to pressure, but resilience  
456might be lost with increasing pressures, leading to rapid changes to different states or  
457regimes. Such transition is thus the result of an accumulation of the disturbing effects of  
458pressures (Folke et al., 2004; Gunderson, 2000; Sasaki et al., 2015). Additionally,  
459ecosystems may exhibit legacy effects of earlier pressures (Folke, 2006; Hughes et al., 2005).  
460Despite the difficulty in studying food webs in their entirety (including large data  
461requirements and advanced computational abilities), emergent trends have been established in  
462food-web ecology at both the community (Fredriksen, 2003; Neira et al., 2009) and  
463ecosystem level (Link et al., 2015).

464           An important aim of EBM is to balance between multiple, often conflicting  
465objectives. How management actions take shape depends on all user groups involved,  
466including stakeholders, indigenous communities, fishers, tourists, NGOs, etc. (Branch et al.,  
4672006; Link, 2010; Marasco et al., 2007). The most successful implementation of EBM will  
468one where user groups are equally engaged, can agree on a set objectives, work towards  
469common economic-social-conservation management goals and ultimately overcome inertia in

470the decision making process (Arkema et al., 2006; deReynier et al., 2010; Espinosa-Romero  
471et al., 2011; Leslie and McLeod, 2007; Link, 2010; Pitcher et al., 2009; Röckmann et al.,  
4722015; Sandström et al., 2015). The set of indicators proposed in this study is an example of  
473how such information can be used to more fully implement EBM by evaluating one facet of  
474marine ecosystem objectives associated with food webs. More so, the process described here  
475is an important means to explore the tradeoffs not only in selecting these indicators but also  
476the underlying objectives and dynamics that each represents.

477 Ecological indicators for conservation (including food-web indicators) are useful to  
478summarize complex information concerning marine ecosystem status (Cury and Christensen,  
4792005; Dulvy et al., 2006; Fulton et al., 2005; Hayes et al., 2015; ICES, 2015; Methratta and  
480Link, 2006b). Clearly defined, consistent metrics at the global scale can provide management  
481in multiple countries with the tools to make EBM more operational (Leslie and McLeod,  
4822007; Lester et al., 2010; Link, 2010; Link et al., 2011; Smith et al., 2007; Thrush and  
483Dayton, 2010). As management efforts continue to implement EBM to meet conservation  
484objectives, having a suite of indicators, a process to select them and ensuring that they map to  
485clear management needs will remain increasingly important.

486

#### 487**Acknowledgements**

488Many thanks to the participants of the WKfooWI (ICES CM 2014\ACOM:48): S.  
489Greenstreet, S. Libralato, N. Niquil, M. Tomczak, A. Belgrano, J-N. Druon, P. Levin, S.  
490Large, J. Lesutiene, M. Johansen, K. van de Wolfshaar, I. Preciado, J. Patricio, A. Paliaxis,  
491A. Bundy, P. Tett, F. Pranovi, E. Gorokhova, M-J. Rochet, M. Dickey-Collas, G. Johansen, J.  
492Houle, A. Rindorf. This work was supported by a NOAA postdoctoral fellowship to JCT.

## 493References

- 494 Andersen, T., Carstensen, J., Hernández-García, E., Duarte, C.M., 2009. Ecological  
495 thresholds and regime shifts: approaches to identification. *Trends Ecol. Evol.* 24, 49–57.  
496 doi:10.1016/j.tree.2008.07.014
- 497 Anderson, C.N.K., Hsieh, C., Sandin, S. a, Hewitt, R., Hollowed, A., Beddington, J., May,  
498 R.M., Sugihara, G., 2008. Why fishing magnifies fluctuations in fish abundance. *Nature*  
499 452, 835–9. doi:10.1038/nature06851
- 500 Andrews, K.S., Ballance, L.T., Barcelo, C., Barlow, J.P., Bellman, M.A., Steven, J., Brodeur,  
501 R.D., Brown, C.J., Chivers, S.J., Cope, J.M., Crone, P.R., Beukelaer, S. De, Dereynier,  
502 Y., Devogelaere, A., Dunsmore, R., Robert, L., Gertseva, V. V, Good, T.P., Gray, I.A.,  
503 Haltuch, M.A., Hamel, O.S., 2013. Integrated Ecosystem Assessment of the California  
504 Current.
- 505 Arkema, K.K., Abramson, S.C., Dewsbury, B.M., 2006. Marine Ecosystem-Based  
506 Management: From Characterization to Implementation. *Front. Ecol. Environ.* 4, 525–  
507 532.
- 508 Badalamenti, G., Anna, G.D., Pinnegar, J.K., Polunin, N.V.C., 2002. Size-related  
509 trophodynamic changes in three target fish species recovering from intensive trawlin.  
510 *Mar. Biol.* 141, 561–570.
- 511 Baker, M.E., King, R.S., 2010. A new method for detecting and interpreting biodiversity and  
512 ecological community thresholds. *Methods Ecol. Evol.* 1, 25–37. doi:10.1111/j.2041-  
513 210X.2009.00007.x
- 514 Baker, M.R., Hollowed, A.B., 2014. Delineating ecological regions in marine systems:  
515 Integrating physical structure and community composition to inform spatial  
516 management in the eastern Bering Sea. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 109,  
517 215–240. doi:10.1016/j.dsr2.2014.03.001
- 518 Belkin, I.M., Cornillon, P.C., Sherman, K., 2009. Fronts in large marine ecosystems. *Prog.*  
519 *Oceanogr.* 81, 233–236.
- 520 Blanchard, J.L., Coll, M., Trenkel, V.M., Vergnon, R., Yemane, D., Jouffre, D., Link, J.S.,  
521 Shin, Y.-J., 2010. Trend analysis of indicators: a comparison of recent changes in the  
522 status of marine ecosystems around the world. *ICES J. Mar. Sci.* 67, 732–744.  
523 doi:10.1093/icesjms/fsp282
- 524 Bondavalli, C., Ulanowicz, R.E., Bondini, A., 2000. Insights into the processing of carbon in  
525 the South Florida Cypress Wetlands: A whole-ecosystem approach to using network  
526 analysis. *J. Biogeogr.* 27, 697–710.
- 527 Boyd, H., Charles, A., 2006. Creating community-based indicators to monitor sustainability  
528 of local fisheries. *Ocean Coast. Manag.* 49, 237–258.  
529 doi:10.1016/j.ocecoaman.2006.03.006
- 530 Branch, T. a, Hilborn, R., Haynie, A.C., Fay, G., Flynn, L., Griffiths, J., Marshall, K.N.,  
531 Randall, J.K., Scheuerell, J.M., Ward, E.J., Young, M., 2006. Fleet dynamics and  
532 fishermen behavior: lessons for fisheries managers. *Can. J. Fish. Aquat. Sci.* 63, 1647–  
533 1668. doi:10.1139/f06-072
- 534 Branch, T.A., Watson, R., Fulton, E.A., Jennings, S., McGilliard, C.R., Pablico, G.T., Ricard,  
535 D., Tracey, S.R., 2010. The trophic fingerprint of marine fisheries. *Nature* 468, 431–5.  
536 doi:10.1038/nature09528

- 537Buckland, S.T., Studeny, A., Magurran, A.C., Illian, J.B., Newson, S.E., 2011. The geometric  
538 mean of relative abundance indices: a biodiversity measure with a difference. *Ecosphere*  
539 2, art100.
- 540Bundy, A., Fanning, P., Zwanenburg, K.C.T., 2005. Balancing exploitation and conservation  
541 of the eastern Scotian Shelf ecosystem: application of a 4D ecosystem exploitation  
542 index. *ICES J. Mar. Sci.* 62, 503–510.
- 543Bundy, A., Shannon, L.J., Rochet, M.-J., Neira, S., Shin, Y.-J., Hill, L., Aydin, K., 2010. The  
544 Good(ish), the Bad and the Ugly: a tripartite classification of ecosystem trends. *ICES J.*  
545 *Mar. Sci.* 67, 745–768.
- 546Burns, T.P., 1989. Lindeman's contradiction and the trophic structure of ecosystems. *Ecology*  
547 70, 1355–1362.
- 548Caldwell, L.K., 1970. Ecosystem as a criterion for public land policy. *Nat. Resour. J.* 10,  
549 203–221.
- 550Chassot, E., Bonhommeau, S., Dulvy, N.K., Melin, F., Watson, R., Gascuel, D., Le Pape, O.,  
551 2010. Global marine primary production constrains fisheries catches. *Ecol. Lett.* 13,  
552 495–500.
- 553Chaudhuri, P., Marron, J.S., 1999. SiZer for exploration of structures in curves. *J. Am. Stat.*  
554 *Assoc.* 94, 807–823.
- 555Christensen, N.L., Bartuska, A.M., Brown, J.H., Carpenter, S., Antonio, D., Francis, R.,  
556 Franklin, J.F., Macmahon, J.A., Noss, R.F., David, J., Peterson, C.H., Turner, M.G.,  
557 Woodmansee, R.G., 1996. The report of the Ecological Society of America Committee on  
558 the Scientific Basis for Ecosystem Management. *Ecol. Appl.* 6, 665–691.
- 559Christensen, V., 1995. Ecosystem maturity - Towards quantification. *Ecol. Modell.* 77, 3–32.
- 560Christensen, V., Pauly, D., 2008. *Ecopath with Ecosim: a user's guide.*
- 561Christensen, V., Pauly, D., 1993. *Trophic models of aquatic ecosystems*, 26th ed. Worldfish.
- 562Christensen, V., Walters, C.J., Ahrens, R., Alder, J., Buszowski, J., Bang, L., Cheung,  
563 W.W.L., Dunne, J., Froese, R., Karpouzi, V., Kaschner, K., Kearney, K., Lai, S., Lam,  
564 V., Palomares, M.L.D., Peters-mason, A., Piroddi, C., Sarmiento, J.L., Steenbeek, J.,  
565 Sumaila, R., Watson, R., Zeller, D., Pauly, D., 2009. Database-driven models of the  
566 world's Large Marine Ecosystems. *Ecol. Modell.* 220, 1984–1996.  
567 doi:10.1016/j.ecolmodel.2009.04.041
- 568Coll, M., Libralato, S., 2012. Contributions of food web modelling to the ecosystem approach  
569 to marine resource management in the Mediterranean Sea. *Fish Fish.* 13, 60–88.  
570 doi:10.1111/j.1467-2979.2011.00420.x
- 571Coll, M., Libralato, S., Tudela, S., Palomera, I., Pranovi, F., 2008. Ecosystem overfishing in  
572 the ocean. *PLoS One* 3, e3881. doi:10.1371/journal.pone.0003881
- 573Coll, M., Shannon, L.J., Yemane, D., Link, J.S., Ojaveer, H., Neira, S., Jouffre, D., Labrosse,  
574 P., Heymans, J.J., Fulton, E.A., Shin, Y.-J., 2010. Ranking the ecological relative status  
575 of exploited marine ecosystems. *ICES J. Mar. Sci.* 67, 769–786.
- 576Constable, A.J., 2011. Lessons from CCAMLR on the implementation of the ecosystem  
577 approach to managing fisheries. *Fish Fish.* 12, 138–151. doi:10.1111/j.1467-  
578 2979.2011.00410.x
- 579Constable, A.J., de la Mare, W.K., Agnew, D.J., Everson, I., Miller, D., 2000. Managing  
580 fisheries to conserve the Antarctic marine ecosystem: practical implementation of the

581 Convention on the Conservation of Antarctic Marine Living Resources (CCAMLR).  
582 ICES J. Mar. Sci. 57, 778–791. doi:10.1006/jmsc.2000.0725

583Curran, K., Bundy, A., Craig, M., Hall, T., Lawton, P., Quigley, S., 2012. Recommendations  
584 for Science, Management and an Ecosystem Approach to Fisheries and Oceans Canada,  
585 Maritimes Region. Can. Sci. Advis. Secr. Res. Doc. 2012/061, 48.

586Curtin, R., Prellezo, R., 2010. Understanding marine ecosystem based management: A  
587 literature review. Mar. Policy 34, 821–830. doi:10.1016/j.marpol.2010.01.003

588Cury, P., Boyd, I.L., Bonhommeau, S., Anker-Nilssen, T., Crawford, R.J., Furness, R.W.,  
589 Mills, J.A., Murphy, E.J., Osterblom, H., Paleczny, M., Piatt, J.F., Roux, J.-P., Shannon,  
590 L., Sydenman, W.J., 2011. Global Seabird response to forage fish depletion - one-third  
591 for the birds. Science (80-. ). 334, 1703–1706.

592Cury, P., Christensen, V., 2005. Quantitative ecosystem indicators for fisheries management.  
593 ICES J. Mar. Sci. 62, 307–310. doi:10.1016/j.icesjms.2005.02.003

594Cury, P., Shannon, L., Roux, J., Daskalov, G., Jarre, a, Moloney, C., Pauly, D., 2005.  
595 Trophodynamic indicators for an ecosystem approach to fisheries. ICES J. Mar. Sci. 62,  
596 430–442. doi:10.1016/j.icesjms.2004.12.006

597Daan, N., Gislason, H., Pope, J.G., Rice, J.C., 2005. Changes in the North Sea fish  
598 community: evidence of indirect effects of fishing. ICES J. Mar. Sci. 62, 177–188.

599Dale, V.H., Beyeler, S.C., 2001. Challenges in the development and use of ecological  
600 indicators. Ecol. Indic. 1, 3–10. doi:10.1016/S1470-160X(01)00003-6

601Dame, J.K., Christina, R.R., 2007. A statistical test of network analysis: can it detect  
602 differences in food web properties. Ecosystems 10, 906–923.

603de Leiva Moreno, J.I., Agostini, V.N., Caddy, J.F., Carocci, F., 2000. Is the pelagic-demersal  
604 ratio from fishery landings a useful proxy for nutrient availability? A preliminary data  
605 exploration for the semi-enclosed seas around Europe. ICES J. Mar. Sci. 57, 1091–1102.

606de Ruiter, P.C., Wolters, V., Moore, J., Winemiller, K.O., 2005. Food Web Ecology: Playing  
607 Jenga and Beyond. Science (80-. ). 309, 68–71.

608deReynier, Y.L., Levin, P.S., Shoji, N.L., 2010. Bringing stakeholders, scientists, and  
609 managers together through an integrated ecosystem assessment process. Mar. Policy 34,  
610 534–540. doi:10.1016/j.marpol.2009.10.010

611Druon, J.N., 2010. Habitat mapping of the Atlantic bluefin tuna derived from satellite data:  
612 its potential as a tool for the sustainable management of fisheries. Mar. Policy 34, 293–  
613 297.

614Druon, J.N., Fromentin, J.M., Aulanier, F., Heikkonen, J., 2011. Potential feeding and  
615 spawning habitat of Atlantic bluefin tuna in the Mediterranean Sea. Mar. Ecol. Prog.  
616 Ser. 439, 223–240.

617Druon, J.N., Panigada, S., David, L., Gannier, A., Mayol, P., Arcangeli, A., Canadas, A.,  
618 Laran, S., Di Meglio, N., Gauffier, P., 2012. Potential feeding habitat of fin whales in  
619 the western Mediterranean Sea: an environmentla niche model. Mar. Ecol. Prog. Ser.  
620 464, 289–306.

621Dulvy, N.K., Jennings, S., Rogers, S.I., Maxwell, D.L., 2006. Threat and decline in fishes: an  
622 indicator of marine biodiversity. Can. J. Fish. Aquat. Sci. 63, 1267–1275.  
623 doi:10.1139/f06-035

624Dulvy, N.K., Polunin, N.V.C., Mill, A.C., Graham, N.A.J., 2004. Size structural change in



- 625 lightly exploited coral reef fish communities: evidence for weak indirect effects. *Can. J.*  
626 *Fish. Aquat. Sci.* 61, 466–475. doi:10.1139/f03-169
- 627Einoder, L.D., 2009. A review of the use of seabirds as indicators in fisheries and ecosystem  
628 management. *Fish. Res.* 95, 6–13. doi:10.1016/j.fishres.2008.09.024
- 629Ellis, N., Pantus, F., Welna, A., Butler, A., 2008. Evaluating ecosystem-based management  
630 options: Effects of trawling in Torres Strait, Australia. *Cont. Shelf Res.* 28, 2324–2338.  
631 doi:10.1016/j.csr.2008.03.031
- 632Engelhard, G.H., Lynam, C.P., García-Carreras, B., Dolder, P.J., Mackinson, S., 2015. Effort  
633 reduction and the large fish indicator: spatial trends reveal positive impacts of recent  
634 European fleet reduction schemes. *Environ. Conserv.* 42, 227–236.  
635 doi:10.1017/S0376892915000077
- 636Espinosa-Romero, M.J., Chan, K.M. a., McDaniels, T., Dalmer, D.M., 2011. Structuring  
637 decision-making for ecosystem-based management. *Mar. Policy* 35, 575–583.  
638 doi:10.1016/j.marpol.2011.01.019
- 639EU, 2010. Comission decision of 18 December 2009 adopting multiannual community  
640 programme for the collection, management and use of data in the fisheries sector for the  
641 period 2011-2013 (2010/93/EU).
- 642Fasola, M., Bogliani, G., Saino, N., Canova, L., 1989. Foraging, feeding and time-activity  
643 niches of eight species of breeding seabirds in the coastal wetlands of the Adriatic Sea.  
644 *Ital. J. Zool.* 56, 61–72.
- 645Fay, G., Large, S.I., Link, J.S., Gamble, R.J., 2013. Testing systemic fishing responses with  
646 ecosystem indicators. *Ecol. Modell.* 265, 45–55. doi:10.1016/j.ecolmodel.2013.05.016
- 647Finn, J., 1976. Measures of structure and functioning derived from analysis of flows. *J.*  
648 *Theor. Biol.* 56, 363–380.
- 649Foley, M.M., Martone, R.G., Fox, M.D., Kappel, C. V., Mease, L. a., Erickson, A.L.,  
650 Halpern, B.S., Selkoe, K. a., Taylor, P., Scarborough, C., 2015. Using Ecological  
651 Thresholds to Inform Resource Management: Current Options and Future Possibilities.  
652 *Front. Mar. Sci.* 2, 1–12. doi:10.3389/fmars.2015.00095
- 653Folke, C., 2006. Resilience: The emergence of a perspective for social–ecological systems  
654 analyses. *Glob. Environ. Chang.* 16, 253–267. doi:10.1016/j.gloenvcha.2006.04.002
- 655Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L., Holling, C.S.,  
656 2004. Regime Shifts, Resilience, and Biodiversity in Ecosystem Management. *Annu.*  
657 *Rev. Ecol. Evol. Syst.* 35, 557–581. doi:10.2307/annurev.ecolsys.35.021103.30000021
- 658Frederiksen, M., Furness, R., Wanless, S., 2007. Regional variation in the role of bottom-up  
659 and top-down processes in controlling sandeel abundance in the North Sea. *Mar. Ecol.*  
660 *Prog. Ser.* 337, 279–286.
- 661Fredriksen, S., 2003. Food web studies in a Norwegian kelp forest based on stable isotope  
662 ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) analysis. *Mar. Ecol. Prog. Ser.* 260, 71–81.
- 663Froese, R., 1992. Progress Report on FishBase. ICES Council. Meet. 852, 1–6.
- 664Froese, R., Pauly, D., 2013. FishBase [WWW Document]. URL [www.fishbase.org](http://www.fishbase.org)
- 665Fuchs, H., Franks, P.J.S., 2010. Plankton community properties determined by nutrients and  
666 size-selective feeding. *Mar. Ecol. Prog. Ser.* 413, 1–15.
- 667Fulton, E., Smith, A., Punt, A.E., 2005. Which ecological indicators can robustly detect  
668 effects of fishing? *ICES J. Mar. Sci.* 62, 540–551. doi:10.1016/j.icesjms.2004.12.012

- 669 Fung, T., Farnsworth, K.D., Reid, D.G., Rossberg, A.G., 2015. Impact of biodiversity loss on  
670 production in complex marine food webs mitigated by prey-release. *Nat. Commun.* 6,  
671 6657. doi:10.1038/ncomms7657
- 672 Fung, T., Farnsworth, K.D., Reid, D.G., Rossberg, A.G., 2012. Recent data suggest no further  
673 recovery in North Sea Large Fish Indicator. *ICES J. Mar. Sci.* 69, 235–239.
- 674 Fung, T., Farnsworth, K.D., Shephard, S., Reid, D.G., Rossberg, A.G., 2013. Why the size  
675 structure of marine communities can require decades to recover from fishing. *Mar. Ecol.*  
676 *Prog. Ser.* 484, 155–171.
- 677 Furness, R.W., Camphuysen, K.C.J., 1997. Seabirds as monitors of the marine environment  
678 726–737.
- 679 Gaichas, S., Bundy, a, Miller, T., Moksness, E., Stergiou, K., 2012. What drives marine  
680 fisheries production? *Mar. Ecol. Prog. Ser.* 459, 159–163. doi:10.3354/meps09841
- 681 Garcia, S.M., Kolding, J., Rice, J., Rochet, M., Zhou, S., Arimoto, T., Beyer, J.E., Borges, L.,  
682 Bundy, A., Dunn, D., Fulton, E.A., Hall, M., Heino, M., Law, R., Makino, M., 2012.  
683 Reconsidering the Consequences of Selective Fisheries. *Science* (80-. ). 335, 1045–  
684 1047.
- 685 Garcia, S.M., Rice, J., Charles, A., 2015. Balanced harvesting in fisheries: a preliminary  
686 analysis of management implications. *ICES J. Mar. Sci.* 2–11.
- 687 Garcia, S.M., Staples, D.J., Chesson, J., 2000. The FAO guidelines for the development and  
688 use of indicators for sustainable development of marine capture fisheries and an  
689 Australian example of their application. *Ocean Coast. Manag.* 43, 537–556.
- 690 Garrison, L.P., Link, J.S., 2000. Dietary guild structure of the fish community in the  
691 Northeast United States continental shelf ecosystem. *Mar. Ecol. Prog. Ser.* 202, 231–  
692 240. doi:10.3354/meps202231
- 693 Gascuel, D., Bozec, Y., Chassot, E., Colomb, A., Laurans, M., 2005. The trophic spectrum:  
694 theory and application as an ecosystem indicator. *ICES J. Mar. Sci.* 62, 443–452.
- 695 Gascuel, D., Pauly, D., 2009. EcoTroph: modelling marine ecosystem functioning and impact  
696 of fishing. *Ecol. Modell.* 220, 2885–2898.
- 697 Gislason, H., Rice, J., 1998. Modelling the response of size and diveristy spectra of fish  
698 assemblages to changes in exploitation. *ICES J. Mar. Sci.* 55, 362–370.
- 699 Gowen, R., McQuatters-Gallop, A., Tett, P., Best, M., Bresnan, E., Castellani, C., Cook, K.,  
700 Sherer, C., McKinney, A., 2011. Plankton indicators. A report of a defra workshop held  
701 at AFBI 2nd-3d June 2011. Belfast.
- 702 Greenstreet, S., Rogers, S., 2006. Indicators of the health of the North Sea fish community:  
703 identifying reference levels for an ecosystem approach to management. *ICES J. Mar.*  
704 *Sci.* 63, 573–593. doi:10.1016/j.icesjms.2005.12.009
- 705 Greenstreet, S.P.R., Rogers, S.I., Rice, J.C., Piet, G.J., Guirey, E.J., Fraser, H.M., Fryer, R.J.,  
706 2011. Development of the EcoQO for the North Sea fish community. *ICES J. Mar. Sci.*  
707 68, 1–11.
- 708 Greenstreet, S.P.R., Rossberg, A.G., Fox, C.J., Quesne, W.J.F. Le, Blasdale, T., Boulcott, P.,  
709 Mitchell, I., Millar, C., Moffat, C.F., 2012. Demersal fish biodiversity: species-level  
710 indicators and trends-based targets for the Marine Strategy Framework Directive. *ICES*  
711 *J. Mar. Sci.* 69, 1789–1801.
- 712 Groffman, P.M., Baron, J.S., Blett, T., Gold, A.J., Goodman, I., Gunderson, L.H., Levinson,

713 B.M., Palmer, M. a., Paerl, H.W., Peterson, G.D., Poff, N.L., Rejeski, D.W., Reynolds,  
714 J.F., Turner, M.G., Weathers, K.C., Wiens, J., 2006. Ecological Thresholds: The Key to  
715 Successful Environmental Management or an Important Concept with No Practical  
716 Application? *Ecosystems* 9, 1–13. doi:10.1007/s10021-003-0142-z

717 Gunderson, L.H., 2000. Ecological Resilience - In Theory and Application. *Annu. Rev. Ecol.*  
718 *Syst.* 31, 425–439.

719 Hayes, K.R., Dambacher, J.M., Hosack, G.R., Bax, N.J., Dunstan, P.K., Fulton, E. a.,  
720 Thompson, P. a., Hartog, J.R., Hobday, a. J., Bradford, R., Foster, S.D., Hedge, P.,  
721 Smith, D.C., Marshall, C.J., 2015. Identifying indicators and essential variables for  
722 marine ecosystems. *Ecol. Indic.* 57, 409–419. doi:10.1016/j.ecolind.2015.05.006

723 Heath, M.R., Cook, R.M., Cameron, A.I., Morris, D.J., Speirs, D.C., 2014. Cascading  
724 ecological effects of eliminating fishery discards. *Nat. Commun.* 5, 1–8.  
725 doi:10.1038/ncomms4893

726 HELCOM, 2013. HELCOM core indicators. Final report of the HELCOM CORESET  
727 project. *Balt. Sea Environ. Proc.* 136, 1–74.

728 HELCOM, 2012. Development of a set of core indicators: interim report of the HELCOM  
729 CORESET project. Part A. Description of the selection process. *Balt. Sea Environ. Proc.*  
730 129A.

731 Heymans, J.J., Guenette, S., Christensen, V., 2007. Evaluating network analysis indicators of  
732 ecosystem status in the Gulf of Alaska. *Ecosystems* 10, 488–502.

733 Heymans, S., Coll, M., Libralato, S., Christensen, V., 2012. Ecopath theory, modelling and  
734 application to coastal ecosystems, in: McLusky, D., Woanski, E. (Eds.), *Trestise on*  
735 *Estuarine and Coastal Science*. Elsevier, pp. 93–111.

736 Hilting, A.K., Currin, C.A., Kosaki, R.K., 2013. Evidence for benthic primary production  
737 support of an apex predator – dominated coral reef food web. *Mar. Biol.* 160, 1681–  
738 1695. doi:10.1007/s00227-013-2220-x

739 Hinkley, D. V., 1970. Inference about the change-point in a sequence of a random variable.  
740 *Biometrika* 57, 1–17.

741 Hornborg, S., Belgrano, A., Bartolino, V., Valentinsson, D., Ziegler, F., 2013. Trophic  
742 indicators in fisheries: a call for re-evaluation. *Biol. Lett.* 9, 20121050.  
743 doi:10.1098/rsbl.2012.1050

744 Houle, J.E., Farnsworth, K.D., Rossberg, A.G., Reid, D.G., 2012. Assessing the sensitivity  
745 and specificity of fish community indicators to management action. *Can. J. Fish Aquat.*  
746 *Sci.* 69, 1065–1079.

747 Hughes, T.P., Bellwood, D.R., Folke, C., Steneck, R.S., Wilson, J., 2005. New paradigms for  
748 supporting the resilience of marine ecosystems. *Trends Ecol. Evol.* 20, 380–6.  
749 doi:10.1016/j.tree.2005.03.022

750 ICES, 2015. Report of the Working Group on Biodiversity Science (WGBIODIV) 9-13  
751 February 2015. Copenhagen, Denmark.

752 ICES, 2014. Report of the workshop to develop recommendations for potentially useful food  
753 web indicators (WKFooWI). Copenhagen, Denmark.

754 ICES, 2013a. Report on the working group on the ecosystem effects of fishing activities  
755 (WGECO), 1-8 May 2013, Copenhagen, Denmark.

756 ICES, 2013b. Report on the working group on multispecies assessment methods (WGSAM).

- 757ICES, 2011. Report on the working group on the ecosystem affects of fishing activities  
758 (WGECO). Copenhagen, Denmark.
- 759ICES, 2008. Report of the working group on ecosystem effects of fishing activities  
760 (WGECO) May 6-13 2008, Copenhagen, Denmark.
- 761Institute for European Environmental Policy (IEEP), 2005. A review of the indicators for  
762 ecosystem structure and functioning. INDECO Development of Indicators of  
763 Environmental Performance of Common Fisheries Policy Report.
- 764Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J.,  
765 Bradbury, R.H., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange,  
766 C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J.,  
767 Warner, R.R., 2001. Historical Overfishing and the Recent Collapse of Coastal  
768 Ecosystems. *Science* (80-. ). 293, 629–639.
- 769Jennings, S., Collingridge, K., 2015. Predicting Consumer Biomass, Size-Structure,  
770 Production, Catch Potential, Responses to Fishing and Associated Uncertainties in the  
771 World's Marine Ecosystems. *PLoS One* 10, e0133794.  
772 doi:10.1371/journal.pone.0133794
- 773Jeppesen, E., Noges, P., Davidson, T., Haberman, J., Noges, T., Blank, K., Lauridsen, R.,  
774 Sondergaard, M., Sayer, C., Laugaste, R., Johansson, L.S., Bjerring, R., Amsinck, S.L.,  
775 2011. Zooplankton as indicators in lakes: a scientific-based plea for including  
776 zooplankton in the ecological quality assessment of lakes according to the European  
777 Water Framework Directive (WFD). *Hydrobiologia* 676, 279–297.
- 778Kane, D., Gordon, S.I., Munawar, M., Charlton, M.N., Culver, D.A., 2009. The Planktonic  
779 Index of Biotic Integrity (P-IBI): an approach for assessing ecosystem health. *Ecol.*  
780 *Indic.* 9, 1234–1247.
- 781Kendall, C., Young, M.B., Silva, S.R., 2010. Applications of Stable Isotopes for Regional to  
782 National-Scale Water Quality and Environmental Monitoring Programs.  
783 doi:10.1007/978-90-481-3354-3
- 784Kerr, S.R., Dickie, I.M., 2001. *Biomass Spectrum*. Columbia University Press, New York,  
785 USA.
- 786Kershner, J., Samhouri, J.F., James, C.A., Levin, P.S., 2011. Selecting indicator portfolios for  
787 marine species and food webs: a Puget sound case study. *PLoS One* 6, e25248.  
788 doi:10.1371/journal.pone.0025248
- 789King, R.S., Baker, M.E., 2010. Considerations for analyzing ecological community  
790 thresholds in response to anthropogenic environmental gradients. *J. North Am. Benthol.*  
791 *Soc.* 29, 998–1008. doi:10.1899/09-144.1
- 792Large, S.I., Fay, G., Friedland, K.D., Link, J.S., 2015a. Critical points in ecosystem responses  
793 to fishing and environmental pressures. *Mar. Ecol. Prog. Ser.* 521, 1–17.  
794 doi:10.3354/meps11165
- 795Large, S.I., Fay, G., Friedland, K.D., Link, J.S., 2015b. Quantifying patterns of change in  
796 marine ecosystem response to multiple pressures. *PLoS One* 10, e0119922. doi:10.1371/  
797 journal.pone.0119922
- 798Large, S.I., Fay, G., Friedland, K.D., Link, J.S., 2013. Defining trends and thresholds in  
799 responses of ecological indicators to fishing and environmental pressures. *ICES J. Mar.*  
800 *Sci.* 70, 755–767.
- 801le Fevre, J., 1986. Aspects of the biology of frontal systems. *Adv. Mar. Biol.* 23, 163–299.

- 802Lekve, K., Ottersen, G., Stenseth, N.C., Gjosaeter, J., 2002. Length dynamics in juvenile  
803 coastal Skagerrak cod: effects of biotic and abiotic factors. *Ecology* 83, 1676–1688.
- 804Leslie, H.M., McLeod, K.L., 2007. Confronting the challenges of implementing marine  
805 ecosystem-based management. *Front. Ecol. Environ.* 5, 540–548. doi:10.1890/060093
- 806Lester, S.E., McLeod, K.L., Tallis, H., Ruckelshaus, M., Halpern, B.S., Levin, P.S., Chavez,  
807 F.P., Pomeroy, C., McCay, B.J., Costello, C., Gaines, S.D., Mace, A.J., Barth, J.A.,  
808 Fluharty, D.L., Parrish, J.K., 2010. Science in support of ecosystem-based management  
809 for the US West Coast and beyond. *Biol. Conserv.* 143, 576–587.  
810 doi:10.1016/j.biocon.2009.11.021
- 811Levin, P.S., Fogarty, M.J., Murawski, S. a, Fluharty, D., 2009. Integrated ecosystem  
812 assessments: developing the scientific basis for ecosystem-based management of the  
813 ocean. *PLoS Biol.* 7, e14. doi:10.1371/journal.pbio.1000014
- 814Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest 2, 18–22.
- 815Libralato, S., 2008. System Omnivory Index, in: Jorgensen, S.E., Fath, B.D. (Eds.),  
816 Ecological Indicators Vol. 4 of Encyclopedia of Ecology. Elsevier, Oxford, pp. 3472–  
817 3477.
- 818Libralato, S., Coll, M., Tempesta, M., Santojanni, A., Spoto, M., Palomera, I., Arneri, E.,  
819 Solidoro, C., 2010. Foodweb traits of protected and exploited areas of the Adriatic Sea.  
820 *Biol. Conserv.* 143, 2182–2194.
- 821Libralato, S., Coll, M., Tudela, S., Palomera, I., Pranovi, F., 2008. Novel index for  
822 quantification of ecosystem effects of fishing as removal of secondary production. *Mar.*  
823 *Ecol. Prog. Ser.* 355, 107–129.
- 824Libralato, S., Pranovi, F., Raicevich, S., Da Pointe, F., Giovanardi, O., Pastres, R., Torricelli,  
825 P., Mainardi, D., 2004. Ecological stages of the Venice Lagoon analysed using landing  
826 time series data. *J. Mar. Syst.* 51, 331–334.
- 827Lindeman, R.L., 1942. The trophic-dynamic aspect of ecology. *Ecology* 23, 339–417.
- 828Link, J.S., 2010. Ecosystem-Based Fisheries Management: Confronting Tradeoffs, 1st ed.  
829 Cambridge University Press, New York, USA.
- 830Link, J.S., 2005. Translating ecosystem indicators into decision criteria. *ICES J. Mar. Sci.* 62,  
831 569–576. doi:10.1016/j.icesjms.2004.12.015
- 832Link, J.S., 2002a. What Does Ecosystem-Based Fisheries Management Mean? *Fisheries* 27,  
833 18–21.
- 834Link, J.S., 2002b. Does food web theory work for marine ecosystems? *Mar. Ecol. Prog. Ser.*  
835 230, 1–9.
- 836Link, J.S., Almeida, F.P., 2002. Opportunistic feeding of longhorn sculpin (*Myoxocephalus*  
837 *octodecemspinosus*): are scallo fishery discards an important food subsidy for  
838 scavengers on Georges Bank. *NOAA Fish. Bull.* 100, 381–385.
- 839Link, J.S., Bundy, A., Overholtz, W.J., Shackell, N., Manderson, J., Duplisea, D., Hare, J.,  
840 Koen-Alonso, M., Friedland, K.D., 2011. Ecosystem-based fisheries management in the  
841 Northwest Atlantic. *Fish Fish.* 12, 152–170. doi:10.1111/j.1467-2979.2011.00411.x
- 842Link, J.S., Pranovi, F., Coll, M., Libralato, S., Christensen, V., Legault, C., 2009. Exploring  
843 novel metrics of ecosystem overfishing using energy budget model outputs. *Fish. Cent.*  
844 *Res. Rep.* 17, 153.
- 845Link, J.S., Pranovi, F., Libralato, S., Coll, M., Christensen, V., Solidoro, C., Fulton, E. a,

- 846 2015. Emergent Properties Delineate Marine Ecosystem Perturbation and Recovery.  
847 *Trends Ecol. Evol.* xx, 1–13. doi:10.1016/j.tree.2015.08.011
- 848 Link, J.S., Stockhausen, W.T., Methratta, E.T., 2005. Food web theory in marine ecosystems,  
849 in: Belgrano, A., Scharler, U.M., Dunne, J., Ulanowicz, R.E. (Eds.), *Aquatic Food*  
850 *Webs: An Ecosystem Approach*. Oxford University Press, Oxford, UK, pp. 98–113.
- 851 Litzow, M.A., Bailey, K.M., Prahl, F.G., Heintz, R., 2006. Climate regime shifts and  
852 reorganization of fish communities: the essential fatty acid limitation hypothesis. *Mar.*  
853 *Ecol. Prog. Ser.* 345, 1–11.
- 854 Loh, J., Green, R.E., Ricketts, T., Lamoreux, J., Jenkins, M., Kapos, V., Randers, J., 2005.  
855 The Living Planet Index: using species population time-series to track trends in  
856 biodiversity. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 360, 289–295.
- 857 Longo, C., Hornborg, S., Bartolino, V., Tomczak, M., Ciannelli, L., Libralato, S., Belgrano,  
858 A., 2015. Role of trophic models and indicators in current marine fisheries management.  
859 *Mar. Ecol. Prog. Ser.* 538, 257–272. doi:10.3354/meps11502
- 860 Lougheed, V.L., Chow-Fraser, P., 2002. Development and use of a zooplankton index of  
861 wetland quality in the Laurentian Great Lakes Basin. *Ecol. Appl.* 12, 474–486.
- 862 Mallory, M.L., Robinson, S. a, Hebert, C.E., Forbes, M.R., 2010. Seabirds as indicators of  
863 aquatic ecosystem conditions: a case for gathering multiple proxies of seabird health.  
864 *Mar. Pollut. Bull.* 60, 7–12. doi:10.1016/j.marpolbul.2009.08.024
- 865 Marasco, R.J., Goodman, D., Grimes, C.B., Lawson, P.W., Punt, A.E., Quinn II, T.J., 2007.  
866 Ecosystem-based fisheries management: some practical suggestions. *Can. J. Fish. Aquat.*  
867 *Sci.* 64, 928–939. doi:10.1139/f07-062
- 868 Martin, J., Runge, M.C., Nichols, J.D., Lubow, B.C., Kendall, W.L., 2009. Structured  
869 decision making as a conceptual framework to identify thresholds for conservation and  
870 management. *Ecol. Appl.* 19, 1079–1090.
- 871 McLeod, K.L., Lubchenco, J., Palumbi, S.R., Rosenberg, A.A., 2005. Scientific consensus  
872 statement on Marine Ecosystem-Based Management. *Commun. Parternesh. Sci. Sea.*
- 873 Mendoza, G.A., Martins, H., 2006. Multi-criteria decision analysis in natural resource  
874 management: a critical review of methods and new modelling paradigms. *For. Ecol.*  
875 *Manage.* 230, 1–22.
- 876 Methratta, E., Link, J.S., 2006a. Evaluation of quantitative indicators for marine fish  
877 communities. *Ecol. Indic.* 6, 575–588. doi:10.1016/j.ecolind.2005.08.022
- 878 Methratta, E., Link, J.S., 2006b. Evaluation of quantitative indicators for marine fish  
879 communities. *Ecol. Indic.* 6, 575–588. doi:10.1016/j.ecolind.2005.08.022
- 880 Meuter, F.J., Megrey, B.A., 2006. Using multispecies surplus production models to estimate  
881 ecosystem-level maximum sustainable yeilds. *Fish. Res.* 81, 189–201.
- 882 Moloney, C., Jarre, a, Arancibia, H., Bozec, Y., Neira, S., Roux, J., Shannon, L., 2005.  
883 Comparing the Benguela and Humboldt marine upwelling ecosystems with indicators  
884 derived from inter-calibrated models. *ICES J. Mar. Sci.* 62, 493–502.  
885 doi:10.1016/j.icesjms.2004.11.009
- 886 Monaco, M.E., Ulanowicz, R.E., 1997. Comparative ecosystem trophic structure of three  
887 U.S. mid Atlantic estuaries. *Mar. Ecol. Prog. Ser.* 161, 239–254.
- 888 Neira, S., Moloney, C.L., Cury, P., Mullon, C., Christensen, V., 2009. Mechanisms affecting  
889 recovery in an upwelling food web: The case of the southern Humboldt. *Prog. Oceanogr.*

- 890 83, 404–416.
- 891 Nicholson, M., Jennings, S., 2004. Testing candidate indicators to support ecosystem-based  
892 management: the power of monitoring surveys to detect temporal trends in fish  
893 community metrics. *ICES J. Mar. Sci.* 61, 35–42. doi:10.1016/j.icesjms.2003.09.004
- 894 Olson, B.D.B., Hitchcock, G.L., Mariano, A.J., Ashjian, C.J., Peng, G., Nero, R.W., Podest,  
895 G.P., 1994. Life on the edge: Marine life and fronts. *Oceanography* 7, 52–60.
- 896 Ostrom, E., Burger, J., Field, C.B., Norgaard, R.B., Policansky, D., 1999. Revisiting the  
897 Commons: Local Lessons, Global Challenges. *Science* (80-. ). 284, 278–283.
- 898 Parsons, M., Mitchell, I., Butler, A., Ratcliffe, N., Frederiksen, M., Foster, S., Reid, J.B.,  
899 2008. Seabirds as indicators of the marine environment. *ICES J. Mar. Sci.* 65, 1520–  
900 1526.
- 901 Pauly, D., Christensen, V., 1995. Primary production required to sustain global fisheries.  
902 *Nature* 374, 255–257.
- 903 Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., Jr, F.T., Series, N., Feb, N., 1998.  
904 Fishing down marine food webs. *Science* (80-. ). 279, 860–863.
- 905 Pauly, D., Christensen, V., Walters, C., 2000. Ecopath, Ecosim, and Ecospace as tools for  
906 evaluating ecosystem impact of fisheries. *ICES J. Mar. Sci.* 57, 697–706.  
907 doi:10.1006/jmsc.2000.0726
- 908 Pauly, D., Palomares, M., 2005. Fishing down marine food web: It is far more pervasive than  
909 we thought. *Bull. Mar. Sci.* 76, 197–211.
- 910 Pauly, D., Watson, R., 2005. Background and interpretation of the “Marine Trophic Index” as  
911 a measure of biodiversity. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 360, 415–23.  
912 doi:10.1098/rstb.2004.1597
- 913 Perez-Espana, H., Arreguin-Sanchez, F., 1999. Complexity related to behavior of stability in  
914 modelled coastal-zone ecosystems. *Aquat. Ecosyst. Heal. Manag.* 2, 129–135.
- 915 Pershing, A.J., Greene, C.H., Jossi, J.W., Brien, L.O., Brodziak, J.K.T., Bailey, B.A., 2005.  
916 Interdecadal variability in the Gulf of Maine zooplankton community, with potential  
917 impacts on fish recruitment. *ICES J. Mar. Sci.* 62, 1511–1523.
- 918 Piet, G., Jennings, S., 2005. Response of potential fish community indicators to fishing. *ICES*  
919 *J. Mar. Sci.* 62, 214–225. doi:10.1016/j.icesjms.2004.09.007
- 920 Piroddi, C., Teixeira, H., Lynam, C.P., Smith, C., Alvarez, M.C., Mazik, K., Andonegi, E.,  
921 Churilova, T., Tedesco, L., Chifflet, M., Chust, G., Galparsoro, I., Garcia, A.C., Kämäri,  
922 M., Kryvenko, O., Lassalle, G., Neville, S., Niquil, N., Papadopoulou, N., Rossberg,  
923 A.G., Suslin, V., Uyarra, M.C., 2015. Using ecological models to assess ecosystem  
924 status in support of the European Marine Strategy Framework Directive. *Ecol. Indic.* 58,  
925 175–191. doi:10.1016/j.ecolind.2015.05.037
- 926 Pitcher, C.R., Lawton, P., Ellis, N., Smith, S.J., Incze, L.S., Wei, C.-L., Greenlaw, M.E.,  
927 Wolff, N.H., Sameoto, J.A., Snelgrove, P.V.R., Cadotte, M., 2012. Exploring the role of  
928 environmental variables in shaping patterns of seabed biodiversity composition in  
929 regional-scale ecosystems. *J. Appl. Ecol.* 49, 670–679. doi:10.1111/j.1365-  
930 2664.2012.02148.x
- 931 Pitcher, T.J., Kalikoski, D., Short, K., Varkey, D., Pramod, G., 2009. An evaluation of  
932 progress in implementing ecosystem-based management of fisheries in 33 countries.  
933 *Mar. Policy* 33, 223–232. doi:10.1016/j.marpol.2008.06.002

- 934Polis, G.A., Strong, D.R., 1996. Food web complexity and community dynamics. *Am. Nat.*  
935 147, 813–846.
- 936Polivina, J.J., Howell, E., Kobayashi, D.R., Seki, M.P., 2001. The transition zone chlorophyll  
937 front, a dynamic global feature defining migration and forage habitat for marine  
938 resources. *Prog. Oceanogr.* 49, 469–483.
- 939Pranovi, F., Libralato, S., Zucchetto, M., Link, J.S., 2014. Biomass accumulation across  
940 trophic levels: analysis of landings for the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 512,  
941 201–216. doi:10.3354/meps10881
- 942Pranovi, F., Link, J.S., Fu, C., Cook, A.M., Liu, H., Gaichas, S., Friedland, K.D., Rong Utne,  
943 K., Benoît, H.P., 2012. Trophic-level determinants of biomass accumulation in marine  
944 ecosystems. *Mar. Ecol. Prog. Ser.* 459, 185–201. doi:10.3354/meps09738
- 945Prasad, A.M., Iverson, L.R., Liaw, A., 2006. Newer Classification and Regression Tree  
946 Techniques: Bagging and Random Forests for Ecological Prediction. *Ecosystems* 9,  
947 181–199. doi:10.1007/s10021-005-0054-1
- 948Pretty, J., 2003. Social Capital and the Collective Management. *Science* (80-. ). 302, 1912–  
949 1914.
- 950Rice, J., 2003. Environmental health indicators. *Ocean Coast. Manag.* 46, 235–259.  
951 doi:10.1016/S0964-5691(03)00006-1
- 952Rice, J., 2000. Evaluating fishery impacts using metrics of community structure. *ICES J.*  
953 *Mar. Sci.* 57, 682–688. doi:10.1006/jmsc.2000.0735
- 954Rice, J., Gislason, H., 1996. Patterns of change in the size spectra of numbers and diversity of  
955 the North Sea fish assemblage, as reflected in surveys and models. *ICES J. Mar. Sci.*  
956 53, 1214–1225.
- 957Rice, J., Rochet, M., 2005a. A framework for selecting a suite of indicators for fisheries  
958 management. *ICES J. Mar. Sci.* 62, 516–527. doi:10.1016/j.icesjms.2005.01.003
- 959Rice, J., Rochet, M., 2005b. A framework for selecting a suite of indicators for fisheries  
960 management. *ICES J. Mar. Sci.* 62, 516–527. doi:10.1016/j.icesjms.2005.01.003
- 961Rochet, M., Rice, J., 2005. Do explicit criteria help in selecting indicators for ecosystem-  
962 based fisheries management? *ICES J. Mar. Sci.* 62, 528–539.  
963 doi:10.1016/j.icesjms.2005.01.007
- 964Rochet, M.-J., Collie, J.S., Trenkel, V.M., 2013. How do fishing and environmental effects  
965 propagate among and within functional groups? *Bull. Mar. Sci.* 89, 285–315.
- 966Rochet, M.-J., Trenkel, V.M., 2003. Which community indicators can measure the impact of  
967 fishing? A review and proposals. *Can. J. Fish. Aquat. Sci.* 60, 86–99. doi:10.1139/f02-  
968 164
- 969Röckmann, C., van Leeuwen, J., Goldsborough, D., Kraan, M., Piet, G., 2015. The  
970 interaction triangle as a tool for understanding stakeholder interactions in marine  
971 ecosystem based management. *Mar. Policy* 52, 155–162.  
972 doi:10.1016/j.marpol.2014.10.019
- 973Rockstrom, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S.I., Lambin, E.F., Lenton,  
974 T.M., Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T.,  
975 van der Leeuw, S., Rodhe, H., Sörlin, S., Snyder, P.K., Costanza, R., Svedin, U.,  
976 Falkenmark, M., Karlberg, L., Corell, R.W., Falbry, V.J., Hansen, J., Walker, B.,  
977 Liverman, D., Richardson, K., Crutzen, P., Foley, J.A., 2009. A safe operating space for  
978 humanity. *Nature* 461, 472–475.



- 979Rodinov, S.N., 2004. A sequential algorithm for testing climate regime shifts. *Geophys. Res.*  
980 *Lett.* 31, L09204.
- 981Rogers, S.I., Casini, M., Cur, P., Heat, M., Irigoe, X., Kuos, H., Scheida, M., Sko, H.,  
982 Stergio, K., Trenkel, V.M., Wikner, J., Yunev, O., 2010. Marine Strategy Framework  
983 Directive Task Group 4 Report Food Webs.
- 984Rosenfeld, J.S., 2002. Functional Redundancy in Ecology and Conservation. *Oikos* 98, 156–  
985 162.
- 986Rossberg, A.G., 2013. Food webs and biodiversity: foundations, models, data. Wiley,  
987 Oxford, UK.
- 988Rossberg, A.G., 2012. Food Webs, in: Hastings, A., Gross, L. (Eds.), *Encyclopedia of*  
989 *Theoretical Ecology*. University of California Press, Berkeley, CA. USA., pp. 1–13.
- 990Rossberg, A.G., Farnsworth, K.D., Satoh, K., Pinnegar, J.K., 2011. Universal power-law diet  
991 partitioning by marine fish and squid with surprising stability-diversity implications.  
992 *Proc. Biol. Sci.* 278, 1617–25. doi:10.1098/rspb.2010.1483
- 993Rossberg, A.G., Yanagi, K., Amemiya, T., Itoh, K., 2006. Estimating trophic link density  
994 from quantitative but incomplete diet data. *J. Theor. Biol.* 243, 261–72.  
995 doi:10.1016/j.jtbi.2006.06.019
- 996Samhouri, J.F., Lester, S.E., Selig, E.R., Halpern, B.S., Fogarty, M.J., Longo, C., McLeod,  
997 K.L., 2012. Sea sick? Setting targets to assess ocean health and ecosystem services.  
998 *Ecosphere* 3, 41.
- 999Samhouri, J.F., Levin, P.S., Ainsworth, C.H., 2010. Identifying thresholds for ecosystem-  
1000 based management. *PLoS One* 5, e8907. doi:10.1371/journal.pone.0008907
- 1001Sandström, A., Bodin, Ö., Crona, B., 2015. Network Governance from the top – The case of  
1002 ecosystem-based coastal and marine management. *Mar. Policy* 55, 57–63.  
1003 doi:10.1016/j.marpol.2015.01.009
- 1004Sasaki, T., Furukawa, T., Iwasaki, Y., Seto, M., Mori, A.S., 2015. Perspectives for ecosystem  
1005 management based on ecosystem resilience and ecological thresholds against multiple  
1006 and stochastic disturbances. *Ecol. Indic.* 57, 395–408. doi:10.1016/j.ecolind.2015.05.019
- 1007Scott, B.E., Sharples, J., Wanless, S., Ross, O., Frederiksen, M., Daunt, F., 2006. The use of  
1008 biologically meaningful oceanographic indices to separate the effects of climate and  
1009 fisheries on seabird breeding success, in: Boyd, I.I., Wanless, S., Camphuysen, C.J.  
1010 (Eds.), *Management of Marine Ecosystems*. Cambridge University Press, Cambridge,  
1011 UK., pp. 46–62.
- 1012Shackell, N.L., Bundy, A., Nye, J.A., Link, J.S., 2012. Common large-scale responses to  
1013 climate and fishing across Northwest Atlantic ecosystems. *ICES J. Mar. Sci.* 69, 151–  
1014 162.
- 1015Shannon, L.J., Coll, M., Bundy, A., Gascuel, D., Heymans, J.J., Kleisner, K., Lynam, C.,  
1016 Piroddi, C., Tam, J., Travers-Trolet, M., Shin, Y.-J., 2014. Trophic level-based  
1017 indicators to track fishing impacts across marine ecosystems. *Mar. Ecol. Prog. Ser.* 512,  
1018 115–140. doi:10.3354/meps10821
- 1019Shannon, L.J., Coll, M., Neira, S., 2009. Exploring the dynamics of ecological indicators  
1020 using food web models fitted to time series of abundance and catch data. *Ecol. Indic.* 9,  
1021 1078–1095. doi:10.1016/j.ecolind.2008.12.007
- 1022Shephard, S., Fung, T., Houle, J.E., Farnsworth, K.D., Reid, D.G., Rossberg, A.G., 2012.  
1023 Size-selective fishing drives species composition in the Celtic Sea. *ICES J. Mar. Sci.* 69,

- 1024 223–234. doi:10.1093/icesjms/fsr200
- 1025 Shephard, S., Fung, T., Rossberg, A.G., Farnsworth, K.D., Reid, D.G., Greenstreet, S.P.R.,  
 1026 Warnes, S., 2013. Modelling recovery of Celtic Sea demersal fish community size-  
 1027 structure. *Fish. Res.* 140, 91–95. doi:10.1016/j.fishres.2012.12.010
- 1028 Shephard, S., Reid, D.G., Greenstreet, S.P.R., 2011. Interpreting the large fish indicator for  
 1029 the Celtic Sea. *ICES J. Mar. Sci.* 68, 1963–1972. doi:10.1093/icesjms/fsr114
- 1030 Shephard, S., Rindorf, A., Dickey-collas, M., Hintzen, N.T., Farnsworth, K., Reid, D.G.,  
 1031 2014. Assessing the state of pelagic fish communities within an ecosystem approach and  
 1032 European Marine Strategy Framework Directive. *ICES J. Mar. Sci.* 71, 1572–1585.
- 1033 Shin, Y.-J., Bundy, A., Shannon, L.J., Simier, M., Coll, M., Fulton, E.A., Link, J.S., Jouffre,  
 1034 D., Ojaveer, H., Mackinson, S., Heymans, J.J., Raid, T., 2010a. Can simple be useful  
 1035 and reliable? Using ecological indicators to represent and compare the states of marine  
 1036 ecosystems. *ICES J. ...* 67, 717–731.
- 1037 Shin, Y.-J., Rochet, M.-J., Jennings, S., Field, J., Gislason, H., 2005. Using size-based  
 1038 indicators to evaluate the ecosystem effects of fishing. *ICES J. Mar. Sci.* 62, 384–396.  
 1039 doi:10.1016/j.icesjms.2005.01.004
- 1040 Shin, Y.-J., Shannon, L.J., 2009. Using indicators for evaluating, comparing, and  
 1041 communicating the ecological status of exploited marine ecosystems. 1. The IndiSeas  
 1042 project. *ICES J. Mar. Sci.* 67, 686–691. doi:10.1093/icesjms/fsp273
- 1043 Shin, Y.-J., Shannon, L.J., Bundy, A., Coll, M., Aydin, K., Bez, N., Blanchard, J.L., Borges,  
 1044 M.D.F., Diallo, I., Diaz, E., Heymans, J.J., Hill, L., Johannesen, E., Jouffre, D., Kifani,  
 1045 S., Labrosse, P., Link, J.S., Mackinson, S., Masski, H., Mo, C., Neira, S., Ojaveer, H.,  
 1046 Abdallahi, M., Perry, I., Thiao, D., Yemane, D., Cury, P.M., 2010b. Using indicators for  
 1047 evaluating, comparing, and communicating the ecological status of exploited marine  
 1048 ecosystems. 2. Setting the scene. *ICES J. Mar. Sci.* 67, 692–716.
- 1049 Slocombe, D.S., 1998. Lessons from experience with ecosystem-based management. *Landsc.*  
 1050 *Urban Plan.* 40, 31–39. doi:10.1016/S0169-2046(97)00096-0
- 1051 Slocombe, D.S., 1993. Implementing Ecosystem-based Management. *Bioscience* 43, 612–  
 1052 622.
- 1053 Smith, A.D.M., Fulton, E.A., Hobday, A.J., Smith, D.C., Shoulder, P., 2007. Scientific tools  
 1054 to support the practical implementation of ecosystem-based fisheries management. *ICES*  
 1055 *J. Mar. Sci.* 64, 633–639.
- 1056 Sonderegger, D.L., Wang, H., Clements, W.H., Noon, B.R., 2008. Using SiZer to detect  
 1057 thresholds in ecological data. *Front. Ecol. Environ.* 7, 190–195.
- 1058 Soto, K.H., Trites, A.W., Arias-Schreiber, M., 2006. Changes in diet and maternal attendance  
 1059 of South American Sea lions indicate changes in the marine environment and prey  
 1060 abundance. *Mar. Ecol. Prog. Ser.* 312, 277–290.
- 1061 Stige, L.C., Dalpadado, P., Orlova, E., Boulay, A.-C., Durant, J.M., Ottersen, G., Stenseth,  
 1062 N.C., 2014. Spatiotemporal statistical analyses reveal predator-driven zooplankton  
 1063 fluctuations in the Barents Sea. *Prog. Oceanogr.* 120, 243–253.  
 1064 doi:10.1016/j.pocean.2013.09.006
- 1065 Tallis, H., Levin, P.S., Ruckelshaus, M., Lester, S.E., McLeod, K.L., Fluharty, D.L., Halpern,  
 1066 B.S., 2010. The many faces of ecosystem-based management: Making the process work  
 1067 today in real places. *Mar. Policy* 34, 340–348. doi:10.1016/j.marpol.2009.08.003
- 1068 Teichert, N., Borja, A., Chust, G., Uriarte, A., Lepage, M., 2015. Restoring fish ecological

- 1069 quality in estuaries: Implication of interactive and cumulative effects among  
 1070 anthropogenic stressors. *Sci. Total Environ.* 542, 383–393.  
 1071 doi:10.1016/j.scitotenv.2015.10.068
- 1072Tett, P., Carrera, C., Mills, D.K., van Leeuwen, S., Foden, J., Bresnan, E., Gowen, R., 2008.  
 1073 Use of a Phytoplankton Community Index to assess the health of coastal waters. *ICES J.*  
 1074 *Mar. Sci.* 65, 1475–1482.
- 1075Tett, P., Gowen, R.J., Painting, S.J., Elliott, M., Forster, R., Mills, D.K., Bresnan, E.,  
 1076 Capuzzo, E., Fernandes, T.F., Foden, J., Geider, R.J., Gilpin, L.C., Huxham, M.,  
 1077 McQuatters-Gollop, A.L., Malcolm, S.J., Saux-Picart, S., Platt, T., Racault, M.F.,  
 1078 Sathyendranath, S., van der Molen, J., Wilkinson, M., 2013. Framework for  
 1079 understanding marine ecosystem health. *Mar. Ecol. Prog. Ser.* 494, 1–27.  
 1080 doi:10.3354/meps10539
- 1081Thompson, R.M., Brose, U., Dunne, J.A., Hall, R.O., Hladysz, S., Kitching, R.L., Martinez,  
 1082 N.D., Rantala, H., Romanuk, T.N., Stouffer, D.B., Tylianakis, J.M., 2012. Food webs:  
 1083 reconciling the structure and function of biodiversity. *Trends Ecol. Evol.* 27, 689–97.  
 1084 doi:10.1016/j.tree.2012.08.005
- 1085Thrush, S.F., Dayton, P.K., 2010. What can ecology contribute to ecosystem-based  
 1086 management? *Ann. Rev. Mar. Sci.* 2, 419–41. doi:10.1146/annurev-marine-120308-  
 1087 081129
- 1088Toms, J.D., Lesperance, M.L., 2003. A tool for identifying ecological thresholds. *Ecology* 84,  
 1089 2034–2041.
- 1090Toms, J.D., Villard, M., 2015. Threshold detection: matching statistical methodology to  
 1091 ecological. *Avian Conserv. Ecol.* 10, 2.
- 1092Trenkel, V.M., Rochet, M.-J., 2003. Performance of indicators derived from abundance  
 1093 estimates for detecting the impact of fishing on a fish community. *Can. J. Fish. Aquat.*  
 1094 *Sci.* 60, 67–85. doi:10.1139/f02-163
- 1095Ulanowicz, R.E., 2004. Quantitative methods for ecological network analysis. *Comput. Biol.*  
 1096 *Chem.* 28, 321–228.
- 1097Ulanowicz, R.E., 1992. Ecosystem health and trophic flow networks, in: Costanza, R.,  
 1098 Northon, B.G., Haskell, B.D. (Eds.), *Ecosystem Health: New Goals for Environmental*  
 1099 *Management*. Washington, DC, pp. 190–225.
- 1100van Leeuwen, S., Tett, P., Mills, D., van der Molen, J., 2015. Stratified and nonstratified  
 1101 areas in the North Sea: long-term variability and biological and policy implications. *J.*  
 1102 *Geophys. Res.* 120, 4670–4686.
- 1103Vargas, C. a, Escribano, R., Poulet, S., 2006. Phytoplankton food quality determines time  
 1104 windows for successful zooplankton reproductive pulses. *Ecology* 87, 2992–2999.
- 1105Vasconcellos, M., Mackinson, S., Sloman, S., Pauly, D., 1997. The stability of trophic mass-  
 1106 balance models of marine ecosystems: a comparative analysis. *Ecol. Modell.* 100, 125–  
 1107 134.
- 1108Wanless, S., Harris, M.P., Redman, P., Speakman, J.R., 2005. Low energy values of fish as a  
 1109 probable cause of a major seabird breeding failure in the North Sea. *Mar. Ecol. Prog.*  
 1110 *Ser.* 294, 1–8.
- 1111Wilderbuer, T.K., Hollowed, A.B., Ingraham, W.J., Spencer, P.D., Connors, M.E., Bond,  
 1112 N.A., Walters, G.E., 2002. Flatfish recruitment response to decadal climactic variability  
 1113 and ocean condiditons in the eastern Bering Sea. *Prog. Oceanogr.* 55, 235–247.

- 1114 Williams, R.J., Martinez, N.D., 2004. Limits to trophic levels and omnivory in complex food  
1115 webs: Theory and data. *Am. Nat.* 163.
- 1116 Xu, W., Mage, J.A., 2001. A review of concepts and criteria for assessing agroecosystem  
1117 health including a preliminary case study of southern Ontario. *Agric. Ecosyst.*  
1118 *Environ.* 83, 215–233.
- 1119 Zador, S., 2013. Ecosystem considerations 2013. Appendix C in Stock Assessment and  
1120 Fishery Evaluation Report of the Groundfish Resources of the Bering Sea/Aleutian  
1121 Islands Regions. Anchorage, AK. USA.
- 1122 Zador, S., Aydin, K., Barbeaux, S., Batten, S., Bengston, J., Bond, N., Ciciel, K., Fergusson,  
1123 E., Fitzgerald, S., Fritz, L., Gann, J., Garcia-reyes, M., Greig, A., Hatch, S., Hebert, K.,  
1124 Hermann, A., Ladd, C., Lee, J., Litzow, M., Ortiz, I., Overland, J., Piatt, J., Ream, R.,  
1125 Renner, H., Ressler, P., Salo, S., Stabeno, P., Sterling, J., Stockhausen, W., Sydeman,  
1126 B., Thompson, A., Towell, R., Urban, D., Wand, M., Wertheimer, A., Whitehouse, A.,  
1127 Williams, J., Yasumiishi, E., 2014. Eastern Bering Sea 2014 Report Card.
- 1128 Zeileis, A., Kleiber, C., 2005. Validating multiple structural change models: a case study. *J.*  
1129 *Appl. Econom.* 20, 685–690.
- 1130

1131

1132 Table 1. Criteria and sub-criteria used in the selection process for operational food-web  
1133 indicators.

Criteria	Sub-criteria (issues)	Rationale
Availability of underlying data	Existing and ongoing data	Indicators are supported by current or planned monitoring programmes that provide the data necessary to derive the indicator. Ideal monitoring programmes should have a time series capable of supporting baselines and reference point setting. Data should be collected on multiple sequential occasions using consistent protocols.
	Relevant spatial coverage	Data should be derived from an appropriate proportion of the regional sea, at appropriate spatial resolution and sampling design, to which the indicator will apply.
	Relevant temporal coverage	Data should be collected at appropriate sampling frequency and for an appropriate extent of time relevant to the time scale of the process or attribute the indicator describes.
Quality of underlying data	Indicators should be technically rigorous	Indicators should ideally be easily and accurately determined using technically feasible and quality assured methods.
	Reflects changes in ecosystem component that are caused by variation in any specified manageable pressures	The indicator reflects change in the state of an ecological component that is caused by specific significant manageable pressures (e.g. fishing mortality, habitat destruction). The indicator should therefore respond sensitively to particular changes in pressure. The response should be based on theoretical or empirical knowledge, thus reflecting the effect of change in pressure on the ecosystem component in question; signal to noise ratio should be high. Ideally the pressure-state relationship should be defined under both the disturbance and recovery phases.
	Magnitude, direction and variance of indicator is estimable	The indicator should exhibit a predictable direction, exhibit clear sense of magnitude of any change, and estimates of precision should allow for detection of trends or distinct locales - requiring that some measure of sampling error or variance estimator is available.
Conceptual basis	Scientific credibility	Scientific, peer-reviewed findings should underpin the assertion that the indicator provides a true representation of process, and variation thereof, for the ecosystem attribute being examined.
	Associated with key processes	The link between the indicator and a process that is essential to food web functioning should be clear and established, based on our current understanding of trophic dynamics.
	Unambiguous	The indicator responds unambiguously to a pressure.
Communication	Comprehensible	Indicators should be interpretable in a way that is easily understandable by policy-makers and other non-scientists (e.g. stakeholders) alike, and the consequences of variation in the indicator should be easy to communicate.
Management	Relevant to management	Indicator links directly to mandated management needs, and ideally to management response. The relationship between human activity and resulting pressure on the ecological component is clearly understood.
	Management thresholds targets are estimable	Clear targets that meet appropriate target criteria (absolute values or trend directions) for the indicator can be specified that reflect management objectives, such as achieving GES. Ideally control rules can be developed.
	Cost-effectiveness	Sampling, measuring, processing, analysing indicator data, and reporting assessment outcomes should make effective use of limited financial resources.

1134 Table 2a. Assessment of food-web indicators for Energy flow indicators against the criteria in Table 1. A maximum score for  
 1135 Availability of data = 6, Quality of data = 6, Conceptual = 6, Communication = 2, Management = 6 (maximum total score = 26).  
 1136 Asterisks (\*) denote food-web indicators that were selected for current use.

	Food-web indicator	Availability	Quality	Conceptual	Communication	Management	Score	Percentage	Other indicator uses
<b>Energy Flow indicators</b>	*Seabird breeding success	6	3	6	2	5	22	85	Biological diversity
	Mean weight at age of predatory fish species from data	4	5	5	2	5	21	81	Fisheries
	Total mortality Z	4	5	4	1	5	19	73	Fisheries
	Productivity of key predators	6	3	4	1	4	18	69	
	*Primary production required to support fisheries	4	3	6	0	5	18	69	Fisheries, Biological diversity
	Productive pelagic habitat index	6	4	4	1	3	18	69	Eutrophication, Fisheries, Biological diversity
	Ecosystem exploitation	5	3	2	1	5	16	62	Fisheries
	Community condition	3	5	3	2	3	16	62	Fisheries
	*Mean trophic level of catch	4	4	2	1	4	15	58	Fisheries
	*Marine trophic index of the community	4	3	4	1	3	15	58	
	*Mean trophic level of the community	4	3	4	1	3	15	58	
	Disturbance index	4	3	4	1	2	14	54	
	Loss in secondary production index	4	3	4	0	3	14	54	Fisheries
	Cumulative distribution of biomass assessment	4	3	4	0	3	14	54	Fisheries
	Trophic balance index	4	2	3	0	4	13	50	
	Mean transfer efficiency for a given trophic level or size	3	2	4	0	1	10	38	
	Finn cycling index	3	1	4	0	1	9	35	Fisheries

1137

1138

1139

1140

1141 Table 2b. Assessment of food-web indicators for Ecosystem resilience and Structural indicators against the criteria in Table 1. A  
 1142 maximum score for Availability of data = 6, Quality of data = 6, Conceptual = 6, Communication = 2, Management = 6 (maximum  
 1143 total score = 26). Asterisks (\*) denote food-web indicators that were selected for current use.

	Food-web indicator	Availability	Quality	Conceptual	Communication	Management	Score	Percent	Other indicator uses
Ecosystem resilience indicators	*Mean trophic links per species	3	2	4	1	2	12	46	Biological diversity
	Ecological network analysis derived indicators	4	1	4	1	2	12	46	
	Gini-Simpson dietary diversity index	3	2	4	1	1	11	42	
	Herbivory to detritivory ratio	3	1	4	1	1	10	38	
	Ecological network indices of ecosystem status and change	4	1	4	0	1	10	38	
	System omnivory index	3	1	2	0	1	7	27	
Structural indicators	Guild surplus production models	6	6	6	1	6	25	96	Fisheries
	Large fish indicator	6	6	5	2	6	25	96	Fisheries
	*Total biomass of small fish	6	5	5	2	5	23	88	Fisheries
	Proportion of predatory fish	6	3	5	2	6	22	85	Biological diversity, Fisheries
	*Mean length of surveyed community	6	6	4	2	4	22	85	Biological diversity, Fisheries
	Pelagic to demersal ratio	6	5	4	2	4	21	81	Fisheries, Eutrophication
	*Biomass of trophic guilds	4	3	5	2	6	20	77	Biological diversity, Fisheries
	Lifeform-based indicator for the pelagic habitat	6	5	4	1	4	20	77	Biological diversity, Eutrophication, Sea-floor integrity
	Region-specific indicators of abundance and spatial distribution	6	3	4	1	5	19	73	Biological diversity, Fisheries,
	Scavenger biomass	3	5	5	1	5	19	73	Biological diversity, Sea-floor integrity
	Geometric mean abundance of seabirds	6	3	5	1	4	19	73	Biological diversity
	Size spectra slope	6	4	4	1	4	19	73	Biological diversity, Fisheries, Sea-floor integrity
	Fish biomass to benthos biomass from models	4	3	4	2	4	17	65	Biological diversity, Fisheries, Sea-floor integrity
	*Zooplankton spatial distribution and total biomass	4	4	3	2	4	17	65	Biological diversity, Eutrophication,
Zooplankton mean size	4	4	3	2	4	17	65	Biological diversity, Eutrophication	

---

Gini-Simpson diversity index	6	2	2	0	4	14	54	Biological diversity
Species richness index	6	2	2	2	2	14	54	Biological diversity

---



## 1144 **Appendix A:**

1145 Indicators that were examined (details, history, rationale).

### 1146 **Functional Indicators linked to Energy Flow**

#### 1147 **Seabird breeding success**

1148 Many species of seabirds feed on lower trophic level forage species such as krill, squid, and  
1149 pelagic fish. Seabirds summarize changes in these forage species communities that are often  
1150 linked to patterns of exploitation (Cury and Christensen, 2005; Cury et al., 2011). Seabird  
1151 breeding success has been consistently monitored across many ecosystems and provides robust  
1152 estimates of both forage fish abundance and success of charismatic species (Cury et al., 2011).  
1153 Seabird breeding success can be a useful indicator, however, it may overlap with other measures  
1154 of forage fish success.

#### 1155 **Productivity (production per unit biomass) of key predators**

1156 Metrics characterizing productivity of predators at high trophic levels have been identified by  
1157 Rogers et al. (2010) as an important class of food-web indicators. They argued that “[t]he  
1158 abundance of species in the food web will generally be determined by the abundance of suitable  
1159 prey taxa on which they can feed. Some species, or groups of species, may play a significant part  
1160 in food web dynamics and so their population status will effectively summarize the main  
1161 predator-prey processes in the part of the food web that they inhabit.” Food quantity or quality is  
1162 known to affect survival and reproduction of many marine species including birds (Wanless et  
1163 al., 2005), mammals (Soto et al., 2006) and fishes (Litzow et al., 2006). It has been argued (Boyd  
1164 and Charles, 2006; Cury et al., 2011; Rogers et al., 2010) that required prey abundance to  
1165 quantitatively and qualitatively sustain viable populations of predators constitutes a threshold  
1166 value which can serve as a reference point for productivity based indicators. “Productivity  
1167 (production per unit biomass) of key species or trophic groups” was listed among the Criteria for  
1168 GES by the EC (EU, 2010). Among others, it has been implemented in form of the HELCOM  
1169 (2013) core indicators “Pregnancy rates of marine mammals”, “White-tailed eagle productivity”,  
1170 “Abundance of sea trout spawners and parr”, and “Abundance of salmon spawners and smolt”.

#### 1171 **Mean weight at age of predatory fish species from data**

1172 Fish weight and condition metrics provide information on state (e.g., food limitation) in an  
1173 ecosystem. The indicator proposed by Shephard et al. (2014) describes the average “weight  
1174 anomaly” for the pelagic fish community in a given year, which is the deviation around an  
1175 observed long-term mean. The youngest and oldest age groups of each stock are excluded to  
1176 avoid sampling bias. Values are then averaged over all ages for each stock to obtain a mean  
1177 annual anomaly for that stock. Stock anomalies are then averaged by year to obtain a regional  
1178 mean weight anomaly for the whole pelagic or predatory fish communities, respectively, where  
1179 indicator values should fluctuate around zero in the long-term. The comparison between species

1180and stocks can give additional information on whether food becomes limiting in general or  
1181whether just some species or trophic guilds are impacted.

1182Changes in this indicator can be caused by changes in food availability as well as an increase or  
1183decrease in predator populations. The demand for food can be also influenced by temperature.  
1184Therefore, the indicator should only be interpreted in conjunction with additional information  
1185(e.g. biomass of forage fish, benthos, sea temperature, predator abundance, etc.). The indicator  
1186will respond predominantly to non-anthropogenic impacts and to a lesser degree to indirect  
1187anthropogenic impacts through food limitation.

#### 1188 **Total Mortality $Z$ (Production:Biomass ratio)**

1189Total mortality has a large effect on both year-to-year survival, long term reference points such  
1190as  $F_{MSY}$ , and resilience. If mean weight of species in the stock and catch remain constant over  
1191time, this indicator is conceptually equivalent to production/biomass. Further, the inverse of total  
1192mortality is a direct indicator of longevity, an indicator which is often more readily  
1193communicated outside the scientific community. It responds to management through direct  
1194fishing mortality and the abundance of predatory fish (ICES, 2013b).

#### 1195 **Primary Production Required to support fisheries**

1196The energy contained in solar radiation lead to primary production (PP) by phytoplankton and so  
1197fuels marine ecosystems. Subsequently, energy is transferred through food webs by predation  
1198and lost through metabolic processes. Ecosystem production results from the conversion of  
1199organic matter at each trophic level and is dependent on ecological features such as the number  
1200of feeding links, the efficiency of energy transfer from one trophic level to the next, and  
1201temperature (Chassot et al., 2010). Production available to fisheries depends upon fishing  
1202mortality and targeted trophic levels in the food web. Fisheries focusing only on lower trophic  
1203levels may be energetically more efficient than those focused on top predators (Gascuel and  
1204Pauly, 2009; Pauly and Christensen, 1995).

1205Primary Production Required (*PPR*) is the primary production and detritus flows from TL 1 that  
1206are required to sustain fisheries (expressed as t/km<sup>2</sup>/year). This allows the evaluation and  
1207comparison of fishing activities across ecosystems. The *PPR* is obtained by calculating the flows  
1208backwards, expressed in primary production and detritus equivalents, for all pathways from the  
1209caught species down to the primary producers and detritus. The *PPR* increases with fishing  
1210intensity. *PPR* has been analyzed also in reference to PP, to reflect a percentage of PP used to  
1211sustain catches.

#### 1212 **Productive pelagic habitat index (chlorophyll fronts)**

1213Productive fronts (chlorophyll-a fronts) are key large-scale features in marine ecosystems since  
1214they last long enough to sustain zooplankton production and are considered among the main

1215vectors of ocean productivity along the food chain (Belkin et al., 2009; Druon et al., 2012, 2011;  
1216le Fevre, 1986; Olson et al., 1994; Polivina et al., 2001).

1217The frequency of chlorophyll-a fronts with an intermediate range of chlorophyll-a content  
1218identifies the productive features that attract top-predators, i.e. areas of efficient energy transfer  
1219between trophic levels outside of low and high chlorophyll levels (from about 0.1 to 3.0 mg.m<sup>-3</sup>).  
1220Indeed, high chlorophyll levels potentially correspond to eutrophic areas where the food chain is  
1221disrupted and primary production is not available to upper trophic levels. Nutrient availability,  
1222hydrological and atmospheric forcing are captured by this indicator, but it captures in particular  
1223the variability of ecosystem productivity available to high trophic levels, independently of fishing  
1224pressure.

1225The indicator of pelagic productivity results from the demonstrated links between top-predators  
1226and chlorophyll-a fronts observed for fast-moving predators such as Atlantic bluefin tuna  
1227(Druon, 2010; Druon et al., 2011), and fin whale (Druon et al., 2012) and demersal nurseries in  
1228the Mediterranean Sea. The generic index of productive pelagic habitats yet requires a formal  
1229validation at European scale (<https://fishreg.jrc.ec.europa.eu/fish-habitat>).

### 1230 **Ecosystem exploitation (fisheries)**

1231This estimates the level of exploitation, integrated over all trophic levels, as the total yield  
1232divided by total production for all exploited species. Required data: Yield, biomass and  
1233production to biomass ratio for each exploited species.

### 1234 **Community Condition**

1235Community condition is a measure of the overall condition (average weight at length) at the  
1236functional group level, and the overall community condition. Condition reflects food availability:  
1237fish are heavier for their length when food abundance is plentiful and/or competition for food is  
1238low and lighter when food abundance is low and/competition for food is high. It is a reflection of  
1239energy flow, food availability and resilience.

### 1240 **Mean trophic level of the catch**

1241Mean trophic level of the catch is one of a suite of trophic level indicators that is based on the  
1242average biomass weighted trophic level across all species. Initial work considered the mean  
1243trophic level of the catch, based on fishery-dependent catch or landing statistics (Pauly et al.,  
12441998). It describes the average trophic level at which species are removed by the fisheries. As  
1245more valuable upper-trophic level fish stocks are depleted, fishers may target lower-value, lower-  
1246trophic level fish stocks (Pauly et al., 1998). Recent work suggests that this indicator is a better  
1247indicator of fishing pattern and pressure than an indicator of ecosystem state (Shannon et al.,  
12482014).

1249           **Marine trophic index of the community (MTI)**

1250The marine trophic index (MTI) (Pauly and Watson, 2005) is another trophic level indicator,  
1251calculated with a cut-off point of trophic level greater than 3.25. Originally calculated from  
1252fisheries landings data, here it is presented as the MTI of the community, based on scientific  
1253survey data, and is considered an indicator of food web functioning (Shannon et al., 2014). It has  
1254most commonly been applied to fish (and cephalopods), but could be extended to a wider range  
1255of taxa. The marine trophic index of the community, like the mean trophic level of the  
1256community (see below), provides a measure of ecosystem integrity and resilience. Declining  
1257trophic levels may result in shorter food chains, which may leave ecosystems less able to cope  
1258with natural or human-induced change.

1259           **Mean trophic level of the community**

1260Average trophic level (TL) obtained from fishery-independent surveys is a commonly used  
1261metric that can be used to measure status and trends of ecosystem structure and functioning(Shin  
1262et al., 2010). Average TL of the community is expected to decrease in response to fishing, as  
1263fisheries tend to target species at upper trophic levels (Pauly et al., 1998). Additionally, fishing  
1264can also change the structure of marine food webs by reducing the mean TL and might also  
1265influence ecosystem functioning by shortening the length of food chains and releasing predation  
1266on lower trophic level organisms (Shin et al., 2010).

1267           **Disturbance index**

1268The disturbance index (DI) measures the change in trophic (or size) structure of the ecosystem  
1269and is calculated as the sum, across all TLs  $\geq 2$  (or size classes), of the absolute difference in the  
1270relative biomass ( $B_{TL}/B_{Total}$ ) within each TL for each year, relative to a reference period (Bundy  
1271et al. 2005). The reference period can represent a preferred state of the ecosystem, an ideal state,  
1272a theoretical state estimated from an ecosystem model or the beginning to the time period for  
1273which there is data. The DI has been shown to respond directly to fishing pressure, but may also  
1274be affected by other pressures such as environmental change.

1275The DI was originally proposed as one of 4 indicators comprising a 4D ecosystem exploitation  
1276index (Bundy et al., 2005).

1277           **Loss in secondary production index (L index)**

1278The decrease in secondary production was proposed as a proxy for quantifying ecosystem effects  
1279of fishing on the basis of a theoretical development and application to a large set of data  
1280(Libralato et al., 2008). The L index is calculated by integrating the primary production required  
1281to sustain the catches (PPR: Pauly and Christensen, 1995) relative to the primary production (PP)  
1282in the ecosystem, taking account of transfer efficiencies (TE, i.e., the efficiency in the transfer of  
1283energy from a trophic level to another; Lindeman, 1942) and the trophic level of the catches  
1284(TLc; Pauly et al., 1998). Theoretically, these inputs can be combined to measure the loss in

1285secondary production due to fishing (L index) and to evaluate ecosystem effects of all fished  
1286species (Libralato et al., 2008).

1287The application of the L index to a set of well-studied models allowed a probability of being  
1288sustainably fished ( $P_{\text{sust}}$ ) to be associated with each L index value, and, by fixing desired  
1289sustainability levels (e.g., 75% and 95%) it provide the basis for back-estimating the associated  
1290Ecosystem-based Maximum Sustainable Catches (EMSC) (Libralato et al., 2008).

1291Thus L index is formally defined as an index of ecosystem overfishing and allows application of  
1292the index using both landings data and ecosystem models. L index can give rough estimates of  
1293overfishing status and management advice measures allowing definition of a region of viable  
1294solutions (Cury et al., 2005). L index quantification can be adapted to specific spatial scales  
1295(regional spatial assessment) and to large pelagic areas exploiting data from satellite for  
1296estimating PP, catches and available data on diets (for TL estimates).

### 1297 **Cumulative distribution of biomass assessment**

1298Accumulation of biomass has been documented for many marine food webs, with the  
1299intermediate TLs exhibiting the largest increase in the system cumulative biomass (Gascuel and  
1300Pauly, 2009; Link et al., 2009). Changes in this accumulation may reflect shifts in the ecosystem  
1301structure and function. According to these observations, from a theoretical point of view, a  
1302perturbed ecosystem should lower the stored, cumulative biomass and “stretch out” across TLs.  
1303To describe and quantify these changes, the biomass distribution across TLs is fitted to a logistic  
1304nonlinear regression model to estimate the main curve parameters: steepness (that is the slope of  
1305the tangent passing through the inflection point), inflection TL (that is the projection of the  
1306inflection point on the x-axis), inflection CumB (that is the projection of the inflection point on  
1307the y-axis), and the basal biomass (that is the y-axis intercept of the fitted curve). Tests, carried  
1308out by using both surveys and landings data, showed that the method is robust to possible  
1309'sampling errors' (in terms of TL assignment), sensitive to both environmental and anthropogenic  
1310drivers, and when applied to fishery dependent data, responsive (Pranovi et al., 2014, 2012).

### 1311 **Trophic balance index (fishing pattern)**

1312This index measures the evenness (pattern) of exploitation across TLs by comparing their  
1313exploitation rates, which are estimated as the sum of yield (Y) divided by the sum of production  
1314(P) at each TL. The evenness of exploitation is then given by the coefficient of variation of all Y/  
1315P. Required data: Yield, biomass and P/B for each species in the yield.

### 1316 **Mean transfer efficiency for a given TL or size**

1317The transfer efficiency ( $TE_{\text{TL}}$ ) is defined as the fraction of production that is passed from one  
1318integer trophic level to the next (Lindeman, 1942; Pauly and Christensen, 1995). It is thus  
1319quantifiable as the ratio between the production of the trophic level (TL) and the production at  
1320the precedent trophic level (TL-1). Several studies have estimated the pattern of TE by different

1321 trophic level after Lindeman's work (Burns, 1989; Lindeman, 1942). It has been used as a  
1322 diagnostic indicator in some cases (e.g. Libralato et al., 2004) but in most instances the  
1323 ecosystem average is used as an integrated summary statistic.

#### 1324 **Finn Cycling Index**

1325 The Finn's cycling index (FCI: Finn, 1976) is the proportion of the total sum of flows in the food  
1326 web that is recycled in the system. It is measured as the proportion of the total flow that is  
1327 flowing within circular pathways. Recycling is considered to be an indicator of an ecosystem's  
1328 ability to maintain its structure and integrity through positive feedback and is used as an  
1329 indicator of stress and maturity (Christensen, 1995; Monaco and Ulanowicz, 1997; Ulanowicz,  
1330 1992; Vasconcellos et al., 1997). FCI is an indicator of the recovery time of an ecosystem  
1331 through development of routes to conserve nutrients. A high FCI would mean the system would  
1332 recover faster from a perturbation, whereas a system would be expected to take longer to recover  
1333 (lower FCI) when it is in a more degraded state.

#### 1334 **Functional Indicators linked to ecosystem resilience**

##### 1335 **Mean trophic links per species**

1336 The mean number of trophic links per species reflects how connected a food web is and,  
1337 potentially, how stable a food web may be (Link, 2002; Link, 2005; Methratta and Link, 2006).  
1338 Changes to this indicator reflect notable differences in the structure and dynamics of a food web.  
1339 As an understanding of temporal and spatial characteristics of marine trophic interactions it may  
1340 not be entirely complete. This index should be used only as a tool to invoke further precautionary  
1341 action (Link, 2005).

##### 1342 **Ecological Network Analysis derived indicators (overall mean Transfer** 1343 **Efficiency)**

1344 The mean transfer efficiency ( $TE_m$ ) for the food web is calculated as the geometric mean of  
1345 transfer efficiencies for each of the integer trophic levels II to IV from models (Christensen and  
1346 Pauly, 2008; Christensen et al., 2009). It is a variant of the mean transfer efficiency discussed  
1347 above. There have been attempts to estimate average TE also on the basis of catches over trophic  
1348 levels on the assumption that fisheries were in balance for some periods (Pauly and Palomares,  
1349 2005) – which would provide a fishing pressure indicator. Average transfer efficiency by  
1350 ecosystem type based on model outputs have shown some variability across ecosystem types  
1351 (Libralato et al., 2008) and other pressures as shown in Heymans et al. (2012). The indicator has  
1352 been proposed as a descriptor of ecosystem health in lakes (Xu and Mage, 2001).

##### 1353 **Gini-Simpson dietary diversity index**

1354 The Gini-Simpson dietary diversity index is defined as the average, over a representative sample  
1355 of consumer species, of the Gini-Simpson diversity of the contributions of resource species to  
1356 consumer diets, by volume or biomass (ICES, 2013b; Rossberg, 2013; Rossberg et al., 2011). It

1357 can be determined from stomach-content data. The metric attains values between 0 and 1, with 0  
 1358 implying no diversity and 1 high diverse. In practice it is computed as

1359 
$$D_{diet} = 1 - \frac{\sum_{ij} [p_{ij}]^2}{\sum_{ij} p_{ij}}$$

1360 where  $p_{ij}$  is the proportional contribution of species  $i$  to the diet of  $j$ , and the sums run over all  
 1361 diet items resolved to species level (Rossberg, 2013; Rossberg et al., 2011). The indicator may  
 1362 be applied to any component of the ecosystem for which diet data is available, but has so far  
 1363 been computed only for fish (Rossberg et al., 2011). A target for the metric near 0.5 has been  
 1364 proposed (Rossberg, 2013; Rossberg et al., 2011) based on theory and observation data. The  
 1365 indicator may respond to pressures (e.g. Rossberg et al., 2011).

1366 **Herbivory : detritivory ratio**

1367 This indicator, proposed by (Ulanowicz (1992), is the ratio of the values of the detritivory flow  
 1368 (from detritus to level II) divided by the value for the herbivory flow (from primary producers to  
 1369 level II). It is sometimes presented as H/D (or abbreviated HDR). This indicator was inspired by  
 1370 Lindeman (1942) when he referred to the role of saprophageous organisms and heterotrophic  
 1371 bacteria. This ratio has already been tested as a candidate for defining functional indicators of the  
 1372 food web, but results seem to be case sensitive. For example, Ulanowicz (1992) observed a  
 1373 higher H/D ratio in disturbed situations whereas Dame and Christina (2007) observed exactly the  
 1374 opposite trend. Then the disturbed situation showed a shift to a more detritus-based food web.

1375 **Ecological Network indices of ecosystem status and change (Ulanowicz)**

1376 Redundancy ( $R$ ) (Monaco and Ulanowicz, 1997) indicates the system's energy in reserve it  
 1377 describes the distribution of energy flow among the ecosystem pathways, and is an indicator of  
 1378 change in the degrees of freedom of the system (Heymans et al., 2007). Based on the description  
 1379 of  $R$  by (Ulanowicz, 2004), who suggested that, "... it strongly ties to the effective multiplicity  
 1380 of parallel flows by which medium passes between any two arbitrary system components".  
 1381 Redundancy is linked by Christensen and Pauly (2008) with system stability and proposed by  
 1382 Heymans et al. (2007) as an index of food-web resilience. According to Bondavalli et al. (2000)  
 1383 high redundancy signifies that either the system is maintaining a higher number of parallel  
 1384 trophic channels in order to compensate for the effects of environmental stress, or that it is well  
 1385 along its way to maturity. With regard to overall performance and robustness, ecosystem level  
 1386 indicators based on ecological network analysis and food-web analysis are informative on  
 1387 intermediate and long time-scales (Cury et al., 2005; IEEP, 2005; Moloney et al., 2005). But they  
 1388 are difficult to use in annual updates and operational approach, and may be more difficult for  
 1389 stakeholders to understand (IEEP, 2005). In addition, use of food-web models and the ecological  
 1390 network analysis approach to explore different management scenarios, through simulation of  
 1391 fishery and nutrient management, could deliver integrated views at ecosystem level.

## 1392 **System omnivory index**

1393 The system omnivory index (SOI) measures the distribution of feeding interactions among  
1394 trophic levels of food webs, thus SOI evaluates the complexity and connectivity of food webs. It  
1395 has been associated to ecosystem ability to recover from perturbations (Christensen, 1995;  
1396 Libralato, 2008). Given a food web with  $n$  elements, the SOI is calculated as the weighted  
1397 average of the elements' omnivory, the latter calculated as the omnivory index (OI). The OI  
1398 of each consumer element  $i$  with trophic level  $TL_i$  is quantified as the variance of the trophic  
1399 levels of its preys ( $TL_j$ ) (Williams and Martinez, 2004). The SOI of a given trophic network is  
1400 quantified as the weighted average of the OI of all consumers of the network, where the  
1401 weighting factors are taken as the logarithm of each consumer food intake ( $Q_i$ ) (Christensen and  
1402 Pauly, 1993). This allows for accounting of the different strengths of consumer interactions and  
1403 the logarithm is used on the observation that consumption is approximately log-normally  
1404 distributed within systems (Christensen and Pauly, 1993).

1405 The topological configuration of links and their weights affect SOI, but it is quite robust to the  
1406 number of nodes in the web (Libralato, 2008). Comparison of stability and complexity indices  
1407 including SOI for coastal marine food webs highlighted positive correlation between SOI,  
1408 magnitude of change and recovery time, thus suggesting that SOI is inversely related to stability  
1409 of marine ecosystems (Perez-Espana and Arreguin-Sanchez, 1999). Moreover, application of  
1410 SOI and other ecological indicators, on the basis of outputs of protected and fished marine food  
1411 webs standardized by number of elements, suggests that SOI is sensitive to fishing (Libralato et  
1412 al., 2010).

## 1413 **Structural Indicators linked to diversity and 'canary' species**

### 1414 **Guild Surplus Production models**

1415 Guild Surplus Production is tracked in the annual Ecosystem Assessment document for the North  
1416 Pacific Fisheries Management Council (Zador, 2013). Species are grouped into functional guilds  
1417 based on feeding and life history studies. Survey and catch time series for each species are used  
1418 to calculate the surplus production for each guild. To use as a catch limit, in addition to a single-  
1419 species limit for each managed stock, the sum of quotas for each guild cannot exceed the MSY  
1420 for the guild as defined by a standard surplus production model. Per-species reductions to meet  
1421 this overall limit are not proscribed by this index; reductions can be made for stakeholder or  
1422 economic reasons. For Bering Sea (ecosystem-wide) indicator example, see Meuter and Megrey  
1423 (2006). The indicator uses is based on survey biomass and catch of the species within each  
1424 guild.

### 1425 **Total biomass of small fish**

1426 This indicator uses survey catch biomass of predefined small (pelagic) fish to assess exploitation  
1427 levels of commercial stocks. The amount of energy transferred from zooplankton to higher  
1428 trophic levels by pelagic fish is ultimately limited by the biomass of pelagic fish available.



1429Shephard et al. (2014) therefore suggest that both the biomass of individual stocks should be  
1430above precautionary reference points on average and the total stock biomass of all pelagic fish  
1431together should be above a joint community reference point. In practice, the community  
1432reference point is always reached when all individual stocks are above precautionary reference  
1433levels. However, in the case where one or more stocks are substantially below single stock  
1434reference points, additional care should be taken in the exploitation of the remaining stocks in the  
1435area.

#### 1436 **Proportion of Predatory Fish**

1437Predatory fish species are defined as all surveyed fish species that are not largely planktivorous  
1438(i.e. phytoplankton and zooplankton feeders should be excluded: Shin et al., 2010). A fish  
1439species is classified as predatory if it is piscivorous, or if it feeds on invertebrates that are larger  
1440than the macrozooplankton category (0.2 cm). Detritivores should not be classified as predatory  
1441fish. This indicator captures changes in the trophic structure and changes in the functional  
1442diversity of fish in the ecosystem. It is sensitive to fishing pressure, but since it is a ratio, it will  
1443also be subject to changes in non-predatory fish, whose biomass may vary for other reasons (i.e.  
1444environmental driver: Bundy et al., 2010).

1445This indicator is calculated as the biomass of predatory fish surveyed / biomass surveyed, and the  
1446data required are trawl survey data and food habits data (if not available locally, from  
1447information in the literature or from comparable systems).

#### 1448 **Pelagic to demersal ratio**

1449The ratio of pelagic to demersal fish (P:D ratio) obtained from fishery-dependent or -independent  
1450surveys is a commonly used metric that describes trophic energy flow and community structure  
1451(de Leiva Moreno et al., 2000; Link, 2005; Rochet and Trenkel, 2003). Changes in P:D ratio  
1452have been linked to anthropogenic pressures such as fishing and eutrophication. Targeted fishing  
1453can result in notable shifts in this indicator, however, changes may not be entirely clear, as an  
1454increase in the P:D ratio could be caused by an increase in pelagic fish or a relative decrease in  
1455demersal fish. As an indicator of food web properties, P:D ratio may overlap with other large  
1456and/or forage fish indicators, but does capture important trophic relationships.

#### 1457 **Biomass of trophic guilds**

1458Biomass of trophic guilds is a measure of ecosystem structure, estimated as the aggregate  
1459biomass of each trophic guild. Individually they provide a measure of the change in biomass of  
1460trophic guilds. Collectively, they provide a measure of change in overall structure. It can be  
1461applied to all marine species if the information is available, based on survey data or model  
1462results. Work to date has largely focused on fish trophic guilds (Rochet et al., 2013; Shackell et  
1463al., 2012), but could be extended to invertebrates birds, and marine mammals. Measures of  
1464functional diversity could also be developed using these data. Data sources can be from research  
1465surveys or models.

1466           **Lifeform-based indicator for the pelagic habitat**

1467 Ecosystem health theory (reviewed by Tett et al., 2013) suggests that ecosystem resilience, and  
1468 the sustainability of services, depends *inter alia* on the abundance and relationships of non-  
1469 substitutable 'functional groups' or 'lifeforms'. The abundances and trophic structural  
1470 relationships of phytoplankters, and their protozoan and mesozooplankton consumers, change  
1471 seasonally. The Plankton Index (Pi) method takes account of such seasonality and requires the  
1472 plotting of log-transformed lifeform abundances, based on at least monthly samples, in sets of 2-  
1473 D state spaces (Gowen et al., 2011). These plots (Tett et al., 2008) often suggest a fuzzy  
1474 doughnut. Using data for a reference period, an envelope can be drawn to include a fixed  
1475 proportion (usually 90%) of points in this doughnut. Data from other years can be plotted against  
1476 this envelope; the  $Pi[j,t]$  value (for lifeform pair  $j$  and year  $t$ ) is defined as the proportion of new  
1477 points that fall inside the envelope. For a given value of  $t$ , values of  $Pi$  for different lifeform pairs  
1478 can be averaged. A UK project has identified sets of lifeform pairs that may serve for  
1479 assessment of environmental status. The lifeform pairs relevant to Food Webs are: (i) chlorophyll  
1480 concentration and mesozooplankton abundance; (ii) phytoplankton  $\geq 20 \mu\text{m}$  abundance and  
1481 phytoplankton  $< 20 \mu\text{m}$  abundance; (iii) [adult] copepods  $\geq 2 \text{ mm}$  abundance and [adult]  
1482 copepods  $< 2 \text{ mm}$  abundance. Reference conditions for any of the  $Pi$  are expected to be  
1483 dependent on ecohydrodynamic (EHD) conditions (van Leeuwen et al., 2015). The UK is  
1484 currently seeking EHD-specific references at sites in the Celtic or Greater North Sea MSFD  
1485 ecoregions that are, according to expert judgement, in GES. Meanwhile, time-series of  $Pi$  will be  
1486 generated from conditions observed during an agreed (but arbitrary) period of 3 years, and the  
1487 time series will be assessed for (a) significant trends, and (b) significant correlation with relevant  
1488 pressures.

1489           **Region-specific indicators of abundance & spatial distribution,**

1490 Indicators can be selected to track the abundance and spatial distribution of major species which  
1491 represent key community and or/ecosystem properties. Ideally, species representing different  
1492 communities or habitats (benthos, plankton, fish, top predators) should be selected, in this way  
1493 covering a large part of the ecosystem. As ecosystems are typically characterized by few strong  
1494 links and many weak links among species or trophic levels, one (or few) indicator populations  
1495 can describe broader ecosystem state and/or human perturbation. Criteria in the MSFD for  
1496 selecting the groups/species that could be included in this category are those with fast turnover  
1497 rates, groups/species that are targeted by fisheries, the habitat-defining groups/species, those at  
1498 the top of the food web, and those tightly linked to other trophic levels (Rogers et al., 2010).

1499           **Fish biomass : benthos biomass from models**

1500 Ratios are used to measure changes in community structure indicating the distribution of energy  
1501 in the ecosystem. They are a supplement to biomass indicators and have the advantage that they  
1502 do not reflect general increases or decreases in biomass in all components but only changes in  
1503 the relative importance between the two groups. Hence, the ratio pelagic biomass : demersal

1504biomass represents the balance between pelagics and demersals whereas the fish/benthos ratio  
1505reflects the proportion of the biomass which is diverted to benthos, including detritivores. The  
1506indicator captures changes in the trophic structure and changes in the functional diversity of the  
1507ecosystem. It is sensitive to fishing pressure, but since it is a ratio, it will also be subject to  
1508changes in non-manageable benthos, whose biomass may vary for other reasons (e.g.  
1509environmental driver, Bundy et al., 2010). Data sources can be research surveys (mainly nekton)  
1510or models (often benthos, since this is often not surveyed on appropriate spatial and temporal  
1511scales).

#### 1512 **Zooplankton spatial distribution and total biomass**

1513This indicator, which describes the distribution of zooplankton, is still at the developmental stage  
1514with methods, threshold and target values to be developed. The reasoning for this indicator is  
1515that zooplankton constitutes an important link between primary producers and higher trophic  
1516levels in the food web. Zooplankton plays an important role in the energy transfer and nutrient  
1517cycling in the food web. Changes of the composition of the zooplankton community are coupled  
1518to environmental changes and can respond quickly to ecosystem changes. Zooplankton biomass  
1519and abundance can e.g. respond to invasive species and local oil spills.

#### 1520 **Scavenger biomass**

1521Fishery discards provide food subsidies that help maintain fish and seabird populations and may  
1522allow some of these populations to be more abundant than they would naturally be (e.g. Link and  
1523Almeida, 2002; Polis and Strong, 1996). Surveys of non-targeted scavenger biomass or  
1524abundance may provide an index of disturbance (Link and Almeida, 2002; Methratta and Link,  
15252006b). Additionally, some scavenger species might be viewed as “canary” or “iconic” species  
1526that can be used as an early warning of disturbance or excessive fishing pressure.

#### 1527 **Geometric mean abundance of seabirds**

1528The Geometric Mean Abundance of Seabirds is computed in regular intervals (e.g. yearly) as the  
1529geometric mean of the population sizes (e.g. numbers of individuals or breeding pairs) of those  
1530seabirds in the assessment region for which population time series are available, normalised such  
1531that the indicator value at the beginning of the indicator time series is one. The indicator is  
1532designed after the Living Planet Index (LPI: Loh et al., 2005), which now underlies Aichi Target  
15335 of the Convention for Biological Diversity. Modern indicator protocols take into account that  
1534species may enter or leave the set of species for which time series are available, and that  
1535population sizes at low abundances become uncertain. Methods to compute indicator confidence  
1536intervals have been developed (Buckland et al., 2011; Loh et al., 2005). By its definition, the  
1537proportional rate of change of the indicator equals the average population growth rate of all  
1538populations contained in the indicator (here seabirds). Under conditions where populations  
1539fluctuate and turn over but overall biodiversity does not change, the indicator is expected not to  
1540deviate significantly from one. A steady decline of geometric mean abundance signals

1541biodiversity loss. Seabird populations are known to be highly sensitive to food availability (Cury  
1542et al., 2011), and their differentiation of foraging niches (Fasola et al., 1989) is evidence of  
1543competition for food among them. Competitive exclusion resulting from loss of biodiversity  
1544among their marine resources (e.g. forage fish), or even at lower trophic levels (Rossberg, 2013),  
1545can be expected to induce the slow decline of seabird diversity to which this indicator is designed  
1546to be sensitive. Geometric mean abundance of seabirds is therefore sensitive to a collapse of the  
1547pyramidal distribution of species over trophic levels in food webs (de Ruiter et al., 2005).

#### 1548 **Gini-Simpson diversity index (species dominance) of large and small fish by** 1549 **biomass**

1550It is incompatible with GES to bring the foodweb into a state where only a few (large) predator  
1551or prey species dominate the system when the biomass of predators and prey was distributed  
1552more evenly in the system during the reference period. Species richness may be inadequate as an  
1553indicator as it often takes a long time to completely lose a species, while management should be  
1554informed and act earlier. The Gini-Simpson index (1-D) applied to the predator and/or prey  
1555community provides the possibility to detect unwanted changes in diversity. Simpson's Diversity  
1556Index is a measure of diversity which takes into account the number of species present, as well as  
1557the relative abundance of each species. As species richness and evenness increase, so does  
1558diversity (ICES, 2013b).

#### 1559 **Species Richness Index**

1560Species richness measures the number of species within a community. A well-structured and  
1561functioning ecosystem will generally have many species (Fung et al., 2015); as a side effect of  
1562fishing, species richness may decrease (Rice, 2003, 2000). However, as a food web indicator  
1563species richness may provide ambiguous information, since multiple community configurations  
1564may produce similar values, as shown by Gislason and Rice (1998) and Rice and Gislason  
1565(1996). In these studies the index was calculated as the number of species in any year whose  
1566numerical abundance or biomass was larger than some percentage of their value in a reference  
1567year. The IUCN Red List criterion of 20% was used as the reference value. Required data:  
1568species or functional group P/B, and species or functional group biomass/abundance to compare  
1569to reference points.

#### 1570 **The Large Fish Indicator (LFI)**

1571The Large Fish Indicator (LFI) is defined as the proportion by weight of large fish in the sample  
1572of a specified survey (Greenstreet et al., 2011), where large fish are defined as those longer than  
1573a threshold length  $L_{th}$ , a region-specific value. The threshold value is chosen such as to optimize  
1574the responsiveness of the indicator to fishing pressure, as determined from historic data  
1575(Shephard et al., 2011). The LFI takes no account of species identity, only of individual sizes.  
1576However, it was shown to reflect mostly the proportion (by weight) of large-bodied species in  
1577communities (Shephard et al., 2012). Large-bodied species tend to be more vulnerable to fishing,

1578which is why the LFI is sensitive (Engelhard et al., 2015; Greenstreet et al., 2011; ICES, 2011;  
1579Shephard et al., 2013) and specific (Houle et al., 2012) to fishing pressure. Furthermore, by  
1580expressing the indicator in terms of proportions by weight, and not by numbers, and through  
1581judicious choice of the appropriate length threshold to define large fish, the indicator can be  
1582desensitized to variation in the abundance of small fish. The influence of environmentally driven  
1583recruitment events on indicator values can therefore be minimized (Greenstreet et al., 2011).  
1584Food-web models (Fung et al., 2013; Shephard et al., 2013) and data (Fung et al., 2012) suggest  
1585that recovery of the indicator from pressures can be slow (decadal scale). The LFI, as an OSPAR  
1586EcoQO for the North Sea, is fully operational. It was named as an indicator for food-web GES  
1587(EU, 2010), and has been chosen as a common food-web indicator by HELCOM and OSPAR (in  
1588some OSPAR Subregions as a priority candidate indicator).

### 1589        **Mean length of surveyed community**

1590Mean length (ML) of all species caught in a survey, whether fishery-independent, fishery-  
1591dependent, or based on landings, can be a useful and simple indicator to evaluate the overall  
1592effects of fishing on an ecosystem (Dulvy et al., 2004; Nicholson and Jennings, 2004; Rochet  
1593and Trenkel, 2003; Shin et al., 2005). ML quantifies relative abundances of large and small  
1594individuals and describes the size distribution of a community (Shin et al., 2005). It is relatively  
1595responsive to key pressures (Link, 2010; Pauly et al., 1998). ML is considered measurable and  
1596generally robust, however, the direction of response may be caused by increasing stocks of large  
1597fish or decreasing in stocks of small fish, leading to potential ambiguity. Whilst the metric is  
1598sensitive to fishing pressure, it can also be strongly influenced by environmentally driven  
1599recruitment events that introduce large numbers of small fish into the community (Badalamenti et  
1600al., 2002; Lekve et al., 2002; Wilderbuer et al., 2002).

### 1601        **Size spectrum slope**

1602Various measures of the change in size can be a useful indicator to describe composition of  
1603communities (Nicholson and Jennings, 2004). Size spectrum slope measures the relationships  
1604between the biomass (y) of individuals within a body size class and body size (x), both normally  
1605plotted on logarithmic scales. Frequently a logarithmic transformation is applied to body size,  
1606particularly when weight classes are used. When applied to fish communities, the slope of the  
1607relationship becomes increasingly negative in response to fishing pressure; fisheries reduce the  
1608abundance of large fish, through the direct effect of fishing and, as a consequence of reduced  
1609predation pressure from large fish, the abundance of small fish increases (Daan et al., 2005;  
1610Gislason and Rice, 1998; Nicholson and Jennings, 2004; Rice and Gislason, 1996). The size  
1611spectrum slope is considered measurable and robust. However, the direction of the response may  
1612not be entirely clear (Trenkel and Rochet, 2003), as the steepening of the slope could indicate a  
1613decrease of large fish or an increase of small fish. The slope is particularly sensitive to changes  
1614in the abundance of small fish, which markedly affect the intercept of the regression line, as such

1615the size spectrum slope can be influenced by environmentally driven recruitment events  
1616(Badalmenti et al., 2002; Lekve et al., 2002; Wilderbuer et al., 2002).

1617        **Zooplankton Size-Biomass index**

1618This is a zooplankton indicator reflecting both mean individual size and total biomass of the  
1619zooplankton community. The indicator represents food web capacity to sustain fish feeding  
1620conditions and grazing on primary producers. The rationale is that both mean body size in the  
1621community and total community biomass are positively related to fish feeding conditions,  
1622whereas total biomass alone is just representative of grazing pressure and trophic transfer  
1623efficiency (Fuchs and Franks, 2010). The effects of zooplankton community structure on energy  
1624transfer and food web resilience have been demonstrated in both freshwater and marine systems  
1625(Jeppesen et al., 2011; Kane et al., 2009; Lougheed and Chow-Fraser, 2002). The index is  
1626currently considered as a core indicator for the Baltic Sea (HELCOM, 2013, 2012). In semi-  
1627enclosed seas, such as the Baltic Sea, with strong salinity and temperature gradients, no single  
1628zooplankton group can adequately reflect community properties (Remm 1984), hence the need  
1629for this two-dimensional index. The index value decreases with increasing fishing pressure.  
1630Protocols for indicator assessment have been developed by HELCOM Zooplankton Expert  
1631Network (ZEN) using nine long-term monitoring datasets in the Baltic Sea (HELCOM, 2013,  
16322012). In all datasets, the indicator was found to predict deviations from GES conditions.  
1633Determination of GES boundaries for the indicator is straightforward and based on the regional  
1634basin-specific Environmental Quality Ratios for chlorophyll accepted within Water Framework  
1635Directive and weight-at-age for zoo-planktivorous fish (HELCOM, 2013, 2012).