

## Research



**Cite this article:** Faulkes CG, Eykyn TR, Aksentijevic D. 2019 Cardiac metabolomic profile of the naked mole-rat—glycogen to the rescue. *Biol. Lett.* **15**: 20190710. <http://dx.doi.org/10.1098/rsbl.2019.0710>

Received: 25 September 2019

Accepted: 25 October 2019

### Subject Areas:

cellular biology, molecular biology, systems biology, biochemistry, developmental biology, behaviour

### Keywords:

naked mole-rat, cardiac metabolism, metabolomics, glycogen, hypoxia

### Author for correspondence:

Dunja Aksentijevic

e-mail: [d.aksentijevic@qmul.ac.uk](mailto:d.aksentijevic@qmul.ac.uk)

<sup>†</sup>Authors contributed equally to this work.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.4725266>.

# Cardiac metabolomic profile of the naked mole-rat—glycogen to the rescue

Chris G. Faulkes<sup>1,†</sup>, Thomas R. Eykyn<sup>2,†</sup> and Dunja Aksentijevic<sup>1</sup>

<sup>1</sup>School of Biological and Chemical Sciences, Queen Mary University of London, G.E. Fogg Building, Mile End Road, London, UK

<sup>2</sup>Department of Imaging Chemistry and Biology, School of Biomedical Engineering and Imaging Sciences, King's College London, St Thomas Hospital, London, UK

**id** CGF, 0000-0002-5228-9075; TRE, 0000-0003-1768-3808; DA, 0000-0002-8480-6727

The African naked mole-rat (*Heterocephalus glaber*) is unique among mammals, displaying extreme longevity, resistance to cardiovascular disease and an ability to survive long periods of extreme hypoxia. The metabolic adaptations required for resistance to hypoxia are hotly debated and a recent report provides evidence that they are able to switch from glucose to fructose driven glycolysis in the brain. However, other systemic alterations in their metabolism are largely unknown. In the current study, a semi-targeted high resolution <sup>1</sup>H magnetic resonance spectroscopy (MRS) metabolomics investigation was performed on cardiac tissue from the naked mole-rat (NMR) and wild-type C57/BL6 mice to better understand these adaptations. A range of metabolic differences was observed in the NMR including increased lactate, consistent with enhanced rates of glycolysis previously reported, increased glutathione, suggesting increased resistance to oxidative stress and decreased succinate/fumarate ratio suggesting reduced oxidative phosphorylation and ROS production. Surprisingly, the most significant difference was an elevation of glycogen stores and glucose-1-phosphate resulting from glycogen turnover, that were completely absent in the mouse heart and above the levels found in the mouse liver. Thus, we identified a range of metabolic adaptations in the NMR heart that are relevant to their ability to survive extreme environmental pressures and metabolic stress. Our study underscores the plasticity of energetic pathways and the need for compensatory strategies to adapt in response to the physiological and pathological stress including ageing and ischaemic heart pathologies.

## 1. Introduction

The naked mole-rat (NMR) (*Heterocephalus glaber*) is a mouse-sized eusocial African rodent that displays a range of unusual physiological characteristics from resistance to cardiovascular disease to extreme longevity [1–3]. Unlike other mammals, they do not conform to Gompertzian laws of age-related mortality as adults show no age-related change in mortality risk [4]. NMRs live in colonies that may number up to 300 individuals, in extensive underground burrows. Although parts of their burrows may often be normoxic, oxygen is likely to become scarce in crowded nest chambers where animals huddle together and sleep [4,5]. To date, the gaseous composition of an NMR nest has not been sampled in the wild. However, these nest chambers are often more than 0.5 m deep, and may be some distance from brief openings to the surface at the ephemeral mole-hills [6]. NMRs are thus adapted to be resistant to hypoxia and in laboratory experiments are able to survive anoxia for up to 18 min [2]. The mechanism underlying this may be their ability to switch from glucose

to fructose driven glycolysis in the brain as the source of lactate. Other studies have revealed a number of cellular adaptations enabling tolerance to hypoxia in the brain [7–10]. The NMR can undergo rapid increases in metabolic rate to meet energy demand associated with digging through compacted soils in its xeric natural habitat with patchy food distribution [11]. They display a low baseline metabolic rate, and a recent study by Pamerter *et al.* [10] has shown that NMRs exhibit a clear decrease in metabolic rate in situations of acute hypoxia, emphasizing their ability to physiologically react to the prevailing conditions. NMRs also have low basal cardiac function accompanied by morphological traits such as cardiomyocyte hypertrophy, which is commonly associated with cardiac pathology in the Murinae and in humans [12–15]. However, they do not develop the cardiac disease and unlike pathologically remodelled hearts have enhanced contractile reserve upon increased demand [12–15]. What fuels these critical functional adaptations as well as what kind of metabolic adaptations and associated mechanisms render NMR hearts resistant to hypoxic injury and senescence remain unknown.

## 2. Methods

### (a) Animals

The non-breeding male adult NMRs used in this study were second-generation or more captive-born, descended from animals captured in Kenya in the 1980s. Colonies were maintained using artificial burrow systems as previously described [16]. The ages selected for this study allowed for physiological age matching between species such that both were at equivalent percentages of maximum lifespan and therefore not the same chronological age.

### (b) High resolution $^1\text{H}$ nuclear magnetic resonance spectroscopy metabolomic profiling

Myocardial tissue was collected from adult, non-breeding NMRs ( $n=5$  males, body weight  $37 \pm 7$  g, age  $\sim 7$  years), adult C57/BL6 mice ( $n=5$ , Charles River, UK, male, 27 g, 9 weeks) and adult Wistar rats ( $n=5$ , Charles River UK, male 450 g, 10 weeks) post-euthanasia. Frozen, weighed and pulverized hearts were subject to methanol/water/chloroform dual-phase extraction and high resolution  $^1\text{H}$  nuclear magnetic resonance spectroscopy (MRS) metabolomic profiling adapted from Chung *et al.* [17] (described in the electronic supplementary material).

## 3. Results

High resolution  $^1\text{H}$  MRS based metabolomic analysis of NMR heart tissue (representative spectra shown in figure 1) revealed distinctly different metabolomic profiles compared to C57/BL6 hearts (figure 2b) and changes in several key metabolites were observed (figure 2c,d). Despite reduced contractile performance and baseline hypertrophy [12–15], NMR hearts were not energetically compromised as the levels of creatine and ATP were comparable to C57/BL6 wild-type hearts. Myocardial glycogen and glucose-1-phosphate levels (G-1-P) resulting from glycogen turnover were significantly higher in the NMR hearts and were undetectable in C57/BL6 mouse hearts ( $p < 0.001$ ; figure 2e). Liver glycogen content was also found to be markedly higher in NMRs than C57/BL6 ( $4.1 \pm 1.1$  versus  $0.31 \pm 0.08 \mu\text{mol g}^{-1}$ ;

$p < 0.001$ ). Surprisingly myocardial glycogen in the NMR was even greater than the mouse liver which is the principal glycogen storage organ with highest intracellular deposits ( $1.2 \pm 0.02$  versus  $0.4 \pm 0.08 \mu\text{mol g}^{-1}$ ;  $p < 0.001$ ). A significant difference in myocardial glycogen levels was also observed in NMRs compared to Wistar rats, the latter was not significantly different to that found in C57/BL6 mice (thus representing another hypoxia-intolerant rodent). Glycogen regulation involves a complex interplay between multiple signalling pathways including 5' adenosine monophosphate-activated protein kinase (AMPK) [18]. Acute AMPK activation stimulates glucose transport and glycolysis while inhibiting glycogen synthase activity. However, chronic AMPK activation has been reported to cause increased glycogen accumulation consistent with AMP levels being elevated 2.5-fold in NMR hearts (figure 2c;  $p < 0.01$ ) compared to C57/BL6 mouse hearts [19].


Previous work has identified myocardial fumarate and succinate as key mediators of intracellular ROS damage during reperfusion [20,21]. In NMRs, differences in fumarate and succinate were not significant, however, an overall decrease in succinate/fumarate ratio was significant, when compared to C57/BL6. This is indicative of reduced reliance on OXPHOS and oxygen for ATP provision, and in agreement with the elevated NMR cardiac levels of intracellular lactate (2.5-fold;  $p < 0.02$ ), indicative of higher glycolytic versus oxidative flux. There was also a twofold increase in levels of acetate ( $p < 0.01$ ) indicative of higher pyruvate turnover [22]. Despite the regular hypoxic conditions experienced by NMRs in their colony, lack of myocardial succinate elevation could be an important cardioprotective mechanism, as succinate has been identified as a metabolic mediator of hypoxia-related ROS formation and elimination at the cross roads of several metabolic pathways [23].

Furthermore, increased glutathione levels were also observed (figure 2c; 2.6-fold,  $p < 0.01$ ), which is consistent with the observations of Munro *et al.* [24,25], where ROS scavenging and antioxidant defences were found to be elevated in the NMR, conferring a protective role against oxidative stress and ROS damage [26].

Differences in amino acid metabolism were also observed. The levels of glutamate were significantly higher in NMRs (approx. twofold;  $p$ -value  $< 0.01$ ), reflecting an alteration in anaplerotic pathways feeding the TCA cycle, while glutamate is also critical in glutathione synthesis. Reduced myocardial tyrosine levels corresponded to reduced circulating tyrosine levels previously reported in NMRs [27]. Levels of isoleucine were significantly elevated ( $p < 0.01$ ), providing an important nutrient source. Isoleucine is also a potent signalling molecule. Branched-chain amino acid (BCAA) L-leucine is a highly effective activator of mTOR signalling, [28] a key pathway for regulation of protein synthesis in hypertrophy which is exhibited in NMR heart [29–31]. mTOR activation also triggers metabolic changes in muscle, liver and other tissues by altering insulin sensitivity [32]. Therefore, it is plausible that the elevation of local BCAA concentration observed can lead to chronic induction of cardiac mTOR activity, in turn, promoting cardiac hypertrophy observed under normal physiological conditions in NMRs.

## 4. Discussion

We applied a semi-targeted metabolomic analysis approach and identified unique metabolic signatures characterized by




**Figure 1.** Representative cardiac  $^1\text{H}$  magnetic resonance spectra (MRS) of (a) naked mole-rat (NMR) compared to (b) control C57/BL6 mouse heart. Inset shows a detail in the region of the glucose and glycogen peaks.

oxygen sparing, ROS damage reduction and readily available ATP store strategy in NMR cardiac tissue that are a likely source of cardioprotection during hypoxia and senescence. Mammalian myocardial glycogen is considered an atavistic remnant of our amphibian ancestry [33]. Myocardial glycogen is an important source of ATP under conditions of metabolic stress which would account for the ability of NMR hearts to increase their cardiac reserve upon sudden demand, and maintain low basal metabolic rates even during prolonged hypoxia or decreased nutrition. A large glycogen store is commonly observed in hypoxia-tolerant species. The role of glycogen in hypoxia resistance is most pronounced in amphibian hearts [33], which are particularly rich in this compound [34,35]. Glycogen accounts for 2% of the cell volume in the healthy mammalian adult and 30% in the fetal myocardium [33]. Even though the fetal heart has better hypoxia adaptation than the adult heart, the physiological role of glycogen remains controversial [33]. Some consider it to be the damaging lactate and proton source in the ischaemic or stressed heart, causing severe contractile dysfunction particularly upon reperfusion [33,36]. However, previous work showed that blood and tissue pH levels in NMRs are largely unchanged after acute hypoxia, suggesting the absence of a metabolic acidosis [10]. As glycogen stores are markedly increased in the myocardium of hibernating mammals [37,38] and linked to increased hypoxia survival [39], there are others attributing a cardioprotective effect to

glycogen in the ischaemic heart [40–42]. Glycogen-rich hearts have been characterized by enhanced ischaemia tolerance, decreased protein loss and cell damage upon reperfusion, [33,43] leading to a hypothesis that glycogen acts as an anchoring molecule for other macromolecular cell constituents including adenine nucleotides and proteins [33,44,45] thereby integrating metabolic pathways required for myocardial survival during hypoxia [33].

Our study reveals similarities in glycogen metabolism between the hearts of NMRs and diving seals which have the ultimate physiological adaptation to prolonged and varying degrees of hypoxia [46–51] as well as Himalayan Sherpas [52]. The first lines of myocardial hypoxia defence include suppression of ATP supply and demand pathways [33], thus reducing ROS generation, during which hypoxia-tolerant systems activate protective mechanisms including metabolic reliance on intracellular glycogen [33,47,53]. When exposed to hypoxia, augmented glycogen stores would enable the NMR heart to switch to carbohydrates as the main fuel for respiration thus improving the metabolic efficiency (ATP produced per mole of oxygen consumed) [33,54], responding to acute increases in workload and stress by catabolizing their large intracellular glycogen store. This energetic advantage of glucose oxidation in the heart is well documented [33,55]. MRS and positron emission tomography ( $^{18}\text{F}$ FDG PET) studies in Sherpa hearts have suggested that enhanced glucose uptake, deposition and



**Figure 2.** Cardiac metabolomic profile of naked mole-rats. (a) C57/BL6 mouse and naked mole-rat. (b) Principal component analysis (PCA) of NMR versus C57/BL6 cardiac metabolomic profile showing a good separation between the two groups. (c)  $^1\text{H}$  nuclear magnetic resonance spectroscopy metabolomic profile presented as fold change NMR versus C57/BL6: redox and energetics (blue bars), TCA cycle, glycolysis and lipid metabolism intermediates (red bars) and amino acid metabolism (green bars). (d) Volcano plot showing  $\ln(\text{NMR}/\text{C57/BL6})$  versus  $-\log_{10}(P)$ . (e) Myocardial concentration of glycogen, glucose-1-phosphate (G-1-P) and  $\alpha$ -glucose in C57/BL6, Wistar rat and NMR.  $n = 5/\text{group}$ . \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  versus C57/BL6.

utilization in hypoxia are also advantageous for aerobic metabolism when  $\text{O}_2$  is present but at a premium [52]. Furthermore, the burst in glycogen oxidation is usually quickly followed by enhanced glycogen re-synthesis [54,56], and when an alternative metabolic substrate such as lactate is available for oxidation, glycogen stores are completely preserved [33,57]. Although both fatty acids and glucose are believed to be cycled through tissue-storage pools in the myocardium, the effect of such cycling has different consequences in terms of ATP provision. The incorporation of glucose into glycogen and the incorporation of fatty acids into triglycerides both consume ATP. However, in the breakdown of glycogen the energy expended in the synthesis is almost fully recovered with the efficiency of temporary storage processes close to 100% [58].

The cardioprotective effects of glycogen in NMRs could extend beyond its role as an endogenous metabolic substrate. It is plausible that cellular proteins are protected from glycosylation and in turn glucotoxicity by intracellular free glucose shunting into glycogen, comparable to shunting of fatty acid metabolites into triglycerides to protect the heart from lipotoxicity [33,59]. Furthermore, the rapid glycogen replenishment was shown to restore  $\text{Ca}^{2+}$  sensitivity and maximum  $\text{Ca}^{2+}$ -activated force in glycogen-depleted skeletal muscle [60]. Glycogen and glycogen-metabolizing enzymes co-locate with sarcoplasmic reticulum [61], thus playing a major role in the complex metabolic signalling systems of calcium homeostasis and cell survival [33,46].

## 5. Study limitations

In the current study, we have investigated myocardial metabolomic differences in C57/BL6 mice and NMRs. Significant errors are known to be associated when inferring environmental adaptation from a given observation in two species where there are potential phenotypic differences between the two species at baseline, for example, due to evolution [62]. Our study was performed under basal  $\text{O}_2$  conditions and therefore not representative of the environmental adaptations to hypoxia *per se*. Our study, therefore, likely reflects phenotypic alterations in the two species under normoxic conditions, some of which are likely relevant to their ability to survive extreme environments such as hypoxia. However, further work will be required to elucidate the complex interplay between environmental hypoxia and evolution in these species.

A further limitation of the current study is the use of MRS for quantification of glycogen.  $^1\text{H}$  magnetic resonance gives a measure of the concentration of glucose monomers that are present in the observed peak normalized to the reference trimethylsilyl propanoic acid (TSP) peak. Despite being a large macromolecule, with possible differences in the mobility of glycosyl units, glycogen has been reported to be fully visible by MRS [63]. However, we used a modified dual-phase Folch extraction method used for separating aqueous and lipid metabolites which has not been optimized for the extraction of glycogen *per se*. Thus, a direct comparison of the glycogen

tissue values obtained in this study to the values published by traditional enzymatic techniques may be at variance. Notwithstanding, the application of the same methods to all three species allows an assessment of fold changes between groups and therefore our assessment of between species variation remains valid.

**Ethics.** This investigation conforms to UK Home Office Guidance on the Operation of the Animals (Scientific Procedures) Act, 1986 incorporating European Directive 2010/63/EU (project licence PF75E5F7F and P71E91C8E for rat and mouse work, respectively) and approved by the local Committee for Animal Care and Ethical Review. Naked mole-rats were maintained in the Biological Services Unit at the Queen Mary University of London. Because tissue sample collection was post-euthanasia, additional local ethical approval for NMR work was not required for this study.

**Data accessibility.** Data are provided as an electronic supplementary material Excel file.

**Authors' contributions.** D.A. designed research; D.A., C.G.F., T.R.E. performed research, analysed and interpreted data; wrote the

manuscript. All authors agree to be held accountable for the content therein and approve the final version of the manuscript. C.G.F., T.R.E., D.A. all made substantial contributions to conception and design, acquisition of data, analysis and interpretation of data; drafting the article and revising it critically for important intellectual content.

**Competing interests.** We declare we have no competing interests.

**Funding.** D.A.: QMUL, School of Biological and Chemical Sciences. T.R.E.: NIHR Biomedical Research Centre at Guy's and St Thomas' NHS Foundation Trust and KCL; the Centre of Excellence in Medical Engineering funded by the Wellcome Trust and Engineering and EPSRC (grant no. WT 203148/Z/16/Z); the BHF Centre of Research Excellence (grant no. RE/18/2/34213) and the KCL Comprehensive Cancer Imaging Centre funded by the Cancer Research UK and EPSRC in association with the MRC and the Department of Health (DoH). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or DoH.

**Acknowledgements.** Thanks to Dr H. Toms, QMUL High Field Nuclear Magnetic Resonance Laboratory; Thanks to Lorna Faulkes for the NMR and Prof. Michael Shattock for C57/BL6 photo in figure 2.

## References

- Liang S, Mele J, Wu Y, Buffenstein R, Hornsby PJ. 2010 Resistance to experimental tumorigenesis in cells of a long-lived mammal, the naked mole-rat (*Heterocephalus glaber*). *Aging Cell* **9**, 626–635. (doi:10.1111/j.1474-9726.2010.00588.x)
- Park TJ *et al.* 2017 Fructose-driven glycolysis supports anoxia resistance in the naked mole-rat. *Science* **356**, 307–311. (doi:10.1126/science.aab3896)
- Smith ES, Omerbasic D, Lechner SG, Anirudhan G, Lapatsina L, Lewin GR. 2011 The molecular basis of acid insensitivity in the African naked mole-rat. *Science* **334**, 1557–1560. (doi:10.1126/science.1213760)
- Holtze S *et al.* 2017 The microenvironment of naked mole-rat burrows in East Africa. *Afr. J. Ecol.* **56**, 279–289. (doi:10.1111/aje.12448)
- Faulkes CG, Bennett NC. 2013 Plasticity and constraints on social evolution in African mole-rats: ultimate and proximate factors. *Phil. Trans. R. Soc. Lond. B Biol. Sci.* **368**, 20120347. (doi:10.1098/rstb.2012.0347)
- Brett RA. 1991 The population structure of naked mole-rat colonies. In *The biology of the naked mole-rat* (eds PW Sherman, M Jarvis, RD Alexander), pp. 97–136. Princeton, NJ: Princeton University Press.
- Larson J, Park TJ. 2009 Extreme hypoxia tolerance of naked mole-rat brain. *Neuroreport* **20**, 1634–1637. (doi:10.1097/WNR.0b013e32833370cf)
- Larson J, Drew KL, Folkow LP, Milton SL, Park TJ. 2014 No oxygen? No problem! Intrinsic brain tolerance to hypoxia in vertebrates. *J. Exp. Biol.* **217**(Pt 7), 1024–1039. (doi:10.1242/jeb.085381)
- Peterson BL, Larson J, Buffenstein R, Park TJ, Fall CP. 2012 Blunted neuronal calcium response to hypoxia in naked mole-rat hippocampus. *PLoS ONE* **7**, e31568. (doi:10.1371/journal.pone.0031568)
- Pamenter ME, Dzal YA, Thompson WA, Milsom WK. 2019 Do naked mole rats accumulate a metabolic acidosis or an oxygen debt in severe hypoxia? *J. Exp. Biol.* **222**(Pt 3), jeb191197. (doi:10.1242/jeb.191197)
- Goldman BD, Goldman SL, Lanz T, Magaurin A, Maurice A. 1999 Factors influencing metabolic rate in naked mole-rats (*Heterocephalus glaber*). *Physiol. Behav.* **66**, 447–459. (doi:10.1016/S0031-9384(98)00306-0)
- Grimes KM, Barefield DY, Kumar M, McNamara JW, Weintraub ST, de Tombe PP, Sadayappan S, Buffenstein R. 2017 The naked mole-rat exhibits an unusual cardiac myofilament protein profile providing new insights into heart function of this naturally subterranean rodent. *Pflugers Arch.* **469**, 1603–1613. (doi:10.1007/s00424-017-2046-3)
- Grimes KM, Lindsey ML, Gelfond JA, Buffenstein R. 2012 Getting to the heart of the matter: age-related changes in diastolic heart function in the longest-lived rodent, the naked mole rat. *J. Gerontol. A Biol. Sci. Med. Sci.* **67**, 384–394. (doi:10.1093/geron/67/3/384)
- Grimes KM, Reddy AK, Lindsey ML, Buffenstein R. 2014 And the beat goes on: maintained cardiovascular function during aging in the longest-lived rodent, the naked mole-rat. *Am. J. Physiol. Heart Circ. Physiol.* **307**, H284–H291. (doi:10.1152/ajpheart.00305.2014)
- Grimes KM, Voorhees A, Chiao YA, Han HC, Lindsey ML, Buffenstein R. 2014 Cardiac function of the naked mole-rat: ecophysiological responses to working underground. *Am. J. Physiol. Heart Circ. Physiol.* **306**, H730–H737. (doi:10.1152/ajpheart.00831.2013)
- Faulkes CG, Abbott DH, Jarvis JU. 1990 Social suppression of ovarian cyclicity in captive and wild colonies of naked mole-rats, *Heterocephalus glaber*. *J. Reprod. Fertil.* **88**, 559–568. (doi:10.1530/jrf.0.0880559)
- Chung YL, Leach MO, Eykyn TR. 2017 Magnetic resonance spectroscopy to study glycolytic metabolism during autophagy. *Methods Enzymol.* **588**, 133–153. (doi:10.1016/bs.mie.2016.09.078)
- Chandramouli C, Varma U, Stevens EM, Xiao RP, Stapleton DJ, Mellor KM, Delbridge LM. 2015 Myocardial glycogen dynamics: new perspectives on disease mechanisms. *Clin. Exp. Pharmacol. Physiol.* **42**, 415–425. (doi:10.1111/1440-1681.12370)
- Hunter RW, Treebak JT, Wojtaszewski JF, Sakamoto K. 2011 Molecular mechanism by which AMP-activated protein kinase activation promotes glycogen accumulation in muscle. *Diabetes* **60**, 766–774. (doi:10.2337/db10-1148)
- Ashrafian H *et al.* 2012 Fumarate is cardioprotective via activation of the Nrf2 antioxidant pathway. *Cell Metab.* **15**, 361–371. (doi:10.1016/j.cmet.2012.01.017)
- Chouchani ET *et al.* 2014 Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* **515**, 431–435. (doi:10.1038/nature13909)
- Bose S, Ramesh V, Locasale JW. 2019 Acetate metabolism in physiology, cancer, and beyond. *Trends Cell Biol.* **29**, 695–703. (doi:10.1016/j.tcb.2019.05.005)
- Tretter L, Patocs A, Chinopoulos C. 2016 Succinate, an intermediate in metabolism, signal transduction, ROS, hypoxia, and tumorigenesis. *Biochim. Biophys. Acta* **1857**, 1086–1101. (doi:10.1016/j.bbabi.2016.03.012)
- Munro D, Baldy C, Pamenter ME, Treberg JR. 2019 The exceptional longevity of the naked mole-rat may be explained by mitochondrial antioxidant defenses. *Aging Cell* **18**, e12916. (doi:10.1111/acer.12916)
- Munro D, Pamenter ME. 2019 Comparative studies of mitochondrial reactive oxygen species in animal longevity: technical pitfalls and possibilities. *Aging Cell* **18**, e13009. (doi:10.1111/acer.13009)
- O'Donovan DJ, Fernandes CJ. 2000 Mitochondrial glutathione and oxidative stress: implications for pulmonary oxygen toxicity in premature infants. *Mol. Genet. Metab.* **71**, 352–358. (doi:10.1006/mgme.2000.3063)
- Lewis KN, Rubinstein ND, Buffenstein R. 2018 Correction to: a window into extreme longevity; the

- circulating metabolomic signature of the naked mole-rat, a mammal that shows negligible senescence. *Geroscience* **40**, 357–358. (doi:10.1007/s11357-018-0023-1)
28. Proud CG. 2002 Control of the translational machinery in mammalian cells. *Eur. J. Biochem.* **269**, 5337. (doi:10.1046/j.1432-1033.2002.03289.x)
  29. Katta A, Kundla S, Kakarla SK, Wu M, Fannin J, Paturi S, Liu H, Addagarla HS, Blough ER. 2010 Impaired overload-induced hypertrophy is associated with diminished mTOR signaling in insulin-resistant skeletal muscle of the obese Zucker rat. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **299**, R1666–R1675. (doi:10.1152/ajpregu.00229.2010)
  30. Xu G, Kwon G, Cruz WS, Marshall CA, McDaniel ML. 2001 Metabolic regulation by leucine of translation initiation through the mTOR-signaling pathway by pancreatic  $\beta$ -cells. *Diabetes* **50**, 353–360. (doi:10.2337/diabetes.50.2.353)
  31. Zhang D *et al.* 2010 MTORC1 regulates cardiac function and myocyte survival through 4E-BP1 inhibition in mice. *J. Clin. Invest.* **120**, 2805–2816. (doi:10.1172/JCI43008)
  32. Huang Y, Zhou M, Sun H, Wang Y. 2011 Branched-chain amino acid metabolism in heart disease: an epiphenomenon or a real culprit? *Cardiovasc. Res.* **90**, 220–223. (doi:10.1093/cvr/cvr070)
  33. Taegtmeyer H. 2004 Glycogen in the heart—an expanded view. *J. Mol. Cell. Cardiol.* **37**, 7–10. (doi:10.1016/j.yjmcc.2004.05.001)
  34. Reeves RB. 1963 Control of glycogen utilization and glucose uptake in the anaerobic turtle heart. *Am. J. Physiol.* **205**, 23–29. (doi:10.1152/ajplegacy.1963.205.1.23)
  35. Wasser JS, Meinertz EA, Chang SY, Lawler RG, Jackson DC. 1992 Metabolic and cardiodynamic responses of isolated turtle hearts to ischemia and reperfusion. *Am. J. Physiol.* **262**(3 Pt 2), R437–R443. (doi:10.1152/ajpregu.1992.262.3.r437)
  36. Neely JR, Grotyohann LW. 1984 Role of glycolytic products in damage to ischemic myocardium. Dissociation of adenosine triphosphate levels and recovery of function of reperfused ischemic hearts. *Circ. Res.* **55**, 816–824. (doi:10.1161/01.RES.55.6.816)
  37. Depre C, Vanoverschelde JL, Melin JA, Borgers M, Bol A, Ausma J, Dion R, Wijns W. 1995 Structural and metabolic correlates of the reversibility of chronic left ventricular ischemic dysfunction in humans. *Am. J. Physiol.* **268**(3 Pt 2), H1265–H1275. (doi:10.1152/ajpheart.1995.268.3.h1265)
  38. Kim SJ, Peppas A, Hong SK, Yang G, Huang Y, Diaz G, Sadoshima J, Vatner DE, Vatner SF. 2003 Persistent stunning induces myocardial hibernation and protection: flow/function and metabolic mechanisms. *Circ. Res.* **92**, 1233–1239. (doi:10.1161/01.RES.0000076892.18394.B6)
  39. Dawes GS, Mott JC, Shelley HJ. 1959 The importance of cardiac glycogen for the maintenance of life in foetal lambs and newborn animals during anoxia. *J. Physiol.* **146**, 516–538. (doi:10.1113/jphysiol.1959.sp006208)
  40. Doenst T, Guthrie PH, Chemnitz JM, Zech R, Taegtmeyer H. 1996 Fasting, lactate, and insulin improve ischemia tolerance in rat heart: a comparison with ischemic preconditioning. *Am. J. Physiol.* **270**(5 Pt 2), H1607–H1615. (doi:10.1152/ajpheart.1996.270.5.h1607)
  41. Lagerstrom CF, Walker WE, Taegtmeyer H. 1988 Failure of glycogen depletion to improve left ventricular function of the rabbit heart after hypothermic ischemic arrest. *Circ. Res.* **63**, 81–86. (doi:10.1161/01.RES.63.1.81)
  42. Scheuer J, Stezoski SW. 1970 Protective role of increased myocardial glycogen stores in cardiac anoxia in the rat. *Circ. Res.* **27**, 835–849. (doi:10.1161/01.RES.27.5.835)
  43. Schneider CA, Taegtmeyer H. 1991 Fasting *in vivo* delays myocardial cell damage after brief periods of ischemia in the isolated working rat heart. *Circ. Res.* **68**, 1045–1050. (doi:10.1161/01.RES.68.4.1045)
  44. Depre C, Taegtmeyer H. 2000 Metabolic aspects of programmed cell survival and cell death in the heart. *Cardiovasc. Res.* **45**, 538–548. (doi:10.1016/S0008-6363(99)00266-7)
  45. Taegtmeyer H. 1994 Energy metabolism of the heart: from basic concepts to clinical applications. *Curr. Probl. Cardiol.* **19**, 59–113. (doi:10.1016/0146-2806(94)90008-6)
  46. Henden T, Aasum E, Folkow L, Mjos OD, Lathrop DA, Larsen TS. 2004 Endogenous glycogen prevents  $\text{Ca}^{2+}$  overload and hypercontracture in harp seal myocardial cells during simulated ischemia. *J. Mol. Cell. Cardiol.* **37**, 43–50. (doi:10.1016/j.yjmcc.2004.03.013)
  47. Hochachka PW, Lutz PL. 2001 Mechanism, origin, and evolution of anoxia tolerance in animals. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **130**, 435–459. (doi:10.1016/S1096-4959(01)00408-0)
  48. Hochachka PW, Owen TG, Allen JF, Whittow GC. 1975 Multiple end products of anaerobiosis in diving vertebrates. *Comp. Biochem. Physiol. B* **50**, 17–22. (doi:10.1016/0305-0491(75)90292-8)
  49. Hochachka PW, Storey KB. 1975 Metabolic consequences of diving in animals and man. *Science* **187**, 613–621. (doi:10.1126/science.163485)
  50. Kjekshus JK, Blix AS, Elsner R, Hol R, Amundsen E. 1982 Myocardial blood flow and metabolism in the diving seal. *Am. J. Physiol.* **242**, R97–104. (doi:10.1152/ajpregu.1982.242.1.r97)
  51. Williams TM *et al.* 2015 Exercise at depth alters bradycardia and incidence of cardiac anomalies in deep-diving marine mammals. *Nat. Commun.* **6**, 6055. (doi:10.1038/ncomms7055)
  52. Hochachka PW, Clark CM, Holden JE, Stanley C, Ugurbil K, Menon RS. 1996  $^{31}\text{P}$  magnetic resonance spectroscopy of the Sherpa heart: a phosphocreatine/adenosine triphosphate signature of metabolic defense against hypobaric hypoxia. *Proc. Natl Acad. Sci. USA* **93**, 1215–1220. (doi:10.1073/pnas.93.3.1215)
  53. Webster KA, Discher DJ, Bishopric NH. 1993 Induction and nuclear accumulation of fos and jun proto-oncogenes in hypoxic cardiac myocytes. *J. Biol. Chem.* **268**, 16 852–16 858.
  54. Goodwin GW, Taylor CS, Taegtmeyer H. 1998 Regulation of energy metabolism of the heart during acute increase in heart work. *J. Biol. Chem.* **273**, 29 530–29 539. (doi:10.1074/jbc.273.45.29530)
  55. Korvald C, Elvenes OP, Myrnes T. 2000 Myocardial substrate metabolism influences left ventricular energetics *in vivo*. *Am. J. Physiol. Heart Circ. Physiol.* **278**, H1345–H1351. (doi:10.1152/ajpheart.2000.278.4.H1345)
  56. Russell 3rd RR, Nguyen VT, Mrus JM, Taegtmeyer H. 1992 Fasting and lactate unmask insulin responsiveness in the isolated working rat heart. *Am. J. Physiol.* **263**(3 Pt 1), E556–E561. (doi:10.1152/ajpendo.1992.263.3.e556)
  57. Goodwin GW, Taegtmeyer H. 2000 Improved energy homeostasis of the heart in the metabolic state of exercise. *Am. J. Physiol. Heart Circ. Physiol.* **279**, H1490–H1501. (doi:10.1152/ajpheart.2000.279.4.H1490)
  58. Holden JE, Stone CK, Clark CM, Brown WD, Nickles RJ, Stanley C, Hochachka PW. 1995 Enhanced cardiac metabolism of plasma glucose in high-altitude natives: adaptation against chronic hypoxia. *J. Appl. Physiol.* (1985) **79**, 222–228. (doi:10.1152/jappl.1995.79.1.222)
  59. Listenberger LL, Han X, Lewis SE, Cases S, Farese Jr RV, Ory DS, Schaffer JE. 2003 Triglyceride accumulation protects against fatty acid-induced lipotoxicity. *Proc. Natl Acad. Sci. USA* **100**, 3077–3082. (doi:10.1073/pnas.0630588100)
  60. Chin ER, Allen DG. 1997 Effects of reduced muscle glycogen concentration on force,  $\text{Ca}^{2+}$  release and contractile protein function in intact mouse skeletal muscle. *J. Physiol.* **498**(Pt 1), 17–29. (doi:10.1113/jphysiol.1997.sp021838)
  61. Entman ML, Goldstein MA, Schwartz A. 1976 The cardiac sarcoplasmic reticulum—glycogenolytic complex, an internal  $\beta$  adrenergic receptor. *Life Sci.* **19**, 1623–1630. (doi:10.1016/0024-3205(76)90066-7)
  62. Garland T J, Adolph SC. 1994 Why not to do two-species comparative studies: limitations on inferring adaptation. *Physiol. Zool.* **67**, 797–828. (doi:10.1086/physzool.67.4.30163866)
  63. Zang LH, Rothman DL, Schulman RG. 1990  $^1\text{H}$  NMR visibility of mammalian glycogen in solution. *Proc. Natl Acad. Sci. USA* **87**, 1678–1680. (doi:10.1073/pnas.87.5.1678)