

**Glucagon-like peptide 2 concentrations vary in Zambian children during diarrhoea, in malnutrition and seasonally.**

Ellen **Besa**<sup>1</sup>, Kanta **Chandwe**<sup>1,2</sup>, Rosemary **Banda**<sup>1</sup>, Likando **Munalula**<sup>1</sup>, Lydia **Kalomo**<sup>1</sup>, Beatrice **Amadi**<sup>1,2</sup>, Paul **Kelly**<sup>1,3,4</sup>

1 Tropical Gastroenterology and Nutrition group, University of Zambia School of Medicine, Lusaka, Zambia; 2 Children's Hospital, University Teaching Hospitals, Lusaka, Zambia;

3 Blizzard Institute, Queen Mary University of London, London, UK; 4 London School of Hygiene and Tropical Medicine, London, UK.

**Corresponding author:** Paul Kelly, Tropical Gastroenterology and Nutrition group, University of Zambia School of Medicine, Lusaka, Zambia, Tel: + 260 966 751875, email: m.p.kelly@qmul.ac.uk

**Funding:** Funding was obtained from the Bill & Melinda Gates Foundation (OPP1066118). The funding agency played no part on the analysis of the data or the decision to publish or the writing of the manuscript.

**Conflicts of interest** None

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. <http://creativecommons.org/licenses/by/4.0>

## **Abstract**

*Objectives:* Glucagon Like Peptide 2 (GLP-2) is a 33 amino acid peptide hormone released from enteroendocrine L-cells following nutrient ingestion. It has been shown to exert trophic effects on the gut. We set out to measure GLP-2 concentrations in blood in children with diarrhoea and malnutrition.

*Methods:* GLP-2 levels were measured in blood samples collected from five different groups of children (n=324) at different time points: (1) those with acute diarrhoea, during illness and 3 weeks after recovery; (2) persistent diarrhoea and severe acute malnutrition (SAM); (3) controls contemporaneous for diarrhoea; (4) stunted children from the community; (5) controls contemporaneous for the stunted children. Stool biomarkers and pathogen analysis was carried out on the children with stunting.

*Results:* GLP-2 concentrations were higher during acute diarrhoea (median 3.1 ng/ml, interquartile range, IQR 2.1, 4.4) than on recovery (median 1.8, IQR 1.4, 3.1;  $P=0.001$ ), but were not elevated in children with persistent diarrhoea and SAM. In stunted children, there was a progressive decline in GLP-2 levels from 3.2 ng/ml (1.9, 4.9) to 1.0 (0.0, 2.0;  $P<0.001$ ) as the children became more stunted. Measures of seasonality (rainfall, temperature, food price index and Shiga toxin-producing *Escherichia coli*) were found to be significantly associated with GLP-2 concentrations in multivariable analysis. We also found a correlation between stool inflammatory biomarkers and GLP-2.

*Conclusions:* In diarrhoea, GLP-2 levels increased in acute but not persistent diarrhoea. Malnutrition was associated with reduced concentrations. GLP-2 displayed seasonal variation consistent with variations in nutrient availability.

**Keywords:** diarrhea, GLP-2

### **What is known**

- Diarrhoeal disease in children is still a major cause of morbidity and mortality globally
- Highest rates of diarrhoea-related mortality occur among malnourished children
- Glucagon-like peptide 2 (GLP-2) is a hormone which can be used to improve intestinal adaptation and repair.

### **What is new**

- GLP-2 levels in blood rose during acute diarrhoea but not persistent diarrhoea, in which levels were similar to healthy controls
- GLP-2 levels fell progressively in children with stunting
- GLP-2 levels exhibited a seasonal pattern, with evidence of an influence of nutrient availability

ACCEPTED

## Introduction

Diarrhoeal disease and malnutrition continue to contribute significantly to global childhood morbidity and mortality. Diarrhoeal disease is the second leading cause of death in children under the age of 5, killing around 525 000 children yearly<sup>1</sup>. In the period 2000-2015, it accounted for 8.6% of the total deaths in children aged 1-59 months<sup>2</sup>. Treatment with oral rehydration solutions, intravenous fluids and zinc has led to a reduction in mortality<sup>3</sup>, but sub-Saharan Africa and Southern Asia continue to have the highest numbers of under-5 deaths despite efforts to try and reduce the disease burden<sup>4</sup>. The majority of diarrhoea cases are of acute onset, but 3 - 20% of these cases progress to persistent diarrhoea (PD) which contributes disproportionately to morbidity and mortality<sup>5,6,7</sup>. Diarrhoeal illness is also known to affect ponderal and linear growth, especially in cases of recurrent illness<sup>8</sup>. The problem of diarrhoea is further compounded by undernutrition, which is the largest single underlying cause of death among children under five<sup>9</sup>. There is a vicious malnutrition-infection cycle as malnutrition lowers immune function, increasing susceptibility to infections which then contribute to undernutrition by reducing energy intake, as well as nutrient loss and malabsorption<sup>10</sup>. Undernutrition may also be exacerbated by the poor living conditions which make one more prone to infections and disease. Children living under unsanitary conditions tend to have a generalized disturbance of small intestinal structure and function described as enteropathy. It has been implicated in growth failure through different mechanisms such as heightened permeability, gut inflammation, bacterial translocation and nutrient malabsorption<sup>8</sup>. Severe enteropathy has been observed in children with severe acute malnutrition (SAM) and PD<sup>11,12</sup> contributing to reduced surface area available for nutrient absorption and impaired barrier function<sup>13</sup>.

Glucagon Like Peptide-2 (GLP-2) is a 33-amino acid peptide growth factor that is secreted in the lower gastrointestinal tract following nutrient ingestion, particularly carbohydrates and lipids<sup>14,15</sup>. It was first described as 'enteroglucagon'<sup>16</sup> in a patient with a renal tumour which induced intestinal hypertrophy. GLP-2 is generated via post-translational processing of glucagon and is co-secreted with GLP-1, oxyntomodulin, and glicentin in a 1:1:1 ratio<sup>17,18</sup>. Once GLP-2 has been released, its effects are mediated through the specific binding of the hormone to its receptor, GLP-2R which localizes to enteric neurons and endocrine cells<sup>19,20</sup>. GLP-2 secretion is regulated by nutritional, hormonal and neural factors and it has been suggested that many of its actions are exerted indirectly through secondary mediators such as growth factors from enteroendocrine, neural and lamina propria cells. Some of its actions include upregulating motility, nutrient absorption and cell proliferation while reducing epithelial permeability and apoptosis in the gut<sup>18,21,22</sup>. Post prandial GLP-2 secretion is regulated in a complex manner consisting of both direct (via nutrients) and indirect (via endocrine and/or neural) pathways<sup>17</sup>. Nutrient stimulated GLP-2 release

has been found to result in a 2 – 5fold increase in plasma levels and as such can easily be detected in fed samples<sup>23</sup>.

We have recently shown that low circulating concentrations of GLP-2 were associated with high levels of bacterial translocation, suggesting that impaired secretion may contribute to mucosal dysfunction during diarrhoea infection<sup>24</sup>. The study set out to investigate what changes, if any occur in the levels of GLP-2 in children with diarrhoea and malnutrition, and whether GLP-2 release has any link to seasonality, as nutrient availability and transmission of enteropathogens are seasonal in Zambia.

## **Methods**

### *Study Participants*

Five different groups of children were studied over the period May 2014 to April 2019. Approval to conduct these studies was obtained from the University of Zambia Biomedical Research Ethics Committee (UNZABREC).

### *Group 1 - Acute Diarrhoea*

28 children with acute diarrhoea, defined as 3 or more loose stools in a 24-hour period (median duration 3 days, interquartile range (IQR) 2-4) were recruited when they presented to St. Lawrence clinic (Misisi, Lusaka). The children were assessed by Integrated Management of Childhood Illness (IMCI) trained nurses and managed accordingly<sup>25</sup>. Informed written consent was obtained from the caregiver, and a 2ml blood sample collected at the time of recruitment. The children were then followed up and a convalescent blood sample obtained three weeks after the diarrhoea episode ended. UNZABREC approval (007-11-15, dated 12<sup>th</sup> January 2016).

### *Group 2 – Persistent Diarrhoea*

These children were undergoing treatment in the malnutrition ward of the University Teaching Hospital, Lusaka, Zambia. They had PD (diarrhoea for more than 14 days, including skip days, median duration 21days, IQR 14-28) and SAM (Weight for length/height (WLZ/WHZ) < -3SD and/or mid upper arm circumference; MUAC < 115mm and/or bilateral pitting oedema of nutritional origin). No pathogens were

detected on first line investigations (stool microscopy and culture). All the admitted children with SAM were managed according to World Health Organisation (WHO) guidelines on inpatient management of SAM<sup>26</sup>. A total of 21 children were recruited and informed written consent obtained from the caregiver. Blood samples were collected from the children. UNZABREC approval (006-01-13, dated 15<sup>th</sup> April 2013).

#### *Group 3 – Controls for Group 1 & 2*

75 stunted but clinically healthy children were recruited from Misisi compound as contemporaneous controls for Groups 1 & 2. They were children who were screened for malnutrition in a previously described nutrition outreach programme<sup>27</sup>. Children from the community were screened for recent episodes of diarrhoea (within one month), NSAID use (within one month), antibiotic use (within one month) and excluded and treated if needed. These children were selected on the basis that they were not acutely malnourished, but 56% had stunting (against a background of 40% stunting across the whole country)<sup>28</sup>

#### *Group 4 – Stunted children*

5660 children aged 0 to 18 months of age, resident in Misisi compound were screened for wasting (MUAC <125mm and/or WLZ/WHZ z-score of < -2 SD) and stunting (length/height-for-age (LAZ/HAZ) z-score of < -2 SD) in order to identify those with malnutrition or growth faltering. 297 of those identified, were then recruited into a study of biomarkers of environmental enteropathy in children (BEECH), provided with nutritional rehabilitation and followed up to the age of 24 months or for a minimum of 12 months. Blood samples were collected at recruitment and at 3 months. For those children who did not respond to 3 months of nutritional rehabilitation, fed and fasted samples were collected on consecutive days. Non-response was defined as lack of improvement in anthropometric parameters (WLZ/LAZ/WAZ > -2SD), provided the child had not been ill in this period. Children were fasted for at least 4 hours before a sample was collected, as they were being prepared for endoscopy. Stool samples were also collected from the children as they came in for their first visit. 181 baseline blood samples were tested for GLP-2. UNZABREC approval (006-02-16, dated 3<sup>rd</sup> January 2018)

#### *Group 5 – Controls for Group 4*

Contemporaneous controls (n=46) were recruited for the stunted children to control for the time lapse between the studies, as well as temporal changes, seasonal changes and an improvement in stunting from 40% to 35%<sup>28</sup>. These children were well nourished (WLZ > -1) and resident in Misisi compound. 19 baseline blood samples were tested for GLP-2.

#### *Assessment of Nutritional status*

A paediatrician and trained study nurses carried out nutritional status assessment of all the participants based on WHO child growth standards<sup>29</sup>. This was done using anthropometric measurements of mid-upper arm circumference (MUAC), weight and length/height. The measurements were done using a Mother-Child scale (SECA 874, Hamburg Germany) for weight, Infantometers (SECA 416, Hamburg Germany) for length, UNICEF height boards for height and MUAC tapes. Raw data were entered into the WHO Anthro software v.3.2.2<sup>30</sup> which then calculated the anthropometry scores.

#### *Measurement of GLP-2, stool markers and stool pathogens*

Plasma samples were tested for GLP-2 by Enzyme Linked Immunosorbent Assay (ELISA) (Millipore Corporation, St Charles Missouri, USA). The assay is a sandwich ELISA that only measures total GLP-2.

Fecal Myeloperoxidase and Calprotectin were measured as markers of inflammation in children with stunting. Myeloperoxidase was measured using the EDI™ Quantitative Fecal/Urine Myeloperoxidase ELISA Kit (Epitope diagnostics, Inc. San Diego, USA) and Calprotectin using the IDK® Calprotectin ELISA (Immundiagnostik AG, Bensheim, Germany). Both assays utilize the two-site sandwich technique with two selected monoclonal antibodies binding to the protein of interest.

Stool pathogen analysis was done using the qualitative, multiplex polymerase chain reaction (PCR) based Luminex x-TAG® gastrointestinal pathogen panel (Luminex Corporation, Austin, Texas, USA). This assay is able to simultaneously detect 15 enteric pathogens with a sensitivity of 94.3%, for 12 of the 15 pathogens, and a specificity of 98.5% across all 15 pathogens<sup>31</sup>. Sample processing and assay conditions are described by Chisenga et al<sup>32</sup>. All the assays were run according to the manufacturer's instructions and more detail on the assays is provided in Supplemental Digital Content 1 (<http://links.lww.com/MPG/B783>).

### *Data regarding environmental exposures*

Rainfall and temperature data for the period 2016 – 2018 were obtained from the Lusaka City Airport, Zambia Meteorological Office. Food price index (FPI) was obtained from the Jesuit Centre for Theological Reflection monthly survey<sup>33</sup>. This survey details retail costs of basic food items of a family of five. It is carried out in different towns and it averages the cost of food based on the prices gathered from different places within the town of interest. For the purposes of the paper, the prices indicated were averaged from two markets in the vicinity of our study catchment area namely Chawama market and City market. This was to allow us to get more representative costs as averaging it across the costs of the entire town would not give a true picture.

### *Data analysis and statistical considerations*

GLP-2 concentrations in serum were determined to be non-normally distributed using the Shapiro-Wilk test, so data are presented as median and interquartile range (IQR). The Kruskal-Wallis test was used for hypothesis testing. The Wilcoxon matched-pairs rank sum test was used to test paired samples, and the Friedman test for multiple comparisons. Spearman's correlation was used to test for correlation between GLP-2 and stool inflammatory markers. A  $p$ -value less than 0.05 was considered significant. Univariable and multivariable logistic regression analysis was carried out to determine if there was any association between GLP-2 concentrations and different variables and pathogens. Pathogen results with numbers less than 10 were not included in the final model. Missing data were treated as missing and no imputation was made. Data were analysed using GraphPad Prism 5, and STATA v.15.

## **Results**

### *Participant characteristics*

324 participants were studied over the course of 5 years. Groups 1 and 2 were comparable in sex and age but not in anthropometry measures (Table 1). The majority of the children from Groups 1, 2 and 3 were stunted but those in Group 2 were also wasted. The control group (Group 3) had not had diarrhoea in the month prior to recruitment, were slightly older with more female participants than the other two groups. The stunted children (Group 4) were also older than their contemporaneous controls (Group 5).



### *GLP-2 levels rise in acute diarrhoea but not persistent diarrhoea*

GLP 2 levels in children with acute diarrhoea (Group 1), were elevated during illness and fell after convalescence ( $P=0.001$ ), in all except 5 children (Figure 1A). Circulating GLP-2 concentrations after convalescence (median 1.83ng/ml, interquartile range 1.38,3.12) were the same as in the controls (Group 3) from the same community (median 1.87, IQR 1.40,2.56), confirming that it was the samples taken during acute diarrhoea which were elevated (median 3.06ng/ml, IQR 2.09,4.39; Figure 1B). Children with PD and SAM (Group 2) had concentrations (median 1.93, IQR 1.28,4.07) which were similar to control values ( $P=0.33$ ) (Figure 1B).

### *GLP-2 in children with stunting and non-responders*

In Group 4, GLP-2 levels at recruitment (3.2, IQR 1.9,4.9) did not differ from the 3-month visit (3.5, IQR 2.39,4.83) or controls (3.1, IQR 2.39,4.16), but were significantly lower in the fasted non-responders (Figure 2A). GLP-2 in the fasted state was 1.01ng/ml (IQR 0.2,0.1) and in the fed state was 2.40ng/ml (IQR 2.10,3.44). Fed and fasted values were both lower than baseline GLP-2 value ( $p < 0.001$ ) suggesting that the fall in GLP-2 was not merely due to the fasted state (Figure 2B).

### *GLP-2 in controls*

In the control groups (groups 3 and 5), GLP-2 concentrations in plasma fell with age ( $\rho = -0.35$ ;  $P=0.001$ ). Using an estimate of  $\beta$  from linear regression the change in GLP-2 over time would be a reduction of 0.04 ng/ml per month of increasing age. It is thus very unlikely that the decline in GLP-2 observed in group 4 could be attributable to age.

### *Stool Inflammatory markers*

In the baseline stool samples, median myeloperoxidase stool concentrations were at 102.51ng/ml (IQR 40.37, 169.68) and calprotectin concentrations at 278.38ug/g(IQR 164.73,619.91). There was a negative correlation between GLP-2 and myeloperoxidase ( $\rho = -0.16$ ,  $p = 0.02$ ) and a positive correlation between GLP-2 and Calprotectin ( $\rho = 0.18$ ,  $p = 0.02$ )

### *GLP-2 shows seasonal pattern*

Seasonal variation has been shown in the prevalence of undernutrition<sup>34</sup> and incidence of diarrhoea<sup>35</sup>. Diarrhoeal pathogens have also been shown to vary seasonally, influenced by factors such as rainfall, temperature and humidity<sup>36</sup>. We set out to explore if there is any association between GLP-2 release and these factors that vary seasonally i.e. food availability, rainfall, temperature and pathogens. The FPI, which covers the cost of food items such as maize meal, beans, kapenta, fish, and vegetables, constituents of a typical Zambian meal, was used as a proxy of food availability.

We found an association between GLP-2 and FPI, temperature and rainfall. There was an increase in GLP-2 during the warmer months, and a decrease in GLP-2 when the cost of food was high, and during the rainy season. We also looked at whether there was any association between the different pathogens detected in stool and GLP-2 release as it has been documented that subclinical carriage of pathogens may result in enteropathy<sup>37</sup>. Intestinal damage from enteropathy may result in a deficiency of hormonal response to nutrients. An association was found between GLP-2 and Shiga toxin-producing *Escherichia coli*. Multivariate regression was carried out to confirm these associations (Table 2).

### **Discussion**

Taken together, our results suggest that GLP2 levels are elevated in acute diarrhoea but not in PD and SAM. In children with acute diarrhoea, GLP-2 levels rose and then returned to background levels after recovery from infection. This suggests that in cases of acute diarrhoea, the intestinal response to infection includes an increase in the amount of circulating GLP-2 which would enhance repair of the damaged mucosal epithelium, thereby restoring epithelial integrity. A similar response has also been observed in children with intestinal failure<sup>38</sup> suggestive of the role GLP-2 plays in enhancing the adaptive response. In that study, mean fasted GLP-2 in controls was 11.6ng/ml, and 19.9ng/ml in children with intestinal failure. All the children in our study had considerably lower circulating GLP-2 concentrations than children in Finland<sup>38</sup> but as different ELISA kits were used, the difference noted could be due to methodological as well as biological factors. In PD, the reverse of what is occurring in acute diarrhoea was observed. It is plausible that in the PD children, SAM plays a major role in the inability of the body to respond in a manner similar to that observed in the acute diarrhoea cases. An inability to produce GLP-2 may perpetuate the malnutrition-infection cycle by impairment of mucosal regeneration, and ultimately lead to death in children with SAM. In stunting, a progressive decline was noted in the children who did not respond to nutritional rehabilitation. It appears that malnutrition limits the ability of the gut to respond

to intestinal damage. What is unknown is whether the enteroendocrine cells that sense these nutrients are impaired or have been lost in malnutrition.

One of the difficulties in interpreting these data is the lack of reference range for young children. Progressive increments in GLP-2 concentrations have been noted during early neonatal development and the weaning period<sup>39</sup>. We included two groups of controls to provide a valid comparison. A further problem in children is difficulty in ascertaining the fed/fasted state. While we attempted to be clear on this point, mothers often use breast feeding to console their children and this may lead to misclassification. Composition of feeds also varies widely.

Intestinal GLP-2 release is nutrient stimulated, so nutrient availability partially determines GLP-2 release. We set out to examine whether there is any association between nutrient availability and GLP-2 release. We looked at seasonal parameters such as rainfall and temperature, which are known to affect food availability. We also took into consideration that the majority of Lusaka residents, very definitely those resident in our catchment area, are not farmers, so there is no direct link between produce and food availability. The cost of food and other essential household goods affects ones' purchasing power, and consequently the amount and type of nutrition present in a home. We factored this into our analysis by use of the FPI, matching the cost of food at the time the sample was collected, with the amount of GLP-2 in the sample. Stool pathogens in samples collected over the same time frame were analysed to determine if there was any association to GLP-2. All of the variables stated above were included in a multivariable logistic regression model with GLP-2 as the dependent variable. Rainfall, temperature, FPI and STEC showed a significant association with GLP-2, with temperature having a positive correlation coefficient. According to WHO, the exposure routes for STEC infection include food, animal contact, human to human contact, water and soil<sup>40</sup>. Based on this, we can conclude that our findings of STEC in baseline stool samples are an indication of the levels of environmental contamination present in a population with high levels of environmental enteropathy. The levels of the stool markers found in our samples were indicative of the levels of gut inflammation present in our cohort. Both of these markers have been studied in relation to environmental enteropathy and malnutrition<sup>41,42,43,44,45</sup> with inconsistent results. We observed a negative correlation between GLP-2 and myeloperoxidase in our study, indicative of the high inflammation present in those with low GLP-2. The reverse was observed in Calprotectin, but this result could have been confounded by the age and breastfeeding status of these children<sup>46,47</sup>.

Our findings suggest that there is a link between seasonality and GLP-2 release, consistent with its role as a nutrient responsive hormone. Teduglutide, a GLP-2 analogue has been successfully used in the treatment of IBD, and studies are now underway in severely malnourished children<sup>48</sup>

## References

1. Google. World Health Organization website. <http://www.who.int/en/news-room/fact-sheets/detail/diarrhoeal-disease>. Published May 2, 2017. Accessed September 26, 2018.
2. Liu L, Oza S, Hogan D, et al. Global, regional and national causes of under 5 mortality in 2000 – 15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet* 2016; 388:3027-35.
3. Rahman AE, Moinuddin M, Molla M, et al. Childhood diarrhoeal deaths in seven low- and middle-income countries. *Bull World Health Organ* 2014;92:664–71.
4. The United Nations Children’s Fund (UNICEF)/World Health Organisation (WHO). *Diarrhoea: Why children are still dying and what can be done*. Geneva; 2009.
5. Moore SR. Update on prolonged and persistent diarrhoea in children. *Curr Opin Gastroenterol* 2011;27:19-23.
6. Fagundes-Neto U. Persistent Diarrhoea: Still a Serious Public Health Problem in Developing Countries. *Curr Gastroenterol Rep* 2013;15:345.
7. Bandsma HJ, Sadiq K, Bhutta ZA. Persistent diarrhoea: current knowledge and novel concepts. *Paediatr Int Child H* 2018;39:41-7.
8. Owino V, Ahmed T, Freemark M, et al. Environmental Enteric Dysfunction and Growth Failure/Stunting in Global Child Health. *Pediatrics* 2016;138: e20160641.
9. Müller O, Krawinkel M. Malnutrition and health in developing countries. *Can Med Assoc J* 2005; 173:279-86.
10. Grenov B, Lanyero B, Nabukeera-Barungi N et al. Diarrhea, Dehydration and the Associated Mortality in Children with Complicated Severe Acute Malnutrition: A Prospective Cohort Study in Uganda. *J Pediatr* 2019;1-8.
11. Amadi BC, Kelly P, Mwiya M, et al. Intestinal and Systemic Infection, HIV and mortality in Zambian children with persistent diarrhea and malnutrition. *J Ped Gastroenterol Nutr* 2001;32: 550–4.
12. Salazar-Lindo E, Allen S, Brewster DR et al. Intestinal infections and environmental enteropathy: working group report of the second world congress of pediatric gastroenterology, hepatology, and nutrition. *J Pediatr Gastroenterol Nutr* 2003;39:S662–9.
13. Trehan I, Kelly P, Shaikh N et al. New insights into environmental enteric dysfunction. *Arch Dis Child* 2016; 101:741–4.

14. Engelstoft MS, Egerod KL, Holst B et al. A gut feeling for Obesity: 7TM sensors on enteroendocrine cells. *Cell Metab* 2008;8:447-9.
15. Burrin DG, Petersen Y, Stoll B et al. Glucagon- Like Peptide 2: A Nutrient – Responsive Gut Growth Factor. *J Nutr* 2001;131: 709-12.
16. Bloom SR. An enteroglucagon tumour. *Gut* 1972; 13: 520-3.
17. Drozdowski L, Thompson ABR. Intestinal hormones and growth factors: Effects on the small intestine. *World J Gastroenterol* 2009;15:385-406.
18. Lovshin J, Drucker DJ. New frontiers in the biology of GLP-2. *Regul Pept* 2000; 90:27-32.
19. Guan X, Karpen HE, Stephens J et al. GLP-2 Receptor Localizes to Enteric Neurons and Endocrine Cells Expressing Vasoactive Peptides and Mediates Increased Blood Flow. *Gastroenterology* 2006;130:150-64.
20. Pedersen J, Pedersen NB, Brix SW et al. The glucagon-like peptide 2 receptor is expressed in enteric neurons and not in the intestinal epithelium. *Peptides* 2015; 67:20-8
21. Yazbeck R, Howarth GS, Abbot CA. Growth factor-based therapies and intestinal disease: is Glucagon-like peptide-2 the new way forward? *Cytokine and Growth Factor Reviews* 2009; 20:175-84.
22. Drucker DJ. Glucagon – Like Peptide 2. *J Clin Endocrinol Metab* 2001; 86:1759 – 64C.
23. Janssen P, Rotondo A, Mule F et al. Review article: a comparison of glucagon-like peptides 1 and 2. *Aliment Pharmacol Ther* 2013;37:18-36.
24. Kelly P, Besa E, Zyambo K, et al. Endomicroscopic and Transcriptomic Analysis of Barrier Function and Malabsorption in Environmental Enteropathy. *PLoS Negl Trop Dis* 2016; 10: e0004600.
25. The United Nations Children’s Fund (UNICEF)/World Health Organisation (WHO). Integrated Management of Childhood Illness Chart Booklet 2014.
26. WHO. Guideline. Updates on the management of severe acute malnutrition in infants and children. Geneva: World Health Organization 2013.
27. Amadi B, Imikendu M, Sakala M, et al. Integration of HIV Care into Community Management of Acute Childhood Malnutrition Permits Good Outcomes: Retrospective Analysis of Three Years of a Programme in Lusaka. *PLoS ONE* 2016; 11: e0149218.
28. Central Statistical Office (CSO) [Zambia], Ministry of Health (MOH) [Zambia], and ICF. 2019. Zambia Demographic and Health Survey 2018: Key Indicators. Rockville, Maryland, USA: Central Statistical Office, Ministry of Health, and ICF.
29. WHO Multicentre Growth Reference Study Group, 2006
30. WHO Anthro(Windows). Version 3.2.2. Geneva: World Health Organization; 2011

<http://www.who.int/childgrowth/software/en/>

31. Claas EC, Burnham CD, Mazzulli T et al. Performance of the xTAG<sup>®</sup> Gastrointestinal Pathogen Panel, a Multiplex Molecular Assay for Simultaneous Detection of Bacterial, Viral and Parasitic Causes of Infectious Gastroenteritis. *J Microbiol and Biotechnol* 2013; 23:1041- 5.
32. Chisenga CC, Bosomorph S, Laban NM et al. Aetiology of Diarrhoea in Children Under Five in Zambia Detected Using Luminex xTAG Gastrointestinal Pathogen Panel. *Ped Infect Diseases* 2018; 3:8.
33. Jesuit Centre for Theological Reflection; Basic Needs Basket Google search engine. <http://www.jctr.org.zm>. Accessed September 20, 2018.
34. Egata G, Yemane B, Alemayehu W et al. Seasonal variation in the prevalence of acute undernutrition among children under five years of age in east rural Ethiopia: a longitudinal study. *BMC Pub Health* 2013, 13:864.
35. Shah M, Kathiiko C, Wada A et al. Prevalence, seasonal variation, and antibiotic resistance pattern of enteric bacterial pathogens among hospitalized diarrheic children in suburban regions of central Kenya. *Trop Med Health* 2016, 44:39.
36. Chao DL, Roose A, Roh M et al. The seasonality of diarrhoeal pathogens: A retrospective study of seven sites over three years. *PLoS Negl Trop Dis* 2019;13:e0007211.
37. Rogawski ET, Guerrant RL. The burden of enteropathy and “subclinical” infections. *Pediatr Clin North Am* 2017; 64:815-36.
38. Mutanen A, Pakarinen. Serum fasting GLP-1 and GLP-2 associate with intestinal adaptation in pediatric onset intestinal failure. *Clin Nutr* 2017; 36:1349-54.
39. Burrin DG, Petersen Y et al. Glucagon-Like Peptide 2: A Nutrient – Responsive Gut growth Factor. *J Nutr* 2001; 131:709-12
40. Hald T, Aspinalli W, Devieesschauwer B et al. World Health Organization Estimates of the Relative Contributions of Food to the Burden of Disease Due to Selected Foodborne Hazards:A Structured Expert Elicitation. *PLoS ONE* 2016; 11:e0145839
41. George CM, Burrowes V, Perin J et al. Enteric Infections in Young children are Associated with Environmental Enteropathy and Impaired Growth. *Trop Med Int Health*. 2018; 23:26-33
42. Arndt MB, Richardson BA, Ahmed T et al. Fecal Markers of Environmental Enteropathy and Subsequent Growth in Bangladeshi Children. *Am J Trop Med Hyg*. 2016; 95:694-701.
43. Guerrant RL, Leite AM, Pinkerton R et al. Biomarkers of Environmental Enteropathy, Inflammation, Stunting, and Impaired Growth in Children in Northeast Brazil. *PLoS ONE* 2016;11: e0158772.

44. Prata MG, Havt A, Bolick DT et al. Comparisons between myeloperoxidase, lactoferrin, calprotectin and lipocalin-2 as fecal biomarkers of intestinal inflammation in malnourished children. *J Transl Sci.* 2016;2:134-9.
45. Fahim SM, Das S, Gazi A et al. Association of intestinal pathogens with faecal markers of environmental enteric dysfunction among slum-dwelling children in the first 2 years of life in Bangladesh. *Trop Med Int Health.* 2018; 23: 1242-50.
46. George CM, Oldja L, Biswas S et al. Geophagy is associated with Environmental Enteropathy and Stunting in Children in Rural Bangladesh. *Am J Trop Med Hyg.* 2015;92: 1117-24.
47. Li F, Ma J, Geng S, et al. Fecal calprotectin concentrations in Healthy Children Aged 1-18 Months. *PLoS ONE* 2015; 10: e0119574
48. Li F, Ma J, Geng S, et al. Comparison of the different kinds of feeding on the levels of fecal calprotectin. *Early Hum Dev.*2014; 90;471-5.
49. Kelly P, Bell L, Amadi B, et al. TAME trial: a multi-arm phase II randomized trial of four novel interventions for malnutrition enteropathy in Zambia and Zimbabwe – a study protocol. *BMJ Open* 2019 ;9: e027548.

Figure legends






Figure 1A: GLP-2 levels in children with acute diarrhoea during illness and 3 weeks after their last diarrhoea episode.






**Figure 1B:** GLP-2 levels in acute and persistent diarrhoea compared to controls



**Figure 2A:** GLP-2 levels in stunted children at baseline, 3 months and prior to endoscopy (fasted non-responders) compared to healthy controls from the same community.



**Figure 2B:** Repeated GLP-2 measurements in children designated non-responders at baseline, 3 months, and in the fed and fasted state. Children were fed ad libitum.

## Table Legends

**Table 1:** Baseline characteristics of children

	<b>Group 1 (n=28)</b>	<b>Group 2 (n=21)</b>	<b>Group 3 (n=75)<sup>1</sup></b>	<b>Group 4 (n=181)</b>	<b>Group 5 (n=19)</b>
Months of sample collection	January-February 2016	May 2014-February 2015	August-October 2014	October 2016-September 2018	April 2017-April 2019
Sex: Male, n (%)	17 (66%)	14 (67%)	30 (40%)	107(59%)	6 (32%)
Age (median IQR)	16.5 (11.5,22)	17 (12,21)	20 (16,29)	11 (7,14)	4 (3,5)
Weight for Age (WAZ)	-1.78(-2.55, -0.69)	-3.48(-4.98, -2.7)	-1.21(-1.98, -0.52)	-2.57(-3.14, -2.21)	0.08 (-0.86, 0.55)
Weight for Length (WLZ)	-0.92(-1.68,0.38)	-2.80 (-4.11, -1.61)	-0.27(-1.07,0.32)	-1.23(-1.86, -0.59)	0.73 (0.21,1.49)
Length for Age (LAZ)	-2.24(-2.69, -1.38)	-2.66 (-4.36, -1.72)	-2.25(-2.9, -1.18)	-2.99(-3.56, -2.57)	- 0.81(-1.01, -0.17)
Samples available	Plasma (fed state) during illness and after convalescence	Plasma (fasted state) during illness	Plasma (healthy in fed state)	Serum (fed and fasted state); stool for pathogen analysis	Serum (fed only); stool for pathogen analysis

<sup>1</sup>anthropometric data missing for one healthy control. Group 1 – acute diarrhoea; Group 2 – Persistent diarrhoea; Group 3 – Controls for Groups 1 & 2; Group 4 – Stunted children; Group 5 – Controls for Group 4. Fed state assumed to be fed, not timed in relation to sample collection

**Table 2:** Table of univariable and multivariable logistic regression analysis of GLP-2 concentrations in

Variable	Univariable Analysis		Multivariable Analysis	
	Coefficient (B)	p(>0.05)	Coefficient (B)	p(>0.05)
Food Price Index	-0.002	0.06	-0.002	<b>0.05</b>
Minimum Temperature	0.029	0.67	0.354	<b>&lt;0.01</b>
Maximum Temperature	0.177	<b>&lt;0.01</b>		
Rainfall	-0.004	<b>&lt;0.01</b>	-0.006	<b>&lt;0.01</b>
<i>Adenovirus</i> (n=20)	-0.642	0.25		
<i>Norovirus</i> (n=104)	-0.457	0.22		
<i>Rotavirus</i> (n=41)	-0.558	0.16		
<i>Campylobacter</i> (n=78)	0.289	0.83		
Enterotoxigenic <i>Escherichia coli</i> (n=117)	0.485	0.23		
<i>Salmonella</i> (n=135)	-0.864	0.86		
Shiga toxin-producing <i>Escherichia coli</i> (n=19)	-1.366	<b>0.02</b>	-1.409	<b>0.01</b>
<i>Shigella</i> (n=109)	0.025	0.95		
<i>Cryptosporidium spp</i> (n=43)	-0.318	0.43		
<i>Giardia</i> (n=116)	0.138	0.73		

relation to measures of seasonality