

**Evaluation of the effectiveness of a tailored mobile application in increasing the duration of wear of thermoplastic retainers: a randomised controlled trial**

Journal:	<i>European Journal of Orthodontics</i>
Manuscript ID	EJO-2019-RCT-0395.R3
Manuscript Type:	Randomized Controlled Trial (RCT)
Research Topics:	Clinical trials, systematic reviews & meta-analyses, Stability & retention, basic science

SCHOLARONE™  
Manuscripts

## Evaluation of the effectiveness of a tailored mobile application in increasing the duration of wear of thermoplastic retainers: a randomised controlled trial

**Background:** The 'My Retainers' mobile application is a patient-informed intervention designed to enhance removable retainer wear and improve patient experiences during the retention phase.

**Objectives:** To evaluate the effect of receiving 'My Retainers' application on objectively-assessed thermoplastic retainer (TPR) wear time, stability and periodontal outcomes and participants' experiences and knowledge related to retainers.

**Materials and Methods:** Eighty-four participants planned for removable retention with TPRs, were assigned to either receive the 'My Retainers' application or to a control not receiving electronic reminders during the 3-month period. Randomisation was based on computer-generated random numbers and allocation was concealed using opaque sealed envelopes. **The primary outcome was objectively-assessed retainer wear recorded using an embedded TheraMon® micro-electronic sensor. Secondary outcomes** including irregularity of the maxillary and mandibular incisors, plaque levels, bleeding on probing and probing depth were assessed at baseline and 3-month follow-up; and analysed using a series of mixed-models. Experiences and knowledge related to orthodontic retainers were recorded using questionnaires. The outcome assessor was blinded when possible

**Results:** Receipt of the mobile application resulted in slightly higher median wear time (0.91 hours/day); however, this difference was not statistically significant ( $P= 0.56$ ; 95% confidence interval [CI]: -2.19, 4.01). No significant differences were found between the treatment groups in terms of stability ( $P= 0.92$ ; 95% CI: -0.03, 0.04), plaque levels ( $P= 0.44$ ; 95% CI: -0.07, 0.03), bleeding on probing ( $P= 0.61$ ; 95% CI: -0.05, 0.03) and probing depth ( $P= 0.79$ ; 95% CI: -0.09, 0.07). Furthermore, similar levels of patients' experiences ( $P= 0.94$ ) and knowledge related to retainers ( $P= 0.26$ ) were found. However,

marginally better levels of knowledge was observed in the intervention group.  
No harms were observed.

**Limitations:** A relatively short follow-up period with study conducted within a single-centre in a university-based hospital.

**Conclusions:** Provision of the bespoke 'My Retainers' application did not lead to an improvement in adherence with TPR wear over a 3-month follow-up period. Further refinement and research are required to develop and investigate means of enhancing adherence levels.

**Keywords:** vacuum-formed retainer; Essix-type retainer; compliance; mHealth.

**ClinicalTrials.gov Identifier:** NCT03224481.

## Introduction

Maintenance of post-treatment orthodontic outcomes hinges on the levels of adherence to orthodontic retention. Barriers to removable retainer wear including negative impact on quality of life, forgetfulness and a lack of appreciation of the importance of retainer wear have been identified in previous research (1, 2). The centrality of the patient-clinician relationship in terms of sharing concerns and frequency of follow-up appointments has also been highlighted in qualitative research (1). Notwithstanding this, suboptimal adherence has been exposed in prospective studies with just one-third of participants claiming to be adherent with Essix-type retainer wear 2 years into the retention phase (3). The importance of developing and evaluating relevant interventions to enhance wear and ameliorate negative experiences associated with orthodontic retainers is therefore clear (4).

The unprecedented access to mobile phones has raised the potential for use for personalised healthcare management and delivery of health-related information (5). A total of 241 patient-centred orthodontic mobile applications were developed in 2018, representing a three-fold increase since 2014 (6). However, relatively little prospective assessment of the effectiveness of these approaches in orthodontics has been undertaken (4). In a recent randomised controlled trial, access to moderated WhatsApp groups involving photo sharing and monthly ranking was postulated to improve Hawley retainer wear, based on the superior stability outcomes in terms of inter-canine width at 1-

1  
2  
3  
4 year follow-up (7). However, neither objective nor subjective wear time was  
5  
6 assessed (7). Nevertheless, it is conceivable that receipt of an electronic  
7  
8 reminder can enhance adherence to removable orthodontic retainer wear.  
9  
10  
11 Additionally, receiving a mobile application has been shown to be effective in  
12  
13 terms of improving oral hygiene (8, 9), and reducing the occurrence of white  
14  
15 spot lesions and caries (9), improving attendance and reducing the duration of  
16  
17 treatment (10).  
18  
19  
20  
21  
22  
23

24  
25 In a recent qualitative study, participants advocated the use reminders through  
26  
27 a mobile application to facilitate adherence to removable functional appliance  
28  
29 wear (11). Receipt of electronic reminders is a passive process involving  
30  
31 automatic notification when the reminder is received. Furthermore, these  
32  
33 approaches offer the potential to motivate wear and educate on the importance  
34  
35 of retainer wear. Addressing patients' needs by capturing preferences can  
36  
37 help make the intervention appealing and, therefore, potentially improve  
38  
39 outcomes. The 'My Retainers' mobile application is a patient-informed  
40  
41 intervention and was developed following a rigorous methodology involving  
42  
43 triangulation of the findings of two qualitative methods (12).  
44  
45  
46  
47  
48  
49  
50  
51

52  
53 The primary aim of this study was to analyse the effect of receiving the 'My  
54  
55 Retainers' mobile application on adherence to thermoplastic retainer wear.  
56  
57 The secondary aims were to investigate the effects of receiving the mobile  
58  
59 application on the stability of the outcome and periodontal health following  
60

1  
2  
3  
4 removal of fixed appliances, and patient experience and knowledge related to  
5  
6 orthodontic retainers.  
7  
8  
9

## 10 11 12 13 14 **Materials and methods** 15

16  
17  
18 Ethical approval was obtained from the East of England, Cambridge Central  
19 Research Ethics Committee (16/EE/0189). **The trial protocol was registered**  
20  
21 **prior to study commencement (ClinicalTrials.gov Identifier: NCT03224481).**  
22  
23

24  
25 Participants were recruited prior to planned removal of the appliances at the  
26  
27 Institute of Dentistry, Barts and The London School of Medicine and Dentistry.  
28  
29 The inclusion criteria were: aged 12 to 21 years; planned for removable  
30  
31 retention with thermoplastic retainers (TPRs); on no medication known to have  
32  
33 an effect on gingival health; and in the permanent dentition. The exclusion  
34  
35 criteria were inability to access or peruse a compatible smart phone (iPhone;  
36  
37 **Apple Inc.**); cleft lip and palate or other craniofacial anomalies; and history of  
38  
39 periodontal disease. An information sheet was provided with oral and written  
40  
41 consent obtained from participants agreeing to take part.  
42  
43  
44  
45  
46  
47  
48  
49

50  
51  
52 Based on previous research (13) with a non-adherence rate of 31%  
53  
54 characterised by wear of the appliance for less than 2 hours daily, a minimum  
55  
56 of 68 participants (34 in each group) was required with a power of 80% to  
57  
58 detect a minimum difference of 25% in adherence rates at the 0.05 level of  
59  
60

1  
2  
3 statistical significance. To compensate for a drop-out rate of at least 20%, the  
4  
5  
6 final number enrolled in the trial was 84.  
7  
8  
9

10  
11 Randomisation was based on computer-generated random numbers and was  
12  
13 stratified in a ratio of 1:1 in relation to gender. Allocation was concealed from  
14  
15 the treating clinician using an opaque, sealed envelope system. Participants  
16  
17 in the intervention group received access to the 'My Retainers' mobile  
18  
19 application via a unique identification code (12). Participants in the control  
20  
21 group did not have access to the mobile application.  
22  
23  
24  
25  
26  
27  
28  
29

30 The primary outcome was objective wear time (hours per day). The following  
31  
32 secondary outcomes were also assessed:  
33

- 34 - Maxillary and mandibular Little's irregularity index (14)
- 35  
36 - Periodontal outcomes including: plaque scores, bleeding on probing,  
37  
38 and probing depth
- 39  
40 - Subjective wear time
- 41  
42 - Patient experiences and knowledge related to retention
- 43  
44  
45  
46  
47

48 Standardised oral hygiene instructions were given to all participants at debond  
49  
50 and recall appointments. Information related to oral hygiene practices were  
51  
52 recorded at baseline (T0). Maxillary and mandibular TPRs (Essix ACE® Plastic  
53  
54 1mm in thickness (DENTSPLY)) were fitted 7-10 days following debond. All  
55  
56 participants were instructed to wear TPRs on a full-time basis (22 hours) for 6  
57  
58 months, followed by part-time wear (8 hours) for a further 6 months. A  
59  
60

1  
2  
3  
4 TheraMon<sup>®</sup> micro-electronic sensor (MC Technology GmbH, Hargelsberg,  
5  
6 Austria) was embedded in the maxillary TPR in all participants following a  
7  
8 standardised laboratory technique (KM'L) (Supplementary material 1).  
9  
10  
11 Participants in both groups had a follow-up appointment scheduled at 3  
12  
13  
14 months (T1) following removal of the appliances (T0).  
15  
16  
17  
18

19  
20 A reading station facilitated data transfer to an encrypted cloud database using  
21  
22 TheraMon<sup>®</sup> Azure reader client software (version 1.2.1.1; MC Technology  
23  
24 GmbH, Hargelsberg, Austria). Data were transferred using radio-frequency  
25  
26 identification technology. Appliance wear was recorded within a specific  
27  
28 temperature range (33.5°C and 38.5°C). The TheraMon<sup>®</sup> micro-electronic  
29  
30 sensor records temperature at 15-minute intervals; as such, data could be  
31  
32 restored for up to 100 days. Subjective data pertaining to wear involved  
33  
34 completion of a retainer wear chart in the control group (Figure 1), and use of  
35  
36 a calendar tool within the mobile application in the intervention group (Figure  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
2).

58 Impressions of both dental arches were taken at T0 and T1 using hydrophilic  
59  
60 vinyl polysiloxane (Virtual; Ivoclar Vivadent, Schaan, Lichtenstein) and study  
models were made from Orthodontic Plaster (ISO type 2; Whip Mix  
Corporation, Louisville, KY, USA). Periodontal assessment was undertaken at  
T0 and T1. Each tooth surface was divided into thirds using vertical lines based  
on the morphology and position of the dental papilla. The periodontal



1  
2  
3  
4 measures were scored clinically on the labial/buccal and palatal/lingual  
5  
6 surfaces in both arches from first molar to first molar, at 6 sites per tooth by  
7  
8 one researcher (DA) and included the following:  
9

- 10  
11 - Plaque scores: A liquid disclosing solution (Plaqsearch™, TePe®,  
12  
13 Malmö, Sweden) was applied using a swab pressed against each  
14  
15 papilla, followed by 10ml water rinse. Plaque was scored as present or  
16  
17 absent.  
18  
19
- 20  
21 - Bleeding on probing: A binary assessment of bleeding on probing was  
22  
23 undertaken with a maximum waiting time of 15 seconds.  
24  
25
- 26  
27 - Probing depth: measured to the nearest 0.5mm from the gingival  
28  
29 margin to the base of the gingival sulcus using a Williams probe.  
30  
31

32  
33 Participants in both groups were asked to complete a questionnaire at T1  
34  
35 concerning their experiences and knowledge in relation to TPRs  
36  
37 (Supplementary material 2).  
38  
39  
40  
41

42  
43 Maxillary and mandibular Little's irregularity index (14) were measured by one  
44  
45 researcher (DA) using a digital caliper (150mm DIN 862, ABSOLUTE  
46  
47 Digimatic caliper, model 500-191U; Mitutoyo, Andover, Hampshire, UK) with  
48  
49 a resolution of  $\pm 0.01\text{mm}$  (19). Mean objectively-assessed hours of retainer  
50  
51 wear was obtained from cloud software (TheraMon Azure®, version 1.2.1.11)  
52  
53 and graphical display of the data was evaluated to detect lack of retainer wear  
54  
55  
56  
57  
58 over a period of three consecutive days or more.  
59  
60

1  
2  
3  
4 Participants in both groups were aware of being monitored. Blinding of either  
5  
6 the operator or the participants to the allocated arm during treatment was not  
7  
8 possible for the periodontal assessment. However, the use of coded study  
9  
10 models and data ensured that the researcher was kept blind to the treatment  
11  
12 group when undertaking measurements and during data analysis. The  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Participants in both groups were aware of being monitored. Blinding of either the operator or the participants to the allocated arm during treatment was not possible for the periodontal assessment. However, the use of coded study models and data ensured that the researcher was kept blind to the treatment group when undertaking measurements and during data analysis. The statistician was also kept blind to group allocation.

In cases in which replacement of the TPR was required, reasons were recorded and the same micro-electronic sensor was used, where possible. If a participant opted to have a TPR without a micro-electronic sensor, a new TPR was fitted and the participant was retained in the study.

As the data were not normally distributed, medians and inter-quartile range (IQR) are presented. Imputation of missing data was undertaken to account for losses and to compensate for uncertainty surrounding missing values. Missing baseline data for periodontal (plaque levels, bleeding on probing and probing depths) and stability outcomes were imputed using the corresponding mean for each group (15). Objective data pertaining to retainer wear were imputed by creating new datasets (n= 40 iterations) with 10 values imputed by the software. For each of these datasets, estimates were calculated by fitting a corresponding separate model (16). Consequently, the estimates were combined to produce the average final estimate (17). The linear regression model accounted for treatment group, available subjective data as well as

1  
2  
3 complete observation variables including age and gender. This permitted  
4  
5  
6 imputation of missing values using values drawn from a distribution based on  
7  
8  
9 observed participant values with similar baseline characteristics. A series of  
10  
11 mixed-models were then fitted in the imputed dataset accounting for  
12  
13 correlation. The level of statistical significance was set at 0.05 with all analyses  
14  
15 undertaken using the Stata statistical software package (version 15.1;  
16  
17 StataCorp, College Station, Tex). The exact Mann-Whitney test was used to  
18  
19 compare knowledge and experience outcomes between the treatment groups.  
20  
21  
22  
23  
24 The analysis was performed in R software (18).

25  
26  
27  
28  
29 An online course was completed (DA) to facilitate familiarisation with  
30  
31 measurement of periodontal outcomes. For stability measurements, intra-  
32  
33 examiner reliability was performed on 10 randomly selected study models, 4  
34  
35 weeks after the initial measurement. Intra-examiner reliability in relation to  
36  
37 plaque scoring was assessed by repeating measurements on 10 intra-oral  
38  
39 photographs at a 4-week interval. Probing depth measurements were  
40  
41 repeated on 10 healthy volunteers 30 minutes apart. Differences between the  
42  
43 repeated measurements relating to stability, mean probing depths and mean  
44  
45 plaque scores per tooth were assessed using intraclass correlation. Excellent  
46  
47 agreement was observed for stability (intraclass correlation coefficient; ICC:  
48  
49 0.97) and periodontal outcomes including plaque score (ICC: 0.96) and  
50  
51 probing depth (ICC: 0.93).  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Results

Full trial dataset is available online (*DOI*: <https://doi.org/10.17636/01059856>).

Eighty-four participants were enrolled and randomised with 42 participants per group and equal gender distribution (Table 1, Figure 3). Overall, the groups were well-matched in terms of age, duration of orthodontic treatment and self-reported oral hygiene practices (Table 1). Slightly more participants were treated without extractions in the control group.

Stability and periodontal data were recorded for 80 participants at baseline with missing values imputed, and 64 at 3-month follow-up (Figure 3) with retainer failures recorded (Table 2). The mean duration from T0 to T1 was 100.78 (standard deviation (SD) 23.49) days.

The median duration of objectively-assessed retainer wear was slightly higher in the intervention (7.25 hours/day) than control group (6.21 hours/day). After adjusting for confounders, the median wear was 0.91 hours/day higher in the intervention group ( $P=0.56$ ; 95% CI: -2.19, 4.01 hours/day); however, the between-group difference was not statistically significant ( $P=0.56$ ) (Table 3).

A period of no wear for three consecutive days or more was observed in more than half of the sample in both groups (Table 3). The median percentage of days in which the retainers were worn for less than 8 hours a day and a minimum of 2 hours of continuous use was 44.3% in the intervention group, and 53.3% in the control group (Table 3). Objectively-assessed retainer wear

1  
2  
3 data were available for a mean of 87.41 (SD: 20.1) days. A median  
4  
5  
6 discrepancy of 4.96 hours was found between subjective and objective wear  
7  
8  
9 time, based on 30 participants with both measures available.  
10

11  
12  
13  
14 No significant difference between the treatment groups was observed in terms  
15  
16 of incisor irregularity ( $P= 0.92$ ) and periodontal outcomes including plaque  
17  
18 scores ( $P= 0.44$ ), bleeding on probing ( $P= 0.61$ ) and probing depth ( $P= 0.79$ )  
19  
20  
21 (Table 3).  
22  
23  
24  
25

26  
27 In terms of patient experiences, the highest scores (4 and 5) were most  
28  
29 frequently selected in both groups, indicating similar levels of satisfaction in  
30  
31 both treatment groups (Table 4). Levels of knowledge were marginally better  
32  
33 in the intervention group (Table 4). However, no significant difference was  
34  
35 found between intervention and control group for both outcomes (Table 5).  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Discussion

Receipt of the mobile application did not seem to significantly improve objectively-assessed adherence levels, stability, periodontal outcomes, patient experiences and knowledge related to orthodontic retainers at 3-month follow-up. The limited benefit of interventions directed at enhancing adherence levels with orthodontic retainers has been exposed in previous research (4). This may relate to the complex and multi-faceted nature of adherence with extraneous factors including associated negative impact on quality of life and pragmatic issues related to retainer wear also being important (1).

The multitude of functions built in the 'My Retainers' mobile application were designed to address reported barriers to retainer wear (1,12). For example, a reminder system was included to overcome forgetfulness. An exhaustive list of frequently-asked questions and the ability to contact the researcher were included to address potential concerns related to retainer wear. Furthermore, this intervention was underpinned by key behavioural change theories (12). The potential benefit of utilising a combination of approaches to behaviour change in developing internet-based health-related interventions was highlighted in a previous systematic review (19).

1  
2  
3  
4 The use of supplementary methods for information provision such as written,  
5  
6 audio and visual information has been shown to result in improvement in recall  
7  
8 of orthodontic information (20-22). On the corollary, participants in the mobile  
9  
10 application group exhibited slightly higher levels of knowledge; however,  
11  
12 retainer wear remained suboptimal. Similar findings have been reported within  
13  
14 medical literature with no clear association between patients' knowledge  
15  
16 concerning diabetes and adherence behaviours (23). The limited effect of the  
17  
18 mobile application on adherence may be explained by inadequate usage of  
19  
20 the different features. This was evident in the median number of days in which  
21  
22 the retainer wear was logged (n= 11; IQR: 51) and the limited interaction in  
23  
24 terms of the number of emails sent by participants (n= 6) throughout the study.  
25  
26 However, user engagement with the intervention, the number of times  
27  
28 participants accessed the mobile application, consistency of use and time  
29  
30 spent viewing its content are unclear. Unknown barriers to the limited  
31  
32 effectiveness of the mobile application will be addressed using an explanatory  
33  
34 qualitative study in keeping with previous approaches (24).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 The average wear time was slightly higher in the intervention compared to the  
49  
50 control group; however, the difference was not statistically significant.  
51  
52 Nevertheless, the median objectively-assessed retainer wear was just 28.2%  
53  
54 and 33% of 22 hours stipulated in the control and intervention groups,  
55  
56 respectively. Moreover, participants were aware of being monitored in the  
57  
58  
59  
60

1  
2  
3  
4 current study with the latter thought to lead to artificially high wear levels.  
5  
6 Micro-electronic timers have been shown to under-report wear duration by the  
7  
8 order of 4% (25); this discrepancy was dwarfed by the low objective readings  
9  
10 identified among the present group of participants. In a previous study with  
11  
12 similar stipulated wear time, better levels of adherence (45.5-60%) were found  
13  
14 with Hawley retainers at 3-month follow-up (26). However, details of  
15  
16 randomisation and allocation concealment were not reported in the latter  
17  
18 study. Mean wear rates varied significantly (0-19.9 hours/day) and participants  
19  
20 over-estimated wear by an average of 5.6 hours daily (26). It is also possible  
21  
22 that the visibility of the Hawley retainer with associated labial bow may serve  
23  
24 as a reminder to wear this type of retainer among both patients and peers.  
25  
26  
27  
28  
29  
30  
31  
32  
33

34  
35 A number of participants in the current study relayed concerns in relation to  
36  
37 the appearance and bulk of the retainer associated with the indwelling micro-  
38  
39 electronic sensor. Related data were collected at 6-month follow-up; the latter  
40  
41 will be analysed in future. It is conceivable that this may have contributed to  
42  
43 suboptimal adherence levels. Furthermore, patient motivation and attitudes  
44  
45 towards treatment have been shown to influence adherence levels in  
46  
47 orthodontics, pointing to overlapping patterns of behaviour (27, 28).  
48  
49  
50  
51  
52  
53

54  
55 No significant difference was found between the groups in relation to the  
56  
57 stability outcomes. This may relate to the comparable objectively-assessed  
58  
59  
60



1  
2  
3 adherence levels in both groups and to the relatively short period of follow-up.  
4  
5  
6 Although objectively-assessed retainer wear may provide an overall  
7  
8 assessment of adherence levels over a particular observation period, it does  
9  
10 not reflect patterns, consistency and distribution of wear. Fluctuations in  
11  
12 adherence levels were previously observed with removable and functional  
13  
14 appliances (11, 29). Similar findings were observed in the current study, with  
15  
16 no retainer wear over at least three consecutive days observed in more than  
17  
18 half of the sample. Similarly, a period of no wear has been shown with  
19  
20 headgear (30), and removable functional appliances (31), in 30% and 13% of  
21  
22 the duration of the study, respectively. This period of no wear, negatively  
23  
24 influenced the transverse changes obtained with functional appliances (31).  
25  
26 However, the implications of extended periods of an absence of wear may be  
27  
28 particularly problematic with retainers, with sustained periods of non-  
29  
30 adherence risk irreversible impairment of retainer fit.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 The content of the mobile application also included general dental and oral  
44  
45 health information (12). No significant difference was observed between both  
46  
47 groups in terms of the periodontal measures. In previous research, superior  
48  
49 periodontal outcomes were found at 1-month follow-up in patients receiving a  
50  
51 mobile application including notification messages and access to an  
52  
53 educational video focusing on oral hygiene (8); however, detailed description  
54  
55 of the intervention was not reported. Similarly, an interactive intervention  
56  
57  
58  
59  
60

1  
2  
3 involving WhatsApp group messaging resulted in better periodontal outcomes  
4  
5  
6 at 1-year follow-up; however, the difference was not significant at 3-month  
7  
8  
9 follow-up (9). The use of a mobile application to allow tracking of toothbrushing  
10  
11  
12 frequency and duration did not result in a significant difference in plaque  
13  
14  
15 accumulation and gingival inflammation at 3-month follow-up (32). Therefore,  
16  
17  
18 it seems that differences in periodontal outcomes may be observed at longer  
19  
20  
21 follow-up periods.  
22  
23

24  
25 Fixed retention offers superior preservation of the alignment of mandibular  
26  
27  
28 anterior teeth in the long term (3). However, thermoplastic retainers continue  
29  
30  
31 to be used due to their acceptability, simplicity and cost-effectiveness (33).  
32  
33  
34 Removable retainers may be prescribed for those exhibiting suboptimal oral  
35  
36  
37 hygiene. This might explain the significant plaque accumulation and bleeding  
38  
39  
40 on probing noted at baseline in both groups. Notwithstanding this,  
41  
42  
43 thermoplastic retainers may impede flushing of saliva from dental surfaces  
44  
45  
46 resulting in a significant increase in *Streptococcus mutans* and *Lactobacillus*  
47  
48  
49 counts (34). An initial phase of full-time wear (35, 36) is often prescribed with  
50  
51  
52 removable retention; however, little is known about the effect of prolonged  
53  
54  
55 removable retainer wear on periodontal health. Interestingly, a reduction in  
56  
57  
58 plaque and calculus accumulation, gingival inflammation and bleeding on  
59  
60  
61 probing was noted following transition to part-time thermoplastic retainer wear  
62  
63  
64 in a previous clinical trial (37).

1  
2  
3  
4  
5  
6 The type of material used to fabricate the TPR in the current study (Essix ACE®  
7  
8 Plastic) was found to have superior wear resistance in comparison to other  
9  
10 types of commercially available materials in an *in vitro* study (38). However, a  
11  
12 substantial proportion of retainers required replacement (n= 22) mainly due to  
13  
14 poor fit and breakage, despite the short period of follow-up of the current study.  
15  
16 Lower breakage rates were observed in a previous randomised controlled trial  
17  
18 (RCT), in which only 6.6% of the participants reported breakage with vacuum-  
19  
20 formed retainers in the first 6 months of retention (33). This could be explained  
21  
22 by the difference in the type and thickness of the material used in the previous  
23  
24 study (1.5mm) (33). It is also possible that the incorporation of the micro-  
25  
26 sensor in the present study may have predisposed to fracture.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 The stipulated wear time in the current study was in line with previous research  
38  
39 (39). However, there is some evidence to suggest similar outcomes with part-  
40  
41 time wear (40). Part-time wear is also regarded as more realistic and  
42  
43 achievable with minimal impact on daily activities (1, 11). This is likely to  
44  
45 explain the part-time wear of Twin Blocks despite full-time prescription with  
46  
47 mean wear rates of 12 hours daily observed in the full-time group and 8 hours  
48  
49 daily in the prescribed part-time group (41). It is conceivable that the relatively  
50  
51 disappointing wear times reported with retainers in the present study may  
52  
53 reflect both complacency as well as a lack of understanding of the implications  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 of poor wear in this cohort (42). In the current study, stratified randomisation  
4 was undertaken to ensure balanced gender distribution in the treatment  
5  
6 groups. This was considered important as adherence levels to intra-oral  
7  
8 removable appliance wear have been shown to vary significantly based on  
9  
10 gender (43).  
11  
12  
13  
14  
15  
16  
17  
18

### 19 **Limitations and generalisability**

20  
21 Drop-out rates in orthodontic RCTs is typically of the order of 13% of those  
22 recruited (44). This was accounted for in the current trial statistically by  
23  
24 imputation of missing data as well as by inflation of the sample size by 20% in  
25  
26 order to retain adequate power. However, the drop-out rate was 24%. A  
27  
28 greater proportion of drop-outs are typical of trials concerning retention  
29  
30 particularly as no active treatment is being provided (3), highlighting the  
31  
32 importance of making adequate allowance for drop-outs in future research on  
33  
34 orthodontic retention. Furthermore, loss of objective adherence data was  
35  
36 inevitable due to the capacity of the TheraMon<sup>®</sup> micro-electronic sensor to  
37  
38 restore data up to 100 days with a measurement interval set to 15 minutes.  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 The study was undertaken in one university hospital in which orthodontic  
49  
50 treatment is funded through a national healthcare system. A significant  
51  
52 difference between university hospital and private practice in terms of  
53  
54 adherence levels has been exposed in previous research (43). Therefore, the  
55  
56 applicability to other settings hinges on comparability of patient characteristics.  
57  
58  
59  
60

1  
2  
3  
4 The relatively short follow-up period might limit the holistic evaluation of the  
5  
6 intervention. Notwithstanding this, adherence to removable appliance wear  
7  
8 also tends to reduce over time (1); it is therefore conceivable that the benefit  
9  
10 of the mobile application may become more apparent over a more prolonged  
11  
12 follow-up period. **We therefore plan to follow up participants in the current**  
13  
14 **study up to one year post-treatment.**  
15  
16  
17  
18  
19  
20  
21

## 22 Conclusions

23  
24  
25  
26 Receipt of a bespoke mobile application did not result in improvement in  
27  
28 adherence to thermoplastic retainer wear, stability and periodontal outcomes  
29  
30 and experience with retainers in the short term. Knowledge concerning  
31  
32 orthodontic retainers was slightly higher in the intervention group; however,  
33  
34 the difference was not statistically significant. Evaluation of the effectiveness  
35  
36 of the mobile application over a longer follow-up period as well as further  
37  
38 refinement are required.  
39  
40  
41  
42  
43

44 **Supplementary material is available at European Journal of Orthodontics online.**  
45  
46  
47  
48

## 49 Conflict of interest statement:

50  
51 The authors declare that they have no competing interests.  
52  
53  
54  
55

## 56 Acknowledgements:

57  
58  
59  
60

1  
2  
3 We would like to thank Anas El-Huni, Yan Huang and all orthodontic postgraduates  
4 at the Royal London Hospital for their help in recruitment. We would also like to thank  
5 UCL Health Creatives for their technical assistance in developing the mobile  
6 application.  
7  
8  
9

### 10 Funding:

11  
12 This work was supported by a funding from the European Orthodontic Society. DA's  
13 PhD is funded by the Saudi Arabian Cultural Bureau.  
14  
15  
16  
17  
18  
19  
20  
21

### 22 References

- 23 1. Al-Moghrabi, D., Colonio Salazar, F.B., Johal, A., Fleming, P.S. (2019). Factors  
24 influencing adherence to vacuum-formed retainer wear: A qualitative study. *Journal*  
25 *of Orthodontics*. 46, 212-219
- 26 2. Al-Moghrabi, D., Johal, A., Fleming, P.S. (2017). What are people tweeting about  
27 orthodontic retention? A cross-sectional content analysis. *American Journal of*  
28 *Orthodontics and Dentofacial Orthopedics*. 152, 516-522.
- 29 3. Al-Moghrabi, D., Johal, A., O'Rourke, N., Donos, N., Pandis, N., Gonzales-Marin, C.,  
30 Fleming, P.S. (2018). Effects of fixed vs removable orthodontic retainers on stability  
31 and periodontal health: 4-year follow-up of a randomized controlled trial. *American*  
32 *Journal of Orthodontics and Dentofacial Orthopedics*. 154, 167-174.
- 33 4. Al-Moghrabi, D., Salazar, F.C., Pandis, N., Fleming, P.S. (2017). Compliance with  
34 removable orthodontic appliances and adjuncts: A systematic review and meta-  
35 analysis. *American Journal of Orthodontics and Dentofacial Orthopedics*. 152, 17-32.
- 36 5. Food and Drug Administration (2015). Mobile Medical Applications- Guidance for  
37 Industry and Food and Drug Administration Staff. Available at:  
38 <https://www.fda.gov/media/80958/download> [Accessed July 4th, 2019].
- 39 6. Siddiqui, N.R., Hodges, S., Sharif, M.O. (2019). Availability of orthodontic smartphone  
40 apps. *Journal of Orthodontics*. 46, 235-241
- 41 7. Zotti, F., Zotti, R., Albanese, M., Nocini, P.F., Paganelli, C. (2019). Implementing post-  
42 orthodontic compliance among adolescents wearing removable retainers through  
43 Whatsapp: a pilot study. *Patient Preference and Adherence*. 13, 609-615.
- 44 8. Alkadhi, O.H., Zahid, M.N., Almanea, R.S., Althaqeb, H.K., Alharbi, T.H., Ajwa, N.M.  
45 (2017). The Effect of using Mobile Applications for Improving Oral Hygiene in Patients  
46 with Orthodontic Fixed Appliances: A Randomised Controlled Trial. *Journal of*  
47 *Orthodontics*. 44, 157-163
- 48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
9. Zotti, F., Dalessandri, D., Salgarello, S., Piancino, M., Bonetti, S., Visconti, L., *et al.* (2016). Usefulness of an app in improving oral hygiene compliance in adolescent orthodontic patients. *The Angle Orthodontist*. 86, 101-107.
10. Li, X., Xu, Z.R., Tang, N., Ye, C., Zhu, X.L., Zhou, T., *et al.* (2016). Effect of intervention using a messaging app on compliance and duration of treatment in orthodontic patients. *Clinical Oral Investigations*. 20, 1849-1859.
11. El-Huni, A., Colonio Salazar, F.B., Sharma, P.K., Fleming, P.S. (2019). Understanding factors influencing compliance with removable functional appliances: A qualitative study. *American Journal of Orthodontics and Dentofacial Orthopedics*. 155, 173-181.
12. Al-Moghrabi D, Colonio-Salazar F.B., Johal A, Fleming P.S. (2019). Development of "My Retainers" mobile application: Triangulation of two qualitative methods. *Journal of Medical Internet Research Preprints*. Submitted for publication: 10.2196/preprints.14687.
13. Tsomos, G., Ludwig, B., Grossen, J., Pazera, P., Gkantidis, N. (2014). Objective assessment of patient compliance with removable orthodontic appliances: a cross-sectional cohort study. *The Angle Orthodontist*. 84, 56-61.
14. Little, R.M. (1975). The irregularity index: a quantitative score of mandibular anterior alignment. *American Journal of Orthodontics*. 68, 554-563.
15. Sullivan, T.R., White, I.R., Salter, A.B., Ryan, P. and Lee, K.J., (2018). Should multiple imputation be the method of choice for handling missing data in randomized trials?. *Statistical Methods in Medical Research*, 27,2610-2626.
16. Sterne, J.A., White, I.R., Carlin, J.B., Spratt, M., Royston, P., Kenward, M.G., Wood, A.M., Carpenter, J.R. (2009). Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *British Medical Journal*, 338, b2393.
17. Verbeke, G., Molenberghs, G. (2000). *Linear Mixed Models for Longitudinal Data*. Springer Series in Statistics. 336-338.
18. R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <https://www.r-project.org/> [Accessed July 4th, 2019].
19. Webb, T., Joseph, J., Yardley, L., Michie, S. (2010). Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *Journal of Medical Internet Research*. 12, e4.
20. Thomson, A.M., Cunningham, S.J., Hunt, N.P. (2001). A comparison of information retention at an initial orthodontic consultation. *European Journal of Orthodontics*. 23, 169-178.
21. Kang, E.Y., Fields, H.W., Kiyak, A., Beck, F.M. and Firestone, A.R. (2009). Informed consent recall and comprehension in orthodontics: traditional vs improved readability

- 1  
2  
3 and processability methods. *American Journal of Orthodontics and Dentofacial*  
4 *Orthopedics*. 136, 488.e1-488.e13.  
5  
6  
7 22. Al-Silwadi, F.M., Gill, D.S., Petrie, A., Cunningham, S.J. (2015). Effect of social media  
8 in improving knowledge among patients having fixed appliance orthodontic treatment:  
9 A single-center randomized controlled trial. *American Journal of Orthodontics and*  
10 *Dentofacial Orthopedics*. 148, 231-237.  
11  
12  
13 23. Chan, Y., Molassiotis, A. (1999). The relationship between diabetes knowledge and  
14 compliance among Chinese with non-insulin dependent diabetes mellitus in Hong  
15 Kong. *Journal of Advanced Nursing*. 30, 431-438.  
16  
17  
18 24. O'Cathain, A., Thomas, K., Drabble, S., Rudolph, A., Hewison, J. (2013). What can  
19 qualitative research do for randomised controlled trials? A systematic mapping review.  
20 *BMJ Open*. 3, e002889.  
21  
22  
23 25. Brierley, C.A., Benson, P.E., Sandler, J. (2017). How accurate are TheraMon®  
24 microsensors at measuring intraoral wear-time? Recorded vs. actual wear times in  
25 five volunteers. *Journal of Orthodontics*. 44, 241-248.  
26  
27  
28 26. Goldenberg, M.S. (2016). A randomized clinical trial on the effect of a compliance  
29 calendar on the use of maxillary Hawley retainers. Master's Thesis, Saint Louis  
30 University. Proquest Dissertations and Theses Global Database.  
31  
32  
33 27. Clemmer, E.J., Hayes, E.W. (1979). Patient cooperation in wearing orthodontic  
34 headgear. *American Journal of Orthodontics*. 75, 517-524.  
35  
36  
37 28. Bartsch, A., Witt, E., Sahm, G., Schneider, S. (1993). Correlates of objective patient  
38 compliance with removable appliance wear. *American Journal of Orthodontics and*  
39 *Dentofacial Orthopedics*. 104, 378-386.  
40  
41  
42 29. Schott, T.C., Ludwig, B. (2014). Microelectronic wear-time documentation of  
43 removable orthodontic devices detects heterogeneous wear behavior and  
44 individualizes treatment planning. *American Journal of Orthodontics and Dentofacial*  
45 *Orthopedics*. 146, 155-160.  
46  
47  
48 30. Huanca Ghislazoni, L., Ameer, S., Antonarakis, G.S., Kiliaridis, S. (2019). Headgear  
49 compliance as assessed by a temperature-sensitive recording device: a prospective  
50 clinical study. *European Journal of Orthodontics*. Epub ahead of print:  
51 10.1093/ejo/cjz036.  
52  
53  
54 31. Charavet, C., Gall, M.L., Albert, A., Bruwier, A., Leroy, S. (2019). Patient compliance  
55 and orthodontic treatment efficacy of Planas functional appliances with TheraMon  
56 microsensors. *The Angle Orthodontist*. 89, 117-122.  
57  
58  
59  
60



- 1  
2  
3  
4 32. Samuelson, B. (2017). Beam® Study in Orthodontic Patients. Master's Thesis, The  
5 University of Alabama at Birmingham. Proquest Dissertations and Theses Global  
6 Database.  
7
- 8  
9 33. Hichens, L., Rowland, H., Williams, A., Hollinghurst, S., Ewings, P., Clark, S., *et al.*  
10 (2007). Cost-effectiveness and patient satisfaction: Hawley and vacuum-formed  
11 retainers. *European Journal of Orthodontics*. 29, 372-378.  
12
- 13  
14 34. Türköz, Ç., Bavbek, N.C., Varlik, S.K., Akça, G. (2012). Influence of thermoplastic  
15 retainers on Streptococcus mutans and Lactobacillus adhesion. *American Journal of*  
16 *Orthodontics and Dentofacial Orthopedics*. 141, 598-603.  
17
- 18  
19 35. Valiathan, M., Hughes, E. (2010). Results of a survey-based study to identify common  
20 retention practices in the United States. *American Journal of Orthodontics and*  
21 *Dentofacial Orthopedics*. 137, 170-177.  
22
- 23  
24 36. Meade, M.J., Millett, D. (2013). Retention protocols and use of vacuum-formed  
25 retainers among specialist orthodontists. *Journal of Orthodontics*. 40, 318-325.  
26
- 27  
28 37. Manzon, L., Fratto, G., Rossi, E., Buccheri, A. (2018). Periodontal health and  
29 compliance: A comparison between Essix and Hawley retainers. *American Journal of*  
30 *Orthodontics and Dentofacial Orthopedics*. 153, 852-860.  
31
- 32  
33 38. Raja, T.A., Littlewood, S.J., Munyombwe, T. and Bubb, N.L., 2013. Wear resistance  
34 of four types of vacuum-formed retainer materials: a laboratory study. *The Angle*  
35 *Orthodontist*, 84, 656-664.  
36
- 37  
38 39. Attack, N., Harradine, N., Sandy, J.R., Ireland, A.J. (2007). Which way forward? Fixed  
39 or removable lower retainers. *The Angle Orthodontist*. 77, 954-959.  
40
- 41  
42 40. Littlewood, S.J., Millett, D.T., Doubleday, B., Bearn, D.R., Worthington, H.V. (2016).  
43 Retention procedures for stabilising tooth position after treatment with orthodontic  
44 braces. *Cochrane Database of Systematic Reviews*. 1, CD002283.  
45
- 46  
47 41. Parekh, J., Counihan, K., Fleming, P.S., Pandis, N., Sharma, P.K. (2019).  
48 Effectiveness of part-time vs full-time wear protocols of Twin-block appliance on  
49 dental and skeletal changes: A randomized controlled trial. *American Journal of*  
50 *Orthodontics and Dentofacial Orthopedics*. 155, 165-172.  
51
- 52  
53 42. Kearney, M.K., Pandis, N., Fleming, P.S. (2016). Mixed-methods assessment of  
54 perceptions of mandibular anterior malalignment and need for orthodontic  
55 retreatment. *American Journal of Orthodontics and Dentofacial Orthopedics*. 150,  
56 592-600.  
57
- 58  
59 43. Schäfer, K., Ludwig, B., Meyer-Gutknecht, H., Schott, T.C. (2015). Quantifying patient  
60 adherence during active orthodontic treatment with removable appliances using

1  
2  
3 microelectronic wear-time documentation. *European Journal of Orthodontics*. 37, 73-  
4 80.

- 5  
6 44. Koletsi, D., Fleming, P.S., Eliades, T., Pandis, N. (2015). The evidence from  
7 systematic reviews and meta-analyses published in orthodontic literature. Where do  
8 we stand? *European Journal of Orthodontics*. 37, 603-609.  
9

10  
11 **Figure legends:**

12  
13  
14 Figure 1. Retainer wear chart.

15  
16  
17 Figure 2. Screenshot of the calendar tool in the 'My Retainers' mobile  
18 application.  
19

20  
21  
22 Figure 3. CONSORT diagram showing the flow of participants.

23  
24 Mn: mandibular; Mx: maxillary.  
25  
26  
27  
28

29  
30 **Table captions:**

31  
32  
33 Table 1. Baseline characteristics of the sample (n= 84).  
34

35  
36 Table 2. Thermoplastic retainer failures during the study.  
37

38 Table 3. Data pertaining to retainer wear, stability and periodontal outcomes  
39 in both treatment groups. Data presented as median (interquartile range).  
40

41  
42 Table 4. Responses concerning experiences and levels of knowledge related  
43 to orthodontic retainers.  
44


45  
46 Table 5. Experience and knowledge outcomes in treatment groups (Exact  
47 Mann-Whitney test).  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Retainer wear chart

Please fill in hours of retainer wear

Study number .....



	Mon	Tues	Wed	Thurs	Fri	Sat	Sun
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Figure 1. Retainer wear chart.

339x471mm (96 x 96 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 2. Screenshot of the calendar tool in the 'My Retainers' mobile application




Figure 3. CONSORT diagram showing the flow of participants. Mn: mandibular; Mx: maxillary.

Table 1. Baseline characteristics of the sample (n= 84).

		Overall sample n= 84	Control group n= 42	Intervention group n= 42
<b>Mean age in years ± SD</b>		17.23 ± 1.9	17.20 ± 1.89	17.24 ± 2.00
<b>Gender</b>	Males	n= 42 (50%)	n= 21 (50%)	n= 21 (50%)
	Females	n= 42 (50%)	n= 21 (50%)	n= 21 (50%)
<b>Mean duration (years) of orthodontic treatment ± SD</b>		2.63 ± 0.86	2.72 ± 1.04	2.55 ± 0.64
<b>Treatment protocol</b>	Extraction	n= 51 (60.7%)	n= 29 (69%) (Mx only n= 7; Mn only n= 4; both arches n= 18)	n= 22 (52.4%) (Mx only n= 2; Mn only n= 3; both arches n= 17)
	Non-extraction	n= 33 (39.3%)	n= 13 (31%)	n= 20 (47.6%)
<b>Type of tooth-brush</b>	Manual	n= 60 (71.4%)	n= 30 (71.4%)	n= 30 (71.4%)
	Electric	n= 20 (23.8%)	n= 10 (23.8%)	n= 10 (23.7%)
	NI	n= 4 (4.8%)	n= 2 (4.8%)	n= 2 (4.8%)
<b>Daily tooth-brushing frequency</b>	Once	n= 11 (13.1%)	n= 6 (14.3%)	n= 5 (11.9%)
	Twice	n= 67 (79.8%)	n= 32 (76.2%)	n= 35 (83.3%)
	Three times	n= 2 (2.4%)	n= 2 (4.8%)	n= 0 (0%)
	NI	n= 4 (4.8%)	n= 2 (4.8%)	n= 2 (4.8%)
<b>Time spent tooth-brushing</b>	< 1 minute	n= 3 (3.6%)	n= 2 (4.8%)	n= 1 (2.4%)
	1-2 minutes	n= 56 (66.7%)	n= 29 (69%)	n= 27 (64.3%)
	> 2 minutes	n= 21 (25%)	n= 9 (21.4%)	n= 12 (28.6%)
	NI	n= 4 (4.8%)	n= 2 (4.8%)	n= 2 (4.8%)
<b>Use of other oral hygiene measures</b>	None	n= 45 (53.6%)	n= 20 (47.6%)	n= 25 (59.5%)
	Dental floss	n= 12 (14.3%)	n= 8 (19%)	n= 4 (9.5%)
	Interdental brush	n= 10 (11.9%)	n= 6 (14.3%)	n= 4 (9.5%)
	Toothpick	n= 13 (15.5%)	n= 6 (14.3%)	n= 7 (16.7%)
	NI	n= 4 (4.8%)	n= 2 (4.8%)	n= 2 (4.8%)
<b>Last visit to the dentist</b>	≤ 6 months	n= 18 (21.4%)	n= 9 (21.4%)	n= 9 (21.4%)
	> 6 months - 1 year	n= 15 (17.9%)	n= 8 (19%)	n= 7 (16.7%)
	> 1 year	n= 47 (56%)	n= 23 (54.8%)	n= 24 (57.1%)
	NI	n= 4 (4.8%)	n= 2 (4.8%)	n= 2 (4.8%)
<b>Smokers</b>		n= 4 (4.8%)	n= 2 (4.8%)	n= 2 (4.8%)

<b>Pregnancy</b>	n= 0 (0%)	n= 0 (0%)	n= 0 (0%)
------------------	-----------	-----------	-----------

Mn: mandibular; Mx: maxillary; NI: no information; SD: standard deviation.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 2. Thermoplastic retainer failures during the study.**

Reasons	Maxillary TPR	Mandibular TPR
Poor fit	n= 4	n= 5
Retainer loss	n= 2	n= 2
Breakage of the retainer	n= 7	n= 0
Detachment of the micro-electronic sensor	n= 2	n/a
Total	n= 15	n= 7

n/a: not applicable; TPR: thermoplastic retainer.



Table 3. Data pertaining to retainer wear, stability and periodontal outcomes in both treatment groups. Data presented as median (interquartile range).

Outcomes		Control group*	Intervention group	Coefficient†	95% CI	P-value	
Adherence levels	Objective data (h/d)	6.21 (7.86)	7.25 (6.71)	-0.91	-4.01, 2.19	0.56	
	Percentage of participants with ≥3 consecutive days of no retainer wear	57.6%	53.6%	-			
	Median percentage of days with wear as instructed (8 h/d and a minimum of 2 hours of continuous use)	46.67 (70.26)	55.70 (59.86)	-			
Stability outcomes	Maxilla	T0: 0.12 (0.1) T1: 0.14 (0.17)	T0: 0.16 (0.18) T1: 0.19 (0.22)	0.002	-0.03, 0.04	0.92	
	Mandible	T0: 0.16 (0.14) T1: 0.16 (0.21)	T0: 0.11 (0.12) T1: 0.16 (0.13)				
Periodontal outcomes	Plaque scores	Maxilla	T0: 0.84 (0.27) T1: 0.74 (0.22)	T0: 0.84 (0.18) T1: 0.75 (0.17)	-0.02	-0.07, 0.03	0.44
		Mandible	T0: 0.79 (0.25) T1: 0.76 (0.18)	T0: 0.84 (0.17) T1: 0.77 (0.17)			
	Bleeding on probing	Maxilla	T0: 0.17 (0.18) T1: 0.09 (0.1)	T0: 0.16 (0.17) T1: 0.08 (0.14)	-0.01	-0.05, 0.03	0.61
		Mandible	T0: 0.17 (0.18) T1: 0.1 (0.14)	T0: 0.20 (0.14) T1: 0.11 (0.1)			
	Probing depth	Maxilla	T0: 2.0 (0.18)	T0: 2.0 (0.25)	-0.01	-0.09, 0.07	0.79

			T1: 1.93 (0.24)	T1: 1.92 (0.31)			
		Mandible	T0: 1.7 (0.27)	T0: 1.8 (0.18)			
			T1: 1.62 (0.22)	T1: 1.6 (0.27)			

CI: confidence interval; h/d: hours/day.

\*Reference group

†Effect of treatment group on the outcome variables at T1.

**Table 4. Responses concerning experiences and levels of knowledge related to orthodontic retainers.**

Experiences						
Questions	Treatment group	1. Very dissatisfied	2. Dissatisfied	3. Neither satisfied nor dissatisfied	4. Satisfied	5. Very satisfied
Do you feel involved in the process of wearing and taking care of your retainers?	Control (n= 35)	0 (0%)	0 (0%)	3 (8.57%)	9 (25.71%)	23 (65.71%)
	Intervention (n= 29)	0 (0%)	0 (0%)	4 (13.79%)	6 (20.69%)	19 (65.52%)
How well do you feel you are being looked after since your braces were removed?	Control (n= 35)	0 (0%)	0 (0%)	2 (5.71%)	14 (40%)	19 (54.29%)
	Intervention (n= 29)	1 (3.45%)	0 (0%)	2 (6.9%)	5 (17.24%)	21 (72.41%)
How would you rate your overall experience within the last 3 months in terms of your use of retainers and contact with the clinic?	Control (n= 35)	0 (0%)	0 (0%)	5 (14.29%)	9 (25.71%)	21 (60%)
	Intervention (n= 29)	0 (0%)	2 (6.9%)	2 (6.9%)	11 (37.93%)	14 (48.28%)
Knowledge						
Questions	Treatment group	Percentage of correct responses				
If I wear the retainers really well for the first year, I can stop wearing them after that.	Control group (n= 35)	29/35 (82.86%)				
	Intervention group (n= 28)	25/28 (89.29%)				


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

How many hours a day do you need to wear the retainers?	Control group (n= 35)	21/35 (60%)
	Intervention group (n= 28)	19/28 (67.86%)
If you stopped wearing the retainers, what is likely to happen after a few weeks?	Control group (n= 35)	35/35 (100%)
	Intervention group (n= 28)	28/28 (100%)
How long do you need to wear your retainers for?	Control group (n= 35)	29/35 (82.86%)
	Intervention group (n= 28)	24/28 (85.71%)
What would you do if your retainers no longer fit or if you had problems with wearing them?	Control group (n= 35)	31/35 (88.57%)
	Intervention group (n= 28)	26/28 (92.86%)









Table 5. Experience and knowledge outcomes in treatment groups (Exact Mann-Whitney test).

Outcomes	Treatment group	Scores (median (IQR))	<i>P</i> -value
Patients' experiences (score out of 15)	Control group (n= 35)	14 (3)	0.94
	Intervention group (n= 29)	14 (2)	
Knowledge (score out of 5)	Control group (n= 35)	4 (1)	0.26
	Intervention group (n= 28)	5 (1)	

IQR: interquartile range.



**Supplementary material 1.** Laboratory procedures followed to integrate the TheraMon<sup>®</sup> micro-electronic sensor within the thermoplastic retainer

<p>(a) Each 13 x 9 x 4.5mm micro-electronic sensor (TheraMon<sup>®</sup>) was encapsulated individually or in groups in Essix ACE<sup>®</sup> plastic 1mm in thickness (DENTSPLY) using a Biostar<sup>®</sup> pressure thermoforming machine.</p>	
<p>(b) The Essix ACE<sup>®</sup> Plastic was then trimmed leaving a 2-3mm margin around the TheraMon<sup>®</sup> micro-electronic sensor.</p>	
<p>(c) The posterior buccal aspect of the maxillary working model was covered by a thin layer of plaster to flatten the surface.</p>	
<p>(d) A pre-heated (180°C, 70 seconds) sheet of Essix ACE<sup>®</sup> plastic 1mm in thickness, (DENTSPLY) was then pressure-formed on the working model at 6 bar of air pressure for 180 seconds using Biostar<sup>®</sup> pressure thermoforming machine.</p>	
<p>(e) The Essix ACE<sup>®</sup> Plastic was then trimmed around the base of the working model. Prior to bonding of the encapsulated micro-electronic sensor, the area was isolated using baseplate wax (Anutex<sup>®</sup> Toughened Dental Modelling Wax; Kemdent)</p>	
<p>(f) The micro-electronic sensor was then seated and bonded using auto-polymerising dental acrylic resin (Forestacryl<sup>®</sup>, Forestadent, Buckinghamshire, UK). The baseplate wax helped to prevent seepage of the auto-polymerising dental acrylic resin onto the TPR.</p>	
<p>(g) The auto-polymerising dental acrylic resin was set after placing the working model, TPR and attached micro-electronic sensor in a pressure-curing vessel. The wax was then removed.</p>	
<p>(h) The TPR was then removed from the working model, followed by trimming and polishing.</p>	

**Figure.** Laboratory procedures followed to integrate the TheraMon<sup>®</sup> micro-electronic sensor within the thermoplastic retainer.

TPR: thermoplastic retainer.

**Supplementary material 2.** Questionnaire to assess knowledge and experiences related to retainer wear



**Evaluation of tailored electronic reminders on compliance with removable orthodontic retention: a randomised controlled trial**

Chief Investigator: Dr Padhraig Fleming  
 Reader/Honorary Consultant in Orthodontics  
 Principal researcher: Dr Dalya Al Moghrabi  
 PhD student  
 Barts Health NHS Trust  
 Phone: 020 7882 8629






Version 4 (13/12/2018)  
 IRAS Id (201263)

Study number: .....

**Please fill in the questionnaire below:**

On a scale from 1 to 5 what would you rate the following statements?

(1: very poorly/ not at all satisfied      5: very well/ very satisfied)

Questions	1. 	2. 	3. 	4. 	5. 
1. How well do you understand the reasons for wearing the retainers?					
2. What do you think of the frequency of your follow-up appointments?					
3. Do you feel involved in the process of wearing and taking care of your retainers?					
4. Do you know where to seek advice regarding your retainers?					
5. Do you feel informed about the importance of retainers?					
6. How confident are you that your teeth won't move if you wear the retainers as advised?					
7. How satisfied are you that your questions about retainers are answered?					
8. How well do you remember to wear your retainers?					
9. How well do you feel you are being looked after since your braces were removed?					
10. How well do you remember where to store your retainers?					
11. How would you rate your overall experience within the last 3 months in terms of your use of retainers and contact with the clinic?					



**Supplementary material 2.** Questionnaire to assess knowledge and experiences related to retainer wear

---

1  
2  
3  
4  
5 12. If I wear the retainers really well for the first year, I can stop wearing them after that.

6  True

7  
8  False

9  
10  
11 13. How many hours a day do you need to wear the retainers?

12  12 hours

13  15 hours

14  20 hours

15  22 hours

16  
17  
18 14. If you stopped wearing the retainers, what is likely to happen after a few weeks?

19  Nothing, my teeth are stable especially if I wore the retainers really well previously

20  Gradual changes over time

21  
22 15. How long do you need to wear your retainers for?

23  6 months

24  1 year

25  Long-term

26  
27  
28 16. What would you do if your retainers no longer fit or if you had problems with wearing them?

29  Come for a casualty appointment

30  Wait for the next appointment

31  Stop wearing the retainers