

QUANTUM GRAVITY AND RIEMANNIAN GEOMETRY ON THE FUZZY SPHERE

EVELYN LIRA-TORRES AND SHAHN MAJID

ABSTRACT. We study the quantum geometry of the fuzzy sphere defined as the angular momentum algebra $[x_i, x_j] = 2i\lambda_p \epsilon_{ijk} x_k$ modulo setting $\sum_i x_i^2$ to a constant, using a recently introduced 3D rotationally invariant differential structure. Metrics are given by symmetric 3×3 matrices g and we show that for each metric there is a unique quantum Levi-Civita connection with constant coefficients, with scalar curvature $\frac{1}{2}(\text{Tr}(g^2) - \frac{1}{2}\text{Tr}(g)^2)/\det(g)$. As an application, we construct Euclidean quantum gravity on the fuzzy unit sphere. We also calculate the charge 1 monopole for the 3D differential structure.

1. INTRODUCTION

The angular momentum algebra $U(su_2)$ has been viewed since the 1970s as the quantisation of \mathbb{R}^3 viewed as su_2^* with its Kirillov-Kostant bracket as part of a general theory for any Lie algebra. As such, setting the quadratic Casimir to a constant quantises the coadjoint orbits, again in a standard way. The angular momentum algebra was also proposed as ‘position coordinates’ for Euclideanised 2+1 quantum gravity by ‘t Hooft[15]. We denote it $\mathbb{C}_\lambda[\mathbb{R}^3]$ or ‘fuzzy \mathbb{R}^3 ’ with generators x_i and relations $[x_i, x_j] = 2\lambda \epsilon_{ijk} x_k$ as in [4, 9, 18, 13] to indicate that we consider it a deformation of flat spacetime. Its covariance under the quantum double $D(U(su_2)) = U(su_2) \bowtie \mathbb{C}[SU_2]$ as ‘Poincaré group’ was found in [4] along with a 4D quantum-Poincaré invariant calculus, and further studied in [18][13] among other places. This is by now a well-established picture of a deformed \mathbb{R}^3 in Euclideanised 2+1 quantum gravity with point sources and without cosmological constant, see e.g. [12] at the group algebra level. Moreover, it deforms naturally to the quantum enveloping algebra $U_q(su_2)$ with quantum Poincaré group $D(U_q(su_2)) \cong \mathbb{C}_q[SO_{1,3}]$ in 2+1 quantum gravity with cosmological constant, see [26] for an overview and the relationship to the bicrossproduct model Majid-Ruegg quantum spacetime[25].

In physical terms, $\lambda = i\lambda_p$ where λ_p is a real deformation parameter which, in the above context (but not necessarily), should be of order the Planck scale. The general idea that spacetime geometry is ‘quantum’ or noncommutative was speculated since the 1920s but in modern times was proposed in [17] coming out of ideas for quantum gravity of quantum Born reciprocity or observable-state/Hopf algebra duality. See also subsequent works by many authors, including [1]. Not surprisingly, however, the quantum spacetime $\mathbb{C}_\lambda[\mathbb{R}^3]$ with its natural rotationally invariant quantum metric is flat and admits only the zero Levi-Civita connection in the standard coordinates [9, Example 8.15]. The same is true for the corresponding bicrossproduct model spacetime with quantum Poincaré covariant calculus[9, Prop. 9.20], the two models being related by twisting[23]. For quantum gravity effects with background gravity present we should look at curved quantum spacetimes.

Key words and phrases. Quantum gravity, fuzzy sphere, quantum geometry, fuzzy monopole, noncommutative geometry, angular momentum algebra, coadjoint quantisation.

Ver. 2. The first author was partially supported by CONACyT and Fundación Alberto y Dolores Andrade (México).

Indeed, one might expect such quantum Riemannian geometry to be more interesting for the corresponding ‘fuzzy sphere’ quotient, but it turns out that the differential structure proposed in [4] does not descend to the fuzzy sphere. To address this problem, [9, Example 1.46] recently proposed a different 3D differential structure on the fuzzy sphere and in the present paper we explore its quantum Riemannian geometry with this calculus. We find that it is indeed curved for general metrics, including its natural rotationally invariant ‘round metric’. We will denote the unit fuzzy sphere here by $\mathbb{C}_\lambda[S^2]$. The term ‘fuzzy sphere’ is also used in the literature, e.g. [16], for matrix algebras $M_n(\mathbb{C})$ viewed in our terms as further quotients of $\mathbb{C}_\lambda[S^2]$ for certain values of λ (those values for which the irreducible n -dimensional representations of spin $n/2$ descend).

We use the constructive ‘quantum groups’ approach to quantum Riemannian geometry as in the recent text [9]. This was established in recent years e.g. [5, 6, 8, 19, 21, 22, 24, 2] using particularly (but not only) the notion of a bimodule connection [11, 27]. The formalism is recalled briefly in Section 2 along with the new differential structure on $\mathbb{C}_\lambda[S^2]$ proposed in [9]. Section 3 contains the first new results, namely uniqueness and construction of a quantum-Levi Civita connection for each metric. Metrics here can be chosen freely as symmetric 3×3 matrices in the natural basis of the 3D calculus. In Section 3.2 we look at the curvature as a function of the metric and use this in Section 4 to explore Euclidean quantum gravity on the fuzzy sphere. Conventionally, the Euclidean case, although not usual quantum gravity itself, is nevertheless of interest on any compact Riemannian manifold with boundary [14].

Note also that our approach to quantum Riemannian geometry is very different from that of Connes [10] based on spectral triples as abstract ‘Dirac’ operators, but the two approaches can sometimes be related [7]. A first step for the fuzzy sphere would be to compute the Grassmann or monopole connection, which we do in the short Section 5.

It remains to explain why the cotangent bundle on the fuzzy sphere in this paper is 3D not 2D. Indeed, it often happens in quantum geometry that there is an obstruction to having the same dimension connected differential calculus as classically and preserving symmetries. This is because most highly noncommutative geometries are inner in the sense of a 1-form θ such that the exterior derivative is $d = [\theta, \]$ but this equation has no meaning in classical geometry (the right hand side would be zero) and indeed it is a purely quantum phenomenon. Hence quantum calculi often contain an extra dimension not visible classically, which we could think of as an internally generated ‘time’ direction $\theta = dt$, since quite often the partial derivative in the θ direction turns out to be the natural wave operator or Laplacian. This was explained in [18] in the context of the 4D calculus on $\mathbb{C}_\lambda[\mathbb{R}^3]$, where this external time is not part of spacetime but may be more related to geodesic flow proper time according to recent ideas in [8] or to renormalisation group flow according to ideas in [13]. In our context, for the new calculus on $\mathbb{C}_\lambda[S^2]$ in [9] which we use, there is a more direct geometric interpretation of this extra dimension: there is in fact a (nonconnected) 3D calculus on $\mathbb{C}_\lambda[\mathbb{R}^3]$ with no extra dimension needed there, but when this descends to the sphere quotient, the normal direction does not decouple as it would classically and instead one has $\theta = (2i\lambda_p)^{-2} x_i dx_i$. So one should think of the fuzzy sphere with this calculus as embedded in a fuzzy \mathbb{R}^3 and retaining the normal vector as an additional direction. This then makes its calculus 3D and not 2D.

2. PRELIMINARIES

Here we give a very short introduction to the general formalism, with more details to be found in [9] and references there in. This provides the framework whereby our constructions for the fuzzy sphere should not be seen as ad-hoc but natural within this context. To this end, let A be a unital algebra, possibly noncommutative, over a field \mathbb{K} (we will be mainly interested in \mathbb{C}). By a first order *differential calculus* (Ω^1, d) we mean that:

- (1) Ω^1 is an A -bimodule;
- (2) A linear map $d : A \rightarrow \Omega^1$ such $d(ab) = (da)b + adb$ for all $a, b \in A$;
- (3) $\Omega^1 = \text{span}\{adb \mid a, b \in A\}$;
- (4) (*Optional*) $\ker(d) = \mathbb{K}.1_A$

Here d is called the *exterior derivative*, the condition (3) is the surjectivity condition and (4) is the *connectedness property*, which is not an axiom but is desirable. We also require Ω^1 to extend to an exterior algebra $\Omega = \oplus_i \Omega^i$ of forms of different degree, generated by $A = \Omega^0$ and Ω^1 with d extending by the graded Leibniz rule and $d^2 = 0$. The product of Ω is denoted \wedge . For quantum Riemannian geometry we only need up to Ω^2 . See [9, Chap. 1].

In this context, we define a quantum metric as $g \in \Omega^1 \otimes_A \Omega^1$ such that there exists an inverse $(\ , \) : \Omega^1 \otimes_A \Omega^1 \rightarrow A$ which is a bimodule map. Inverse here means in the usual sense, explicitly $(\omega, g^1)g^2 = \omega = g^1(g^2, \omega)$ for all ω in a notation where $g = g^1 \otimes g^2$ (sum of terms and tensor product over A understood). This implies [9] that g is central, since $ag = g^1(g^2, ag^1) \otimes g^2 = g^1 \otimes (g^2 a, g^1)g^2 = ga$ for all a , where $g = g^1 \otimes g^2$ is another copy of g . We usually (but not always) also require g to be quantum symmetric in the sense $\wedge(g) = 0$. If this does not hold, we speak of an asymmetric or ‘generalised’ metric.

Next, a left connection on Ω^1 for us means a linear map $\nabla : \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$ obeying the left Leibniz rule

$$\nabla(a.\omega) = da \otimes \omega + a.\nabla\omega$$

for all $a \in A$ and $\omega \in \Omega^1$. If $X : \Omega^1 \rightarrow A$ is a right vector field in the sense that it commutes with the action of A from the right, then we may define $\nabla_X = (X \otimes \text{id})\nabla : \Omega^1 \rightarrow \Omega^1$ as an associated covariant derivative-like map. It obeys $\nabla_X(a\omega) = (X(da))\omega + a\nabla_X\omega$ as expected if X also commutes with the action of A from the left. In the classical case with local coordinates x^i , say, we would set $\nabla dx^i = -\Gamma^i_{jk} dx^j \otimes dx^k$ in terms of Christoffel symbols. One can apply a similar definition for any vector bundle in the sense of a left A -module E (typically required to be projective). In our case $E = \Omega^1$ is a bimodule and we demand

$$\nabla(\eta a) = (\nabla\eta)a + \sigma(\eta \otimes da),$$

for a bimodule map $\sigma : \Omega^1 \otimes_A \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$, called the *generalised braiding*. If this exists, it is uniquely determined by this formula, so this is not additional data, just a property of some left connections. We say that ∇ is then a *bimodule connection* [11, 27]. This case is nice because bimodule connections can be tensor producted. Relevant to us is that $\Omega^1 \otimes_A \Omega^1$ gets a bimodule connection

$$\nabla(\omega \otimes \eta) = \nabla\omega \otimes \eta + (\sigma(\omega \otimes ()) \otimes \text{id})\nabla\eta$$

for $\omega, \eta \in \Omega^1$. In this case $\nabla g = 0$ makes sense and we say when that holds that g is *metric compatible* [6]. Explicitly, we need

$$\nabla g := (\nabla \otimes \text{id})g + (\sigma \otimes \text{id})(\text{id} \otimes \nabla)g = 0.$$

See [9, Chap 8] for more details.

Also, for any left connection on Ω^1 and a choice of Ω^2 , we have a canonical notion of torsion, which is the standard notion but written in terms of differential forms as

$$T_\nabla = \wedge \nabla - d : \Omega^1 \rightarrow \Omega^2$$

Given a generalised metric we also have a notion of ‘cotorsion’ define by

$$coT_\nabla = dg^1 \otimes g^2 - g^1 \wedge \nabla g^2 \in \Omega^2 \otimes_A \Omega^1,$$

which classically is a skew-symmetrized version of metric compatibility. We say that a connection is a *quantum Levi-Civita connection* (QLC) for a metric g if it is torsion free and metric compatible. We say that it is a weak QLC (WQLC) if it is cotorsion free and torsion free. One can show that a QLC is necessarily a WQLC so it can be useful to impose the WQLC condition first, being linear in ∇ compared to the QLC condition which is quadratic (due to the σ , which is linear in ∇) and hence much harder to solve.

Finally, over \mathbb{C} , we need everything to be ‘unitary’ or ‘real’ in a suitable sense. It means that A is a $*$ -algebra in the usual sense, and that $*$ extends to Ω in a way that commutes with d and is a graded-order reversing involution (it means there is an extra minus sign on a product of odd degree forms). We require the metric and connection to be ‘real’ in the sense

$$\text{flip}(* \otimes *)g = g, \quad \sigma \circ \text{flip}(* \otimes *)\nabla = \nabla \circ *.$$

The key here is the map $\dagger = \text{flip}(* \otimes *) : \Omega^1 \otimes_A \Omega^1 \rightarrow \Omega^1 \otimes_A \Omega^1$ defined by $(\omega \otimes \eta)^\dagger = \eta^* \otimes \omega^*$, which one can check is well-defined on the tensor product over A understood here, and obeys $(a\omega \otimes \eta b)^\dagger = b^*(\omega \otimes \eta)^\dagger a^*$. It is shown in [9] that $(\sigma \circ \dagger)^2 = \text{id}$ then holds. Just imposing the latter is a weaker notion of $*$ -compatibility as in [5]. In the classical case with self-adjoint local coordinates, the displayed ‘reality’ conditions would ensure that the metric and connection coefficients are real. These are a well-studied set of axioms for which many interesting examples are known, e.g., [6, 21, 22, 24]. Also note that if the reality property for ∇ holds on ω then

$$\sigma((\nabla(a\omega))^\dagger) = \sigma(\omega^* \otimes da^* + (\nabla\omega)^\dagger a^*) = \sigma(\omega^* \otimes da^*) + (\nabla\omega^*)a^* = \nabla(\omega^* a^*) = \nabla((a\omega)^*)$$

using both Leibniz rules, so it also holds on $a\omega$. It follows that it suffices to prove it on a generating set of 1-forms.

2.1. Fuzzy sphere and its 3D differential calculus. We work over \mathbb{C} and start with the enveloping algebra $U(su_2)$ of the angular momentum Lie algebra, with basis x_i normalised so that

$$[x_i, x_j] = 2i\lambda_p \epsilon_{ijk} x_k$$

for a parameter λ_p . We call this $\mathbb{C}_\lambda[\mathbb{R}^3]$ as a quantisation of functions on \mathbb{R}^3 . We take it as a $*$ algebra with $x_i^* = x_i$ and λ_p real. Note that this has finite-dimensional irreducible representations ρ_j labelled by a non-negative half-integer j of dimension $n = 2j + 1$ and in which $\sum_i x_i^2 = (n^2 - 1)\lambda_p^2$ in our normalisation. We define the *unit fuzzy sphere* $A = \mathbb{C}_\lambda[S^2]$ as the quotient $U(su_2)$ modulo the relation

$$\sum_i x_i^2 = 1 - \lambda_p^2$$

which we see descends to the spin j representation precisely when $\lambda_p = 1/n$. We keep λ_p as a free parameter, however. Note that in all cases $A = \mathbb{C}_\lambda[S^2]$ is infinite-dimensional and therefore never a matrix algebra.

Next we define $\Omega(\mathbb{C}_\lambda[\mathbb{R}^3])$ as a free 3D calculus with central basis s^i , $i = 1, 2, 3$. This means we impose $[s^i, x_j] = 0$, and we define a differential

$$dx_i = \epsilon_{ijk} x_j s^k$$

which one can check is translation and rotation invariant calculus, but not connected. Indeed, $d \sum_i x_i^2 = 0$ so there are different connected components according to any constant value of $\sum_i x_i^2$. The calculus is inner with

$$\theta = \frac{1}{2i\lambda_p} x_i s^i = \frac{1}{(2i\lambda_p)^2} x_i dx_i = -\frac{1}{(2i\lambda_p)^2} (dx_i) x_i.$$

Finally, for the exterior algebra we take s^i to be the Grassmann algebra, with

$$s^i \wedge s^j + s^j \wedge s^i = 0, \quad ds^i = -\frac{1}{2} \epsilon_{ijk} s^j \wedge s^k$$

and Ω free over this (so in each degree it has a basis over A given by a basis over \mathbb{C} of the Grassmann algebra). The reader should be warned, however, that this is no longer inner by θ in higher degree. That in turn means it is not the maximal prolongation of the first order calculus, but is a natural quotient.

We then take the same form of calculus and d for $A = \mathbb{C}_\lambda[S^2]$, where we add the unit sphere relation. This is compatible with d for reasons already given and this time we obtain a connected calculus. These facts are all covered in [9, Example 1.46] and one could do the same for a sphere of any fixed radius.

Lemma 2.1. *In $\Omega(\mathbb{C}_\lambda[S^2])$, one has*

$$s^l = \frac{1}{(1-\lambda_p^2)} (2i\lambda_p x_l \theta + \epsilon_{lim} (dx_i) x_m) = \frac{1}{(1-\lambda_p^2)} \left(\frac{1}{2i\lambda_p} x_l x_i dx_i + \epsilon_{lim} (dx_i) x_m \right).$$

Proof. First observe from the form of dx_i that $\epsilon_{lim} dx_i = \epsilon_{lim} \epsilon_{ijk} x_j s^k = x_l s^m - x_m s^l$. Then

$$\begin{aligned} \epsilon_{lim} (dx_i) x_m &= (x_l s^m - x_m s^l) x_m = x_l x_m s^m - x_m x_m s^l = 2i\lambda_p x_l \theta - x_m x_m s^l \\ &= 2i\lambda_p x_l \left(\frac{1}{(2i\lambda_p)^2} x_i dx_i \right) - x_m x_m s^l = \frac{1}{2i\lambda_p} x_l x_i dx_i - x_m x_m s^l \end{aligned}$$

using the two forms of θ . We then use the sphere relation and rearrange as stated. \square

From this we see that s^l does not necessarily have a classical analogue when $\lambda_p \rightarrow 0$, due to the $\frac{1}{2i\lambda_p} x_i dx_i$ in the second expression. However, if we take this limit with θ fixed then $x_i dx_i$ is order λ_p^2 and this ‘purely quantum’ term in s^l then vanishes in the limit. In this sense, we think of s^l as a deformation of the Killing forms

$$s_{\text{class}}^l = \epsilon_{lim} (dx_i) x_m$$

for the action of the rotation group on the classical sphere with its standard round metric. These are the standard orbital angular momentum vector fields on the sphere when viewed via the metric as 1-forms [9, Cor 9.10].

3. MODULI OF QLCs ON THE FUZZY SPHERE

Once we have fixed the calculus it is clear, since $\Omega^1(\mathbb{C}_\lambda[S^2])$ has a central basis, a general metric, as already observed in [9, Example 1.46], takes the form

$$g = g_{ij} s^i \otimes s^j$$

where, since g has to be central, we need the coefficients g_{ij} to be central. Since the centre of $U(su_2)$ is generated by the quadratic Casimir, it follows that $\mathbb{C}_\lambda[S^2]$ has trivial centre, so $g_{ij} \in \mathbb{C}$. For quantum symmetry, we need $0 = g_{ij} s^i \wedge s^j = \sum_{i < j} (g_{ij} s^i \wedge s^j + g_{ji} s^j \wedge s^i) = \sum_{i < j} (g_{ij} - g_{ji}) s^i \wedge s^j$, i.e. g_{ij} symmetric. We assume such symmetry throughout, for simplicity. For the reality property, we note that $s^{i*} = s^i$ so we need g_{ij} to be hermitian which, given the symmetry, means $g_{ij} \in \mathbb{R}$. Finally, we need g_{ij} to be an invertible matrix with inverse g^{ij} , say. Then $(s^i, s^j) = g^{ij}$. The new question, which we address, is what are the QLCs and WQLCs. Part of this was answered in [30], although not our main result Proposition 3.4.

3.1. Quantum Levi-Civita connection. First, we consider an arbitrary connection equivalently in the form

$$\nabla s^i = -\frac{1}{2}\Gamma^i_{jk}s^j \otimes s^k,$$

where $\Gamma^i_{jk} \in A$. As s^i form a basis, this extends by the Leibniz rule to a left connection.

Proposition 3.1. *Let $\Gamma_{ijk} := g_{im}\Gamma^m_{jk}$.*

- (1) ∇ is torsion free if and only if $\Gamma_{ijk} - \Gamma_{ikj} = 2g_{im}\epsilon_{mjk}$.
- (2) ∇ is cotorsion free if and only if $\Gamma_{ijk} - \Gamma_{jik} = 2g_{km}\epsilon_{mij}$.

Proof. (1) is immediate from

$$T_\nabla s^i = \wedge \nabla s^i - ds^i = \wedge (-\frac{1}{2}\Gamma^i_{kl}s^k \otimes s^l) + \frac{1}{2}\epsilon_{ikl}s^k \wedge s^l$$

that torsion freeness needs $\Gamma^i_{kl} - \Gamma^i_{lk} + 2\epsilon_{ilk} = 0$, which we write to match (2) in terms of the lowered index version.

(2) We calculate

$$\begin{aligned} coT_\nabla &= d(g_{ij}s^i) \otimes s^j - g_{ij}s^i \wedge (-\frac{1}{2}\Gamma^j_{kl}s^k \otimes s^l) \\ &= g_{ij}(-\frac{1}{2}\epsilon_{imn}s^m \wedge s^n) \otimes s^j + \frac{1}{2}g_{ij}s^i \wedge (\Gamma^j_{kl}s^k \otimes s^l) \\ &= -\frac{1}{2}(g_{ml}\epsilon_{mik} - g_{ij}\Gamma^j_{kl})(s^i \wedge s^k) \otimes s^l. \end{aligned}$$

So vanishing of cotorsion requires that $g_{ml}\epsilon_{mik} - g_{ij}\Gamma^j_{kl}$ is symmetric in i, k by a similar argument as for symmetry of the metric, which we have written as stated. \square

Therefore, the moduli of WQLCs (where torsion and cotorsion vanish) are given by the two conditions (1) and (2) simultaneously. Note that the two conditions have a very similar form, which is part of the symmetric role of torsion and cotorsion in the WQLC theory. Finally, for the full QLC theory, we need to know when ∇ is a bimodule connection, i.e. when there exists a suitable generalised braiding σ and what it looks like.

Lemma 3.2. *If ∇ is a bimodule connection on $\Omega^1(\mathbb{C}_\lambda[S^2])$ then*

$$\sigma(s^i \otimes s^j) = s^j \otimes s^i + \frac{1}{2(1-\lambda_p^2)}\left(\frac{1}{2i\lambda_p}x_jx_n[\Gamma^i_{lk}, x_n] + \epsilon_{jmn}[\Gamma^i_{lk}, x_m]x_n\right)s^l \otimes s^k$$

Proof. The generalised braiding if it exists, given that $[s^i, x_j] = 0$, is characterised by

$$\sigma(s^i \otimes dx_j) = dx_j \otimes s^i - [\nabla s^i, x_j] = \epsilon_{jmn}x_ms^n \otimes s^i + \frac{1}{2}[\Gamma^i_{mn}, x_j]s^m \otimes s^n,$$

where the tensor is over the algebra A . Now substituting s^n from Lemma 2.1,

$$\begin{aligned} \sigma(s^i \otimes s^n) &= \sigma(s^i \otimes \frac{1}{2i\lambda_p(1-\lambda_p^2)}x_nx_jdx_j + \frac{1}{1-\lambda_p^2}\epsilon_{nmj}(dx_m)x_j) \\ &= \frac{1}{2i\lambda_p(1-\lambda_p^2)}x_nx_j\sigma(s^i \otimes dx_j) + \frac{1}{1-\lambda_p^2}\epsilon_{nmj}\sigma(s^i \otimes dx_m)x_j \\ &= \frac{1}{2i\lambda_p(1-\lambda_p^2)}x_nx_j(\epsilon_{jmk}x_ms^k \otimes s^i + \frac{1}{2}[\Gamma^i_{mk}, x_j]s^m \otimes s^k) \\ &\quad + \frac{1}{1-\lambda_p^2}\epsilon_{nmj}(\epsilon_{mlk}x_ls^k \otimes s^i + \frac{1}{2}[\Gamma^i_{lk}, x_m]s^l \otimes s^k)x_j \\ &= \frac{1}{1-\lambda_p^2}\left(\frac{1}{2i\lambda_p}\epsilon_{jmk}x_nx_jx_m + \epsilon_{nmj}\epsilon_{mlk}x_lx_j\right)s^k \otimes s^i + \\ &\quad + \frac{1}{2(1-\lambda_p^2)}\left(\frac{1}{2i\lambda_p}x_nx_j[\Gamma^i_{lk}, x_j] + \epsilon_{nmj}[\Gamma^i_{lk}, x_m]x_j\right)s^l \otimes s^k \\ &= s^n \otimes s^i + \frac{1}{2(1-\lambda_p^2)}\left(\frac{1}{2i\lambda_p}x_nx_j[\Gamma^i_{lk}, x_j] + \epsilon_{nmj}[\Gamma^i_{lk}, x_m]x_j\right)s^l \otimes s^k \end{aligned}$$

as stated, using the commutation relations in A . \square

It follows that σ has to flip on the s^i basic forms if the $\Gamma^i_{jk} \in A$ are central, i.e. $\Gamma_{ijk} \in \mathbb{C}$ as our algebra has trivial centre and g already has constant coefficients. In this constant coefficients case, $\sigma(s^i \otimes s^j) = s^j \otimes s^i$ has a well-defined extension as a bimodule map since the s^i are a central basis of Ω^1 , and ∇ is indeed then a bimodule connection. Also in this case, the ‘reality’ property with respect to $*$ reduces to $\Gamma_{ijk} \in \mathbb{R}$. We are now ready to consider the condition for full metric compatibility under this assumption.

Lemma 3.3. [30] *The metric compatibility condition assuming Γ are constant coefficients is*

$$\Gamma_{lik} + \Gamma_{kil} = 0.$$

Proof.

$$\begin{aligned} \nabla g &= (\nabla \otimes \text{id})(g_{ij}s^i \otimes s^j) + (\sigma \otimes \text{id})(g_{ij}s^i \otimes \nabla(s^j)) \\ &= -g_{ij} \frac{1}{2} \Gamma^i_{mn} s^m \otimes s^n \otimes s^j - (\sigma \otimes \text{id})(g_{ij}s^i \otimes \frac{1}{2} \Gamma^j_{mn} s^m \otimes s^n) \\ &= -g_{ij} \frac{1}{2} \Gamma^i_{mn} s^m \otimes s^n \otimes s^j - \frac{1}{2} g_{ij} \Gamma^j_{mn} \sigma(s^i \otimes s^m) \otimes s^n \\ &= \left(-\frac{1}{2} (\Gamma_{nlk} + \Gamma_{kln}) - \frac{1}{4(1-\lambda_p^2)} \Gamma_{imn} \left(\frac{x_m x_p}{2i\lambda_p} [\Gamma^i_{lk}, x_p] + \epsilon_{mpq} [\Gamma^i_{lk}, x_p] x_q \right) \right) s^l \otimes s^k \otimes s^n. \end{aligned}$$

In the natural case of constant coefficients of Γ , we can drop the second term. \square

It is a nice check that torsion free and metric compatible (in our constant Γ case) implies cotorsion free, as it must. Indeed, we can write torsion freeness as $\Gamma_{lki} - \Gamma_{lik} = 2g_{lm}\epsilon_{mki}$ by Proposition 3.1. Given metric compatibility in the form just found, this is equivalent to $-\Gamma_{ikl} + \Gamma_{kil} = 2g_{lm}\epsilon_{mki}$, which is the cotorsion free condition in Proposition 3.1. It remains to solve for the moduli of constant coefficient QLC solutions for a given a metric $g_{ij}s^i \otimes s^j$.

Proposition 3.4. *For any metric g_{ij} , there is a unique QLC among those with constant coefficients, namely*

$$\Gamma_{ijk} = 2\epsilon_{ikm}g_{mj} + \text{Tr}(g)\epsilon_{ijk}.$$

There are real, hence the connection is $$ -preserving.*

Proof. We have to solve the joint system

$$\Gamma_{ikl} - \Gamma_{ilk} = 2g_{im}\epsilon_{mkl}, \quad \Gamma_{lik} + \Gamma_{kil} = 0,$$

the second of which is solved by setting $\Gamma_{ijk} = \epsilon_{ikm}\gamma_{mj}$ for some matrix γ . Letting L_i be the matrices $(L_i)_{mn} = \epsilon_{imn}$, the first equation is then

$$L_i \gamma - (L_i \gamma)^t + 2g_{im} L_m = 0$$

as matrices. This is a linear system for γ with a unique solution

$$\gamma = 2g - \text{Tr}(g)\text{id}$$

which translates into the solution stated. (Note that we do not have to solve the cotorsion equation $\Gamma_{kil} - \Gamma_{ikl} + 2g_{lm}\epsilon_{mki} = 0$ as this is implied, as mentioned.) \square

For example, when $g_{ij} = \delta_{ij}$ (the rotationally invariant or ‘round’) metric, we have a unique solution $\Gamma_{ijk} = \epsilon_{ijk}$. The classical limit of this metric with θ fixed is $g_{\text{class}} = dx_i \otimes dx_i$, which is the classical round metric on the unit sphere in this description[9, Sec. 9.3.1].

3.2. Ricci curvature. Now that we understand the moduli of QLCs, we explore their curvature on the fuzzy sphere for general metrics. The curvature in quantum Riemannian geometry is

$$R_{\nabla} : \Omega^1 \rightarrow \Omega^2 \otimes_A \Omega^1, \quad R_{\nabla} = (d \otimes \text{id} - \text{id} \wedge \nabla) \nabla$$

which one can check is well-defined and a left A -module map [9, Lemma 8.1]. In our case, we can necessarily write it in the form

$$R_{\nabla}(s^i) = \rho^i_{jk} \epsilon_{jmn} s^m \wedge s^n \otimes s^k$$

for some curvature coefficients $\rho^i_{jk} \in A$. We are also interested in taking a ‘trace’ for the Ricci tensor and the Ricci scalar, which in the current framework [9] means with respect to a further, but in our case canonical, bimodule ‘lift’ map

$$i : \Omega^2 \rightarrow \Omega^1 \otimes_A \Omega^1, \quad i(s^i \wedge s^k) = \frac{1}{2}(s^i \otimes s^k - s^k \otimes s^i).$$

For this choice of i and form of R_{∇} , we have

$$\begin{aligned} \text{Ricci} &= ((,) \otimes \text{id})(\text{id} \otimes i \otimes \text{id})(\text{id} \otimes R_{\nabla})(g) \\ &= \frac{1}{2} g_{pi} \rho^i_{jk} \epsilon_{jmn} ((s_p,) \otimes \text{id})(s^m \otimes s^n \otimes s^k - s^n \otimes s^m \otimes s^k) \\ &= \frac{1}{2} \rho^i_{jk} \epsilon_{jin} s^n \otimes s^k - \frac{1}{2} \rho^i_{jk} \epsilon_{jmi} s^m \otimes s^k = \rho^i_{jn} \epsilon_{jim} s^m \otimes s^n. \end{aligned}$$

Hence the Ricci tensor defined by $\text{Ricci} = R_{mn} s^m \otimes s^n$ and Ricci scalar $S = (,) \text{Ricci}$ are

$$R_{mn} = \rho^i_{jn} \epsilon_{jim}, \quad S = \rho^i_{jn} \epsilon_{jim} g^{mn}.$$

Finally, we adopt the convention that indices of ϵ can be raised with the inverse metric $(,) = g^{-1}$ with matrix entries g^{ij} .

Proposition 3.5. For $\mathbb{C}_{\lambda}[S^2]$, the scalar curvature for the QLC in Proposition 3.4 is

$$S = \frac{1}{2}(\text{Tr}(g^2) - \frac{1}{2}\text{Tr}(g)^2) / \det(g).$$

Proof. We first compute from its definition as given above that

$$\begin{aligned} R_{\nabla}(s^i) &= -\frac{1}{2}(\Gamma^i_{kl} ds^k + (d\Gamma^i_{kl})s^k) \otimes s^l + \frac{1}{2}\Gamma^i_{kl} s^k \wedge \nabla s^l \\ &= \frac{1}{4}\Gamma^i_{kl} \epsilon_{kmn} s^m \wedge s^n \otimes s^l - \frac{1}{2}d\Gamma^i_{kl} \wedge s^k \otimes s^l - \frac{1}{4}\Gamma^i_{kl} \Gamma^l_{mn} s^k \wedge s^m \otimes s^n \end{aligned}$$

which corresponds to

$$(1) \quad \rho^i_{jk} = \frac{1}{4}\Gamma^i_{jk} - \frac{1}{4}\epsilon_{jmn} \partial_m \Gamma^i_{nk} - \frac{1}{8}\epsilon_{jmn} \Gamma^i_{ml} \Gamma^l_{nk}$$

where $df = (\partial_i f) s^i$ defines the partial derivatives. One can check that this reproduces the same R_{∇} . This applies for any left connection on Ω^1 .

We now specialise to the QLC in Proposition 3.4. Then

$$\rho^i_{jk} = \frac{1}{2}(\epsilon^i_{km} g_{mj} - \epsilon^i_{lp} \epsilon_{jmn} \epsilon^l_{kq} g_{pm} g_{qn}) + \frac{1}{4}\text{Tr}(g)(\epsilon^i_{jk} - \epsilon^j_{ik} + \epsilon^{ij}_{pg} g_{pk}) - \frac{1}{8}\text{Tr}(g)^2 \epsilon^{ij}_k$$

which we contract to obtain Ricci as

$$R_{st} = \frac{1}{2}\epsilon^i_{tm} g_{mj} \epsilon_{jis} + \frac{1}{2}\text{Tr}(g^{-1}) g_{st} - \frac{1}{2}\delta_{st} + \frac{1}{2}\text{Tr}(g)(g^{st} - \text{Tr}(g^{-1})\delta_{st} - \frac{1}{2}\epsilon^{ij}_p \epsilon_{ijs} g_{pt}) + \frac{1}{8}\text{Tr}(g)^2 \epsilon^{ij}_t \epsilon_{ijs}.$$

This then contracts further to

$$S = \text{Tr}(g^{-1}) + \frac{1}{2}\epsilon^{ij}_m (g_{mk} - \frac{1}{2}\text{Tr}(g)\delta_{mk} + \frac{1}{4}\text{Tr}(g)^2 g^{mk}) \epsilon_{ijk} + \frac{1}{2}\text{Tr}(g)(\text{Tr}(g^{-2}) - \text{Tr}(g^{-1})^2).$$

Finally, we identify the middle $\epsilon \dots \epsilon$ expression in terms of $\det g^{-1}$. This is most easily seen assuming that $g = \text{diag}(\lambda_1, \lambda_2, \lambda_3)$, say, but then holds generally. Here $\epsilon^{ij}_k g^{mk} \epsilon_{ijk} = \epsilon^2_{ijk} \lambda_i^{-1} \lambda_j^{-1} \lambda_k^{-1} = 6 \det g^{-1}$ (summing over i, j, k) is well-known, but a similar method gives

$$\begin{aligned} \epsilon^{ij}_k \epsilon_{ijk} &= \epsilon^2_{ijk} \lambda_i^{-1} \lambda_j^{-1} \epsilon^2_{ijk} \lambda_i^{-1} \lambda_j^{-1} \lambda_k^{-1} \lambda_k = 2 \det g^{-1} \text{Tr}(g) \\ \epsilon^{ij}_m g_{mk} \epsilon_{ijk} &= \epsilon^2_{ijk} \lambda_i^{-1} \lambda_j^{-1} \lambda_k = \epsilon^2_{ijk} \lambda_i^{-1} \lambda_j^{-1} \lambda_k^{-1} \lambda_k^2 = 2 \det g^{-1} \text{Tr}(g^2). \end{aligned}$$

In this way, we obtain

$$S = \text{Tr}(g^{-1}) + (\text{Tr}(g^2) + \frac{1}{4}\text{Tr}(g)^2) \det g^{-1} + \frac{1}{2}\text{Tr}(g) (\text{Tr}(g^{-2}) - \text{Tr}(g^{-1})^2)$$

which then simplifies further to the form stated, as one can again check in the diagonal case. In terms of the λ_i , this is

$$(2) \quad S = \frac{\lambda_1^2 + \lambda_2^2 + \lambda_3^2 - 2(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3)}{4\lambda_1\lambda_2\lambda_3},$$

which one can also regard as a function on the space of metrics modulo conjugation if we think of the λ_i as the (not necessarily distinct) eigenvalues of g . \square

Note that the classical limit of S in the noncommutative geometry conventions here is $-\frac{1}{2}$ of the classical value. Hence a classical unit sphere with its usual round metric would in our conventions have $S = -1$. In the fuzzy case, for the rotationally invariant ‘round’ metric $g_{ij} = \delta_{ij}$ on $\mathbb{C}_\lambda[S^2]$ and the unique QLC $\Gamma_{ijk} = \epsilon_{ijk}$ in Proposition 3.4, we have

$$R_{mn} = -\frac{1}{4}\delta_{mn}, \quad S = -\frac{3}{4}.$$

This is not quite the same as the classical value, presumably as a consequence of the extra cotangent direction. If we perturb around this metric by setting $g = \text{id} + \epsilon$ then we can write

$$S = -\frac{3}{4} + \frac{1}{4}\text{Tr}\epsilon - \frac{1}{12}(\text{Tr}\epsilon)^2 + \frac{1}{4}(\epsilon_{12}^2 + \epsilon_{13}^2 + \epsilon_{23}^2) + \frac{1}{24}((\epsilon_{11} - \epsilon_{22})^2 + (\epsilon_{11} - \epsilon_{33})^2 + (\epsilon_{22} - \epsilon_{33})^2) + O(\epsilon^3),$$

showing an unbounded mode for the average of the diagonal entries plus a positive definite part for the ‘fluctuations’ off diagonal or between the diagonal entries.

4. EUCLIDEANISED QUANTUM GRAVITY ON THE FUZZY SPHERE

The scalar curvature found above is the main, but not the only, ingredient for quantum gravity in a functional integral approach. Here we briefly consider the other elements and formulate the theory, although it will remain too hard to compute explicitly. Moreover, this will not be physical gravity since our spherical form of coordinates where $\sum_i x_i^2 = 1 - \lambda_p^2$ are suitable for a Euclidean signature (the actual ‘shape’ depends on the metric but the choice of algebra plays the role of the manifold in some sense) but this is still of interest in certain contexts. In the Euclidean case, the partition function for the functional integral should take the form

$$Z = \int \mathcal{D}g e^{-\frac{2}{G} \int S[g]}$$

(and similarly with operators inserted for expectation values), where G is some real positive coupling constant and

$$(3) \quad S[g] = \frac{g_{11}^2 + g_{22}^2 + g_{33}^2 - 2(g_{11}g_{22} + g_{11}g_{33} + g_{22}g_{33}) + 4(g_{12}^2 + g_{13}^2 + g_{23}^2)}{4 \det g}$$

is the scalar curvature in Proposition 3.5.

We also need for the action a map $\int : A \rightarrow \mathbb{C}$ which classically would be the Lebesgue measure in local spacetime coordinates times $\sqrt{|\det(g)|}$. One could attempt to characterise this map in the quantum case by requiring that it is a positive linear functional (so $\int a^*a \geq 0$ for all a in the algebra) compatible with the Riemannian metric in some way (classically this would be so as to vanish on a total divergence). This is an interesting question but is not needed for our present purposes as $S[g] \in \mathbb{R}$ is a multiple of the constant function, so the only thing we need is $\int 1$. The natural choice here similar to integration on a Riemannian manifold would be power of $|\det g|$ (but not necessarily its square root as classically, given

that our tangent space and metric have the wrong dimension compared to classical sphere). Bearing in mind the $\det(g)$ in the denominator of $S[g]$, the most natural choice here is to cancel this by setting

$$\int 1 = |\det(g)|.$$

Finally, we need a measure $\mathcal{D}g$ for integration over the space of metrics. In our case, this is the 6-dimensional space \mathcal{P}_3 of 3×3 positive-definite symmetric matrices, which has the natural structure of a Riemannian manifold as the noncompact symmetric space $GL_3(\mathbb{R})/O_3(\mathbb{R})$, with an invariant metric $\mathfrak{g}_{\mathcal{P}_3}$ given in line element terms as

$$ds^2 = \text{Tr}((g^{-1}dg)^2).$$

Integration over \mathcal{P}_3 therefore has a canonical Riemannian manifold measure $\mathcal{D}g$ defined relative to Lebesgue measure in local coordinates by $\sqrt{|\det(\mathfrak{g}_{\mathcal{P}_3})|}$. The latter works out at metric g as a factor $|\det(g)|^{-2}$, see [31, Sec 4.1.3].

Putting these ingredients together, we thus define Euclideanised quantum gravity on the fuzzy sphere by partition function

$$(4) \quad Z = \int_{\mathcal{P}_3} \prod_{i \leq j} dg_{ij} |\det(g)|^{-2} e^{-\frac{1}{G}(\text{Tr}(g^2) - \frac{1}{2}\text{Tr}(g)^2)},$$

where G is a real coupling constant. Here expectation values are the ratio of the same expression with operators inserted divided by Z , both parts of which can be expected to diverge given the above initial remarks and the noncompact nature of \mathcal{P}_3 . One can also write

$$\det(g)^{-2} = (2\pi)^{-\frac{3}{2}} \int_{\mathbb{R}^3} d^3\bar{x} e^{-\frac{1}{2}\bar{x}^t g^2 \bar{x}}$$

so that up to a discarded constant and assuming we can swap the order of integration,

$$(5) \quad Z = \int_{\mathbb{R}^3} \prod_i dx_i \int_{\mathcal{P}_3} \prod_{i \leq j} dg_{ij} e^{-\frac{1}{G}(\text{Tr}(g^2) - \frac{1}{2}\text{Tr}(g)^2) - \frac{1}{2}\bar{x}^t g^2 \bar{x}},$$

giving an idea of the formal content of the theory. Moreover, *if* we ignored the restriction to \mathcal{P}_3 , we could then do the dg integration as a Gaussian to give the inverse determinant of a quadratic form built from the x_i . We can also write $(\det g)^{-2} = e^{-2\text{Tr} \ln g}$ in (4) as a non-quadratic ‘interaction’ term. We now look more closely at the theory in a reduced form where we look only at SO_3 -invariant expressions.

Our first step is to parameterize positive symmetric matrices according to the spectral decomposition $g = C^t \text{diag}(\lambda_1, \lambda_2, \lambda_3)C$ for some $\vec{\lambda} \in \mathbb{R}_{>0}^3$ and some $C \in SO_3$. The latter is not unique but the multiplicity is discrete and generically we can convert dg to these new coordinates with an appropriate Jacobean. To do this explicitly, we let $E(\theta, \phi, \psi)$ be the Euler rotation matrix for angles θ, ϕ, ψ and let

$$g = E(\theta, \phi, \psi)^t \text{diag}(\lambda_1, \lambda_2, \lambda_3)E(\theta, \phi, \psi).$$

This change of variables is locally invertible for $\sin \phi \neq 0$ and distinct λ_i , indeed the Jacobean can be computed and we find

$$\prod_{i \leq j} dg_{ij} = d\theta d\phi d\psi |\sin(\phi)| \prod_i d\lambda_i |(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_3)|$$

On the other hand, the action itself does not depend on the SO_3 conjugation (having the value in (2) but without the $\lambda_1 \lambda_2 \lambda_3$ denominator there), so for the partition function and for any insertions that depend only on the λ_i and not on the angles,




FIGURE 1. Expectation values for Euclidean quantum gravity on the fuzzy sphere as a function of the cutoff L . The right graph shows $\langle \lambda_i \lambda_j \rangle$ converging to $\frac{16}{3} \langle \lambda_i \rangle^2$ and $\frac{3}{16} L^2$.

we can do the integration over a dense subset of SO_3 to give a constant, which we ignore. Hence (4) becomes effectively

$$(6) \quad Z = \int_{\epsilon}^L \prod_i d\lambda_i \frac{|(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_3)|}{\lambda_1^2 \lambda_2^2 \lambda_3^2} e^{-\frac{1}{2G}(\lambda_1^2 + \lambda_2^2 + \lambda_3^2 - 2(\lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \lambda_2 \lambda_3))},$$

where we have introduced cut-offs $L \gg \epsilon > 0$ to regulate divergences at both ends. The story here turns out to be very similar to [21] in that the divergence as $\epsilon \rightarrow 0$ does not show up when we look at vacuum expectation values as these are ratios (both top and bottom diverge as $\epsilon \rightarrow 0$ but the ratio is well defined in this limit). Moreover,

$$\langle \lambda_{i_1} \cdots \lambda_{i_n} \rangle \sim \frac{3}{16} L^n$$

for large L , independently of which λ_i are involved (this was checked numerically to several orders of λ), as plotted for $n = 1, 2$ in Figure 1. It follows that we have well-defined ratios

$$\frac{\langle \lambda_{i_1} \cdots \lambda_{i_n} \rangle}{\langle \lambda_i \rangle^n} = \left(\frac{16}{3}\right)^{n-1}$$

in the limit $L \rightarrow \infty$, and in particular that there is a uniform relative uncertainty in the λ_i ,

$$\frac{\Delta \lambda_i}{\langle \lambda_i \rangle} := \frac{\sqrt{\langle \lambda_i^2 \rangle - \langle \lambda_i \rangle^2}}{\langle \lambda_i \rangle} = \sqrt{\frac{13}{3}}$$

similarly to quantum gravity on a square in [21]. Note, however, that these are formal interpretations given that this is a Euclidean theory.

We can also follow the pattern of [21] and look at a partial theory where we regard the average of the λ_i as a background metric with respect to which we are quantising only the differences. Thus, we let

$$\lambda_1 = u - 2v, \quad \lambda_2 = u + v - w, \quad \lambda_3 = u + v + w$$

with inverse

$$u = \frac{1}{3}(\lambda_1 + \lambda_2 + \lambda_3), \quad v = \frac{1}{6}(\lambda_2 + \lambda_3 - 2\lambda_1), \quad w = \frac{1}{2}(\lambda_3 - \lambda_2),$$

which diagonalizes the quadratic form in the action to

$$\lambda_1^2 + \lambda_2^2 + \lambda_3^2 - 2(\lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \lambda_2 \lambda_3) = -3u^2 + 12v^2 + 4w^2.$$

In the partial theory, we leave out the du integral and regard $u > 0$ as a parameter. Putting in the region of u, v, w corresponding to $\lambda_i > 0$, we have the effective theory

for the ‘fluctuation’ variables v, w ,

$$\begin{aligned} Z_u &= 2 \int_{-u}^{\frac{u}{2}} dv \int_{-(u+v)}^{u+v} dw \frac{|(9v^2 - w^2)w|}{(u-2v)^2((u+v)^2 - w^2)^2} e^{-\frac{2}{G}(3v^2+w^2)} \\ &= 4 \int_{-u}^{\frac{u}{2}} dv \int_0^{u+v} dw \frac{|9v^2 - w^2|w}{(u-2v)^2((u+v)^2 - w^2)^2} e^{-\frac{2}{G}(3v^2+w^2)}. \end{aligned}$$

This is still divergent at the boundaries corresponding previously to $\lambda_i = 0$, but has the merit that the inner integral can now be done analytically. The previous partition function

$$Z = \int_0^\infty du e^{\frac{3}{2G}u^2} Z_u$$

still contains the other divergence at $\lambda_i = \infty$, requiring a cut-off. One can make this change for the diagonal entries of any metric and have the off-diagonals as three further Gaussian variables according to (3), but the restriction on the variables for a positive metric is then much harder to describe.

5. FUZZY MONPOLE

We have focussed on the quantum Riemannian geometry of the fuzzy sphere with its new calculus from [9]. However, we could also ask about the monopole connection for this calculus. Classically, this arises naturally on the tautological line bundle over the sphere which in algebraic terms is the Grassmann connection for the rank 1 projector associated with that.

We recall that if A is a unital algebra, a vector bundle can be viewed as, say, a left finitely generated projective module \mathcal{S} . This property is characterised by existence of a finite number n of generators $e^\alpha \in \mathcal{S}$ and corresponding left module maps $e_\beta : \mathcal{S} \rightarrow A$ such that $s = e_\alpha(s)e^\alpha$ for all $s \in \mathcal{S}$. There is an associated projection $P_{\alpha\beta} = e_\beta(e^\alpha)$ obeying $P^2 = P$ as an element of $M_n(A)$ and expressing relations $e^\alpha = P_{\alpha\beta}e^\beta$ in \mathcal{S} . Equivalently, given a projection P we can realise \mathcal{S} concretely as generated by the rows e^α of P and set $e_\beta(e^\alpha) = P_{\alpha\beta}$ extended as left-module maps. The associated Grassmann connection and its curvature in the conventions of [9, Prop. 3.24] are

$$\nabla_{\mathcal{S}} e^\alpha = (dP_{\alpha\beta})P_{\beta\gamma} \otimes e^\gamma, \quad R_{\mathcal{S}}(e^\alpha) = -dP_{\alpha\beta} \wedge (dP_{\beta\gamma})P_{\gamma\delta} \otimes e^\delta.$$

The construction is best-known in the case of the universal calculus, where it underlies the Chern-Connes pairing between K-theory and cyclic cohomology[10].

It was already shown in [20] that the fuzzy sphere (as well as the standard q -sphere and the 2-parameter q -fuzzy sphere) has such a projection matrix $P_{\alpha\beta}$ associated to a tautological rank 1 projective module \mathcal{S} . Here $P \in M_2(A)$ and [9, Example 3.27] covers the Grassmann connection but depending on the choice of calculus; here we carry this through for our choice of $\Omega(\mathbb{C}_\lambda[S^2])$. In terms of natural ‘coordinates’ and relations

$$x = \frac{1}{2}(x_3 + 1 + \lambda_p), \quad z = \frac{1}{2}(x_1 + ix_2); \quad [x, z] = \lambda_p z, \quad [z, z^*] = 2\lambda_p x - \lambda_p(1 + \lambda_p), \quad z^* z = x(1-x),$$

the projection matrix is [20]

$$P = \begin{pmatrix} 1 + \lambda_p - x & z \\ z^* & x \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 + \lambda_p - x_3 & x_1 + ix_2 \\ x_1 - ix_2 & 1 + \lambda_p + x_3 \end{pmatrix}$$

which one can check obeys $P^2 = P$. Starting with P , one can realise \mathcal{S} as spanned by the rows of P with one relation,

$$e^1 = (1, 0).P = (1 + \lambda_p - x, z), \quad e^2 = (0, 1).P = (z^*, x); \quad (x - \lambda_p)e^1 = ze^2$$

and forming dual bases with e_β defined by $e_\beta(e^\alpha) = P_{\alpha\beta}$ as explained. Here, we will stick with the x_i generators of A in order to match the rest of the paper.

Lemma 5.1. *The monopole Grassmann connection on \mathcal{S} for our projection P and calculus on $\mathbb{C}_\lambda[S^2]$ is*

$$\nabla e^\alpha = \left(\frac{1 + \lambda_p}{2} dP_{\alpha\beta} + \lambda_p P_{\alpha\beta} \theta + \iota \frac{1 - \lambda_p^2}{4} Q_{\alpha\beta} - \frac{\lambda_p(1 - \lambda_p)}{2} \delta_{\alpha\beta} \theta \right) \otimes e^\beta,$$

$$\text{where } Q = \begin{pmatrix} -s^3 & s^1 + \iota s^2 \\ s^1 - \iota s^2 & s^3 \end{pmatrix}.$$

Proof. Here $\nabla_{\mathcal{S}} : \mathcal{S} \rightarrow \Omega^1 \otimes_A \mathcal{S}$ obeys similar axioms to those for a left connection on Ω^1 in Section 2. The calculation of the stated formula from the projector is a straightforward from

$$dP = \frac{1}{2} \begin{pmatrix} x_2 s^1 - x_1 s^2 & x_2 s^3 - x_3 s^2 + \iota x_3 s^1 - \iota x_1 s^3 \\ x_2 s^3 - x_3 s^2 - \iota x_3 s^1 + \iota x_1 s^3 & x_1 s^2 - x_2 s^1 \end{pmatrix}$$

and the commutation relations in the algebra. For example,

$$\begin{aligned} ((dP)P)_{11} &= \frac{1}{4} \left((x_2 s^1 - x_1 s^2)(1 + \lambda_p - x_3) + (x_2 s^3 - x_3 s^2 - \iota x_3 s^1 + \iota x_1 s^3)(x_1 - \iota x_2) \right) \\ &= \frac{1}{4} \left((1 + \lambda_p)(x_2 s^1 - x_1 s^2) + \iota x_3(x_1 s^1 + x_2 s^2 + x_3 s^3) - \iota(1 - \lambda_p^2) s^3 + \right. \\ &\quad \left. + [x_3, x_2] s^1 + [x_1, x_3] s^2 + [x_2, x_1] s^3 \right) \\ &= \frac{1}{4} \left((1 + \lambda_p)(x_2 s^1 - x_1 s^2) - 2x_3 \lambda_p \theta + \iota(\lambda_p^2 - 1) s^3 - 2\iota \lambda_p(x_1 s^1 + x_2 s^2 + x_3 s^3) \right) \\ &= \frac{1}{4} \left((1 + \lambda_p) d(-x_3) + \iota(\lambda_p^2 - 1) s^3 - 2\lambda_p x_3 \theta + 4\lambda_p^2 \theta \right) \\ &= \frac{1}{2} (1 + \lambda_p) dP_{11} + \lambda_p P_{11} \theta - \frac{1}{2} (1 - \lambda_p) \lambda_p \theta - \frac{\iota}{4} (1 - \lambda_p^2) s^3. \end{aligned}$$

Similar for the other entries of $dP.P$, the proof of which we omit. \square

The curvature of ∇ was also recalled in general as $-dP \wedge (dP)P$ acting as a 2-form valued operation on our e^α .

Proposition 5.2. *The curvature of the monopole connection is given by*

$$-dP \wedge (dP)P = \frac{(1 - \lambda_p)}{2\iota} \left(\begin{pmatrix} x_1 & \lambda_p \\ \lambda_p & x_1 \end{pmatrix} s^2 \wedge s^3 + \begin{pmatrix} x_2 & \iota \lambda_p \\ -\iota \lambda_p & x_2 \end{pmatrix} s^3 \wedge s^1 + \begin{pmatrix} x_3 - \lambda_p & 0 \\ 0 & x_3 + \lambda_p \end{pmatrix} s^1 \wedge s^2 \right) P.$$

Proof. We similarly compute the matrix-valued coefficients of $s^1 \wedge s^2, s^2 \wedge s^3, s^3 \wedge s^1$ of $dP \wedge (dP)P$, using the commutation relations in the differential algebra. For example,

$$\begin{aligned} 8(dP \wedge (dP)P)_{11} &= 8(dP_{11} \wedge dP_{11} + dP_{12} \wedge dP_{21})P_{11} + 8(dP_{11} \wedge dP_{12} + dP_{12} \wedge dP_{22})P_{21} \\ &= ((x_2 s^1 - x_1 s^2)(x_2 s^1 - x_1 s^2) + (x_2 s^3 - x_3 s^2 + \iota x_3 s^1 - \iota x_1 s^3)(x_2 s^3 - x_3 s^2 + \iota x_1 s^3 - \iota x_3 s^2)) \\ &\quad (1 + \lambda_p - x_3) \\ &\quad + ((x_2 s^1 - x_1 s^2)(x_2 s^3 - x_3 s^2 + \iota x_3 s^1 - \iota x_1 s^3) + (x_2 s^3 - x_3 s^2 + \iota x_3 s^1 - \iota x_1 s^3)(x_1 s^2 - x_2 s^1)) \\ &\quad (x_1 - \iota x_2) \end{aligned}$$

and from this we pick out the coefficient of $s^1 \wedge s^2$ as

$$\begin{aligned} &([x_1, x_2] - 2\iota x_3^2)(1 + \lambda_p - x_3) + (-[x_2, x_3]x_2 - 2x_3 x_2 + \iota[x_1, x_3] + 2\iota x_3 x_1)(x_1 - \iota x_2) \\ &= (2\iota \lambda_p x_3 - 2\iota x_3^2)(1 + \lambda_p - x_3) + (-2\iota \lambda_p x_1 - 2x_3 x_2 + 2\lambda_p x_2 + 2\iota x_3 x_1)(x_1 - \iota x_2) \\ &= 2\iota(x_3 - \lambda_p)(x_3(-1 - \lambda_p + x_3) + (x_1 + \iota x_2)(x_1 - \iota x_2)) \\ &= 2\iota(1 - \lambda_p)(x_3 - \lambda_p)(1 + \lambda_p - x_3) \end{aligned}$$

using the commutation relations and the quadratic relation for the fuzzy sphere. By comparison, the coefficient of $s^1 \wedge s^2$ on the right hand side of the stated result is

$$\frac{(1 - \lambda_p)}{4\iota} \begin{pmatrix} (x_3 - \lambda_p)(1 + \lambda_p - x_3) & (x_3 - \lambda_p)(x_1 + \iota x_2) \\ (x_3 + \lambda_p)(x_1 - \iota x_2) & (x_3 + \lambda_p)(1 + \lambda_p + x_3) \end{pmatrix}$$

which thus verifies the top left corner entry. Similarly for the other matrix entries. One then repeats for the coefficient of $s^2 \wedge s^3$ and of $s^3 \wedge s^1$. Note that the factorisation into matrices in the stated answer is not unique as P has a kernel; we have chosen a factorisation which is much simpler than taking the matrix coefficients of $dP \wedge dP$ as the factor in front of P . \square

Note that the right hand side in the proposition reduces as $\lambda_p \rightarrow 0$ to FP , where

$$F = \frac{1}{2i}(x_1 s^2 \wedge s^3 + x_2 s^3 \wedge s^1 + x_3 s^1 \wedge s^2) = \frac{1}{4i} \epsilon_{ijk} x_i s^j \wedge s^k = \frac{1}{4i} \epsilon_{ijk} x_i dx^j \wedge dx^k$$

if we use the classical limit of the s^l with θ fixed as discussed after Lemma 2.1. This agrees with a classical monopole where the curvature is an imaginary multiple of the volume form. The last expression is indeed a multiple of the classical volume form by rotational invariance.

6. CONCLUDING REMARKS

Fuzzy- \mathbb{R}^3 in the form of the angular momentum algebra $U(su_2)$ has a long history as a ‘quantisation’ as well as clear applications such as at the heart of 3D Euclideanised quantum gravity without cosmological constant. Its quotient the fuzzy sphere is likewise well known as the quantisation of a coadjoint orbit. Although less relevant perhaps to 3D quantum gravity, it is nevertheless related to Penrose’ spin network geometry[28], and could also have a role for the geometry of angular momentum in actual quantum systems. A differential calculus that works with it was, however, only recently proposed[9, Example 1.46] and it is significant that, using this, we have now solved its quantum Riemannian geometry (found a natural quantum Levi-Civita connection) for any quantum metric, including the canonical rotationally invariant ‘round’ metric given by δ_{ij} .

After understanding this moduli of quantum Riemannian geometries on the fuzzy sphere, we went on and constructed Euclidean quantum gravity on it, where we integrate over all quantum geometries. Even though the fuzzy sphere is an infinite-dimensional algebra which becomes functions (in some form) on the usual sphere in the classical limit, its quantum geometry turned out to be much more rigid and to admit only quantum metrics of the form of a single 3×3 real symmetric matrix g_{ij} transported over the whole algebra. It turned out that the quantum Levi-Civita connection, hence the whole moduli of quantum Riemannian geometries could similarly be developed with constant coefficients, hence the quantum geometry behaves effectively like just one point. This was not put in by hand, but forced by the rigidity of the axioms of noncommutative geometry and the noncommutativity of our particular algebra. Moreover, our results were strikingly similar to quantum gravity on a quadrilateral in [21] even though the details are completely different not to mention that that model was Lorentzian with ι in action, whereas ours is Euclidean. In both cases, the functional integral over all metrics of the natural action built from the quantum Ricci scalar has UV and IR divergences, appearing in our case at $\lambda_i = 0, \infty$. In both cases, the divergence at metric zero modes cancels in the ratio of functional integrals when we look at expectation values. In both cases, the other divergence is controlled by a cut-off L , and in both cases the expectation value of an n -th power of the field diverges as L^n with the result that ratios of expectation values can still be defined as $L \rightarrow \infty$. In both cases, we found in this way a uniform relative uncertainty in the quantisation of the metric components (in our case, we quantised the metric eigenvalues but one can also think of this as quantising diagonal metrics).

There are many interesting directions that one could further explore. On the quantum gravity front, one could introduce matter and see how some kind of Einstein equation emerges out of quantum gravity with matter. A first step here would

be to better understand the geometric approach to the stress-energy tensor. One can consider quantum matter fields on curved FLRW cosmologies $\mathbb{R} \times S^2$ where the S^2 is now fuzzy. Although the field theory stress tensor is not clear, the analysis for a perfect fluid is possible and will be done elsewhere in the spirit of [2], where $\mathbb{R} \times \mathbb{Z}_n$ was done using quantum geometry on the polygon \mathbb{Z}_n (this work also solved Euclidean quantum gravity on \mathbb{Z}_n). Finally, one could look at quantum geodesics on the fuzzy sphere using the recent formalism in [8].

We have also constructed a natural fuzzy monopole and there are potentially many applications that could be related to that. In physical terms, this could be relevant to any quantum system where classically one has angular momentum at play, for example the effective geometry around a quantised hydrogen atom. In mathematical terms, an important application would be towards the programme of ‘geometric realisation’ - constructing examples of Connes’ notion of spectral triple[10] or ‘axiomatic Dirac operator’ but in a geometric manner starting with the quantum differential structure, a spinor bundle with connection, a quantum metric and a Clifford structure. This was done for the q -sphere in [7] and one might try to follow the same steps. Thus, the spinor bundle on the sphere should be of the form $\mathcal{S} = \mathcal{S}_+ \oplus \mathcal{S}_-$ where \mathcal{S}_+ is the charge 1 monopole line bundle as found in Section 5 and \mathcal{S}_- is its dual. The Clifford structure is a ‘Clifford action’ map $\Omega^1 \otimes_A \mathcal{S} \rightarrow \mathcal{S}$ obeying certain axioms [7, 9] of compatibility with the $*$ -structure, with the connection on \mathcal{S} and with the quantum Levi-Civita connection. On the other hand, the Clifford structure for the q -sphere case in [7] was found from the holomorphic structure of its 2D calculus, which does not apply here. This nevertheless merits further study and will be attempted elsewhere. If a geometrically-realised spectral triple can be constructed on the fuzzy sphere, it may (or may not) descend when $\lambda_p = 1/n$ to the quotient $c_\lambda[S^2]$ isomorphic to M_n , i.e. to the reduced matrix fuzzy spheres. This may then complement (or perhaps relate to) the finite fuzzy Dirac operators constructed in [3] using Connes formalism and starting from the reduced noncommutative torus.

Conflict of interest. On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

- [1] G. Amelino-Camelia and S. Majid, Waves on noncommutative spacetime and gamma-ray bursts, *Int. J. Mod. Phys. A* 15 (2000) 4301–4323
- [2] J. Argota Quiroz and S. Majid, Quantum gravity on polygons and $\mathbb{R} \times \mathbb{Z}_n$ FLRW model, *Class. Quant. Gravity* 37 (2020) 245001 (43pp)
- [3] J. W. Barrett, Matrix geometries and fuzzy spaces as finite spectral triples, *J. Math. Phys.* 56 (2015) 082301
- [4] E. Batista and S. Majid, Noncommutative geometry of angular momentum space $U(su_2)$, *J. Math. Phys.* 44 (2003) 107–137
- [5] E.J. Beggs and S. Majid, $*$ -compatible connections in noncommutative Riemannian geometry, *J. Geom. Phys.* 61 (2011) 95–124
- [6] E.J. Beggs and S. Majid, Gravity induced by quantum spacetime, *Class. Quantum. Grav.* 31 (2014) 035020 (39pp)
- [7] E.J. Beggs and S. Majid, Spectral triples from bimodule connections and Chern connections, *J. Noncomm. Geom.*, 11 (2017) 669–701
- [8] E.J. Beggs and S. Majid, Quantum geodesics in quantum mechanics, arXiv:1912.13376 (math-ph)
- [9] E.J. Beggs and S. Majid, *Quantum Riemannian Geometry*, Grundlehren der mathematischen Wissenschaften, Vol. 355, Springer (2020) 809pp
- [10] A. Connes, *Noncommutative Geometry*, Academic Press, Inc., San Diego, CA, 1994
- [11] M. Dubois-Violette and P.W. Michor, Connections on central bimodules in noncommutative differential geometry, *J. Geom. Phys.* 20 (1996) 218–232

- [12] L. Freidel and E. R. Livine, Ponzano-Regge model revisited. III: Feynman diagrams and effective field theory, *Class. Quantum Grav.* 23 (2006) 2021
- [13] L. Freidel and S. Majid, Noncommutative harmonic analysis, sampling theory and the Duflo map in 2+1 quantum gravity, *Class. Quant. Gravity* 25 (2008) 045006 (37pp)
- [14] S. Hawking and G. Gibbon, eds. *Euclidean Quantum Gravity*, World Scientific (1993)
- [15] G. 't Hooft, Quantization of point particles in 2+1 dimensional gravity and space- time discreteness, *Class. Quant. Grav.* 13 (1996) 1023
- [16] J. Madore, *An introduction to noncommutative differential geometry and its physical applications*, LMS Lecture Note Series, 257, C.U.P. 1999.
- [17] S. Majid, Hopf algebras for physics at the Planck scale, *Class. Quantum Grav.* 5 (1988) 1587–1607
- [18] S. Majid, Noncommutative model with spontaneous time generation and Planckian bound, *J. Math. Phys.* 46 (2005) 103520 (18pp)
- [19] S. Majid, Noncommutative Riemannian geometry of graphs, *J. Geom. Phys.* 69 (2013) 74–93
- [20] S. Majid, q-Fuzzy spheres and quantum differentials on $B_q[SU_2]$ and $U_q(su_2)$, *Lett. Math. Phys.* 98 (2011) 167–191
- [21] S. Majid, Quantum gravity on a square graph, *Class. Quantum Grav* 36 (2019) 245009 (23pp)
- [22] S. Majid, Quantum Riemannian geometry and particle creation on the integer line, *Class. Quantum Grav.* 36 (2019) 135011 (22pp)
- [23] S. Majid and P.K. Osei, Quasitriangular structure and twisting of the 2+1 bicrossproduct model, *J. High Energ. Phys.* 1 (2018) 147 (28pp)
- [24] S. Majid and A. Pachol, Digital finite quantum Riemannian geometries, *J. Phys. A* 53 (2020) 115202 (40pp)
- [25] S. Majid and H. Ruegg, Bicrossproduct structure of the κ -Poincaré group and non-commutative geometry, *Phys. Lett. B.* 334 (1994) 348–354
- [26] S. Majid and B. Schroers, q-Deformation and semidualisation in 3D quantum gravity, *J. Phys A* 42 (2009) 425402 (40pp)
- [27] J. Mourad, Linear connections in noncommutative geometry, *Class. Quant. Grav.* 12 (1995) 965–974
- [28] R. Penrose, Angular momentum: an approach to combinatorial spacetime, in *Quantum Theory and Beyond*, T. Basin, ed., Cambridge University Press, Cambridge (1971)
- [29] H. S. Snyder, Quantized space-time, *Phys. Rev. D* 67 (1947) 38–41
- [30] I. Staden, Noncommutative differential structures, MSc thesis, QMUL (2019)
- [31] A. Terras, *Harmonic Analysis on Symmetric Spaces and Applications, Vol II*, Springer (2012)

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY UNIVERSITY OF LONDON, MILE END RD,
LONDON E1 4NS

E-mail address: `e.y.liratorres@qmul.ac.uk`, `s.majid@qmul.ac.uk`