

Supplementary file 1: Patient interval by ethnicity – first symptoms

Symptom	Total cohort				White men				Black men				Asian men			
	1-3 months	3-6 months	6-12 months	>12 months	1-3 months	3-6 months	6-12 months	>12 months	1-3 months	3-6 months	6-12 months	>12 months	1-3 months	3-6 months	6-12 months	>12 months
Any symptom	157 (57.3)	37 (13.5)	32 (11.7)	46 (16.8)	108 (54.8)	29 (14.7)	22 (11.2)	37 (18.8)	24 (58.5)	5 (12.2)	9 (21.9)	2 (4.9)	25 (69.4)	3 (8.3)	1 (2.8)	7 (19.4)
Hesitancy	25 (83.3)	1 (3.3)	2 (6.7)	2 (6.7)	22 (81.5)	1 (3.7)	2 (7.4)	2 (7.4)	2 (100)	0	0	0	1 (100)	0	0	0
Frequency	21 (55.3)	5 (13.6)	3 (7.9)	9 (23.7)	13 (56.5)	3 (13.0)	1 (4.4)	6 (26.1)	5 (55.6)	1 (11.1)	2 (22.2)	1 (11.1)	3 (50.0)	1 (16.7)	0	3 (33.3)
Urgency	6 (75.0)	1 (12.5)	0	1 (12.5)	4 (80.0)	0	0	1 (20.0)	1 (100)	0	0	0	1 (50.0)	1 (50.0)	0	0
Urgency-with-dribble	4 (66.7)	1 (16.7)	0	1 (16.7)	1 (100)	0	0	0	1 (50.0)	1 (50.0)	0	0	2 (66.7)	0	0	1 (33.3)
Nocturia	34 (40.9)	14 (16.9)	15 (18.1)	20 (24.1)	21 (32.8)	12 (18.8)	13 (20.3)	18 (28.1)	8 (66.7)	2 (16.7)	2 (16.7)	0	5 (71.4)	0	0	2 (28.6)
Retention	2 (66.7)	0	0	1 (33.3)	1 (50.0)	0	0	1 (50.0)	0	0	0	0	1 (100)	0	0	0
Haematuria	29 (93.6)	2 (6.5)	0	0	22 (91.7)	8.3	0	0	3 (100)	0	0	0	4 (100)	0	0	0
Erectile dysfunction	25 (42.4)	12 (20.3)	10 (16.7)	11 (18.6)	18 (41.7)	11 (25.6)	5 (11.6)	8 (18.6)	2 (28.6)	0	4 (57.1)	1 (14.3)	5 (55.7)	1 (11.1)	1 (11.1)	2 (22.2)
Other	11 (73.3)	1 (6.7)	1 (6.7)	2 (13.3)	6 (75.0)	0	0	2 (25.0)	2 (50.0)	1 (25.0)	1 (25.0)	0	3 (100)	0	0	0

Values are n (%)