

# A palliative care goals model for people with dementia and their family: Consensus achieved in an international Delphi study

*Palliative Medicine*  
2024, Vol. 38(4) 457–470  
© The Author(s) 2024



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/02692163241234579  
journals.sagepub.com/home/pmj



Mayumi Nishimura<sup>1</sup> , Karen Harrison Dening<sup>2</sup>,  
Elizabeth L Sampson<sup>3,4</sup>, Edison Iglesias de Oliveira Vidal<sup>5</sup> ,  
Miharu Nakanishi<sup>6</sup>, Nathan Davies<sup>7</sup> , Wilson Abreu<sup>8</sup>,  
Sharon Kaasalainen<sup>9</sup>, Yvonne Eisenmann<sup>10</sup>, Laura Dempsey<sup>11</sup>,  
Kirsten J Moore<sup>12,13</sup>, Sascha R Bolt<sup>14</sup>, Judith MM Meijers<sup>15,16,17</sup>,  
Natashe Lemos Dekker<sup>18</sup>, Mitsunori Miyashita<sup>19</sup> , Takeo Nakayama<sup>1</sup>  
and Jenny T van der Steen<sup>20,21,22</sup>

## Abstract

**Background:** Advance care planning in dementia includes supporting the person and their family to consider important goals of care. International research reports the importance of psycho-social-spiritual aspects towards end of life.

**Aim:** To develop a multidimensional international palliative care goals model in dementia for use in practice.

**Design:** International Delphi study integrating consensus and evidence from a meta-qualitative study. The Delphi panel rated statements about the model on a 5-point agreement scale. The criteria for consensus were pre-specified.

**Setting/participants:** Seventeen researchers from eight countries developed an initial model, and 169 candidate panellists were invited to the international online Delphi study.

<sup>1</sup>Department of Health Informatics, School of Public Health, Graduate School of Medicine, Kyoto University, Kyoto, Kyoto Prefecture, Japan

<sup>2</sup>Dementia UK, London, UK

<sup>3</sup>Royal London Hospital and Centre for Psychiatry and Mental health, East London NHS Foundation Trust, London, UK

<sup>4</sup>Wolfson Institute of Population Health, Queen Mary University London, London, UK

<sup>5</sup>Botucatu Medical School, Sao Paulo State University (UNESP), Botucatu, State of São Paulo, Brazil

<sup>6</sup>Department of Psychiatric Nursing, Tohoku University Graduate School of Medicine, Sendai, Miyagi, Japan

<sup>7</sup>Research Department of Primary Care and Population Health, University College London, London, UK

<sup>8</sup>Center for Health Technology and Services Research, University of Porto (ESEP/CINTESIS), Porto, Portugal

<sup>9</sup>School of Nursing, McMaster University, Hamilton, ON, Canada

<sup>10</sup>Department of Palliative Medicine, University Hospital of Cologne, Cologne, North Rhine-Westphalia, Germany

<sup>11</sup>Department of Nursing and Healthcare, Technological University of the Shannon: Midlands Midwest, Athlone, Ireland

<sup>12</sup>National Ageing Research Institute, Parkville, VIC, Australia

<sup>13</sup>Medicine-Royal Melbourne Hospital, The University of Melbourne, Melbourne, VIC, Australia

<sup>14</sup>Tilburg School of Social and Behavioral Sciences, Tilburg University, Tilburg, North Brabant, The Netherlands

<sup>15</sup>Department of Health Services Research, CAPHRI School for Public Health and Primary Care, Maastricht University, Maastricht, Limburg, The Netherlands

<sup>16</sup>Living Lab in Ageing and Long-Term Care, Maastricht, The Netherlands

<sup>17</sup>Zuyderland Care, Zuyderland Medical Center, Sittard-Geleen, The Netherlands

<sup>18</sup>Institute of Cultural Anthropology and Development Sociology, Leiden University, Leiden, South Holland, The Netherlands

<sup>19</sup>Department of Palliative Nursing, Health Sciences, Tohoku University Graduate School of Medicine, Sendai, Miyagi, Japan

<sup>20</sup>Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, South Holland, The Netherlands

<sup>21</sup>Department of Primary and Community Care, Radboud University, Nijmegen, The Netherlands

<sup>22</sup>Cicely Saunders Institute, King's College London, UK

## Corresponding author:

Jenny T van der Steen, Department of Public Health and Primary Care, Leiden University Medical Center, Albinusdreef 2, 2333 ZA Leiden, The Netherlands.

Email: jtvandersteen@lumc.nl

**Results:** Panellists (107; response 63.3%) resided in 33 countries. The model comprised four main care goals: (1) Comfort ensured; (2) Control over function maintained; (3) Identity protected and personhood respected and (4) Coping with grief and loss—person and caregiver supported. The model reflects how needs and care goals change over time with the progression of dementia, concluding with bereavement support. The first version of the model achieved a consensus after which it was slightly refined based on feedback. We did not achieve a consensus on adding a goal of life prolongation, and on use of the model by people with dementia and family themselves.

**Conclusion:** A new palliative care goals model for people with dementia and their families includes relationship aspects for use by professionals and achieved a consensus among a panel with diverse cultural background. The position of life prolongation in relation to palliative care goals needs further research.

### Keywords

Dementia, cognitive disorders, Delphi technique, model, biopsychosocial, advance care planning, interprofessional education, palliative care

#### What is already known about this topic?

- A white paper from the European Association for Palliative Care (EAPC) presented a care goals model for people with dementia covering changes of the goals of life prolongation; maintenance of functioning and maximisation of comfort towards death.
- Advance care planning discussions have gained momentum also for dementia, and recent Western and Asian studies indicated the relevance of psycho-social and spiritual care goals in addition to physical care goals.

#### What this study adds?

- A newly developed multidimensional palliative care goals model adds psycho-social-spiritual goals such as: identity protected and personhood respected; coping with grief and loss – person and caregiver supported.
- While the EAPC care goals model did not achieve a full consensus in the original Delphi study, the new multidimensional model did.
- The study highlights remaining controversy around life prolongation as a goal of care for persons with dementia.

#### Implications of research for clinical practice, theory or policy

- The model may broaden the perspective of healthcare providers to consider goals of care that may change as dementia progresses, focusing on care goals that are important in maintain good relationships and connectedness.
- The model aims to promote communication on how important each goal is in the context of advance care planning and how the goals can be achieved for individuals.
- Further research is needed into how to apply the model in practice and how the goals relate to a care goal of life prolongation.

## Introduction

Dementia is a progressive condition with people becoming increasingly dependent upon long-term care, while behaviour often changes and decision-making capacity decreases.<sup>1,2</sup> This can affect the person's relationships with their family and the community,<sup>3–5</sup> and a palliative care approach including early initiation of advance care planning and continuous communication about care goals is recommended.<sup>2,3,6,7</sup> Living with a progressive disease, often for years, implies that prioritised goals of care likely shift over time.<sup>8</sup>

A dementia-specific care goals model was developed in a Delphi study by the European Association for Palliative Care (EAPC) reported in the EAPC white paper on palliative dementia care. It presented goals of 'prolongation of life' decreasing in priority towards the end of life, the goal of 'maintenance of function' remaining relevant and

gradually prioritising 'maximisation of comfort'<sup>9,10</sup> (Figure 1(a)). However, the model achieved only a moderate consensus, with criticism, for example, that families were not included in the model (unpublished finding, personal communication).

Since the EAPC care goals model was published in 2014, caring for people with dementia and models of care have become more of a global topic.<sup>11–13</sup> Whereas the EAPC care goals model focuses on medical goals of care, studies from diverse countries have emphasised the importance of psycho-social-spiritual needs.<sup>14–16</sup> Our eight-country meta-qualitative study showed that a good end of life with dementia in Western, Southern and Asian countries included multidimensional concepts such as: Care for caregivers; Identity being preserved; and Being connected.<sup>17</sup> Other studies from Western and Asian countries also support the importance of relationships such as providing compassionate

care, and respecting personhood and dignity for people with dementia.<sup>18–21</sup> Therefore, expanding of current care goals models for use in care planning is timely and relevant. In addition, international models can support communication about values and concepts that risk being overlooked in individual countries.

This study aimed to develop a multidimensional international palliative care goals model specific to dementia based on evidence and consensus among a Delphi panel of experts. A secondary aim was to explore if support for the model varied with panellists' characteristics.

## Methods

We performed a Delphi study to develop a multidimensional palliative care goals model, aimed at achieving consensus among experts on the model and its potential use. Delphi studies are based on evidence and consensus in developing of content, multiple rounds of anonymous evaluation, with feedback to the panellists after each successive round.<sup>22–24</sup> An online Delphi study allows input from experts from diverse geographical locations in a transparent manner.<sup>22</sup> We used five rounds with input alternating between an international project team of researchers,<sup>17</sup> and a Delphi expert panel (Table 1). This Delphi study was embedded in a larger Delphi study on advance care planning in dementia with surveys conducted in 2021–2022.<sup>25</sup>

### *Delphi expert panel members*

The inclusion criteria were: (1) expertise in advance care planning, dementia care or advance care planning in dementia more specifically, through practice, policy, research or clinical experience and (2) ability to complete online surveys in English.

### *Sampling and recruitment*

Potential Delphi experts were identified via research and clinical networks, and from relevant publications in PubMed. We aimed for diversity in terms of expertise, continent of residence and occupation, with the aim of recruiting approximately 100 participants.

*Round 1: Developing a multidimensional palliative care goals model and preparation of evaluation items.* The project team of 17 researchers from eight countries, who had conducted the meta-qualitative study on a good end of life with dementia<sup>17</sup> developed an initial model (process of development in Figure 2).

The initial model was based on a previous eight-country meta-qualitative study in which the project team synthesised qualitative interview and observational data about experiences at the end of life of 121 people with

dementia and 292 families.<sup>17</sup> The data had been collected during 2009–2020 in the UK, Ireland, Portugal, Germany, the Netherlands, Canada, Brazil and Japan.<sup>14,16,26–36</sup> The synthesis process identified nine important components towards end of life<sup>17</sup> shown in the left column of Figure 2.

Based on the components, the project team discussed possible palliative goals of care that meet five criteria that were formulated during our discussions to consider the characteristics of appropriate care goals: (1) relevant to persons with dementia and family caregivers in different cultures—whether more individual or relationship centred—and in particular towards the end of life; (2) preferences tend to differ between individuals – therefore, cannot be taken for granted and needs to be discussed for each individual; (3) require appropriate effort by professionals and family caregivers to set feasible and achievable goals; (4) require variable effort over time indicating the notion of dynamic goals; (5) can involve trade-offs with other, competing goals of care which necessitates prioritisation.

Additionally, we reviewed panellists' comments regarding the Delphi study for the EAPC white paper in dementia (Figure 1(a)). There were comments on the lack of inclusion of family members' needs, preferences differing between individuals not being visualised and different importance attached to the three goals and change over time, in particular the goal of life prolongation. From these two sources (the meta-qualitative study and the earlier Delphi study), the team considered several drafts to present the care goals and also developed accompanying explanatory text.

*Round 2: Survey Delphi panel and interim analyses.* In September 2021, the potential panellists received an email inviting them to join the panel. An online pilot test was conducted with Leiden University Medical Center researchers on advance care planning in dementia. The online survey was developed with Castor Electronic Data Capture (EDC) System for Clinical Research Trials (Amsterdam, New York).

The evaluation items covered the proposed palliative care goals model, explanatory text, alternative formats for listing the goals of care and a comparison with the EAPC white paper care goals model (Figure 1(a)). We used statements that had been used to evaluate the EAPC white paper care goals model,<sup>10</sup> including those with reverse phrasing, such as 'the interpretation of the model is problematic.' The Delphi panellists rated levels of agreement on 1–5 agreement scales: strongly disagree (1); moderately disagree (2); neither agree nor disagree (3); moderately agree (4); and strongly agree (5). Based on median, inter-quartile range (IQR) and percent agreement, the consensus criteria (footnote to Table 1) were defined a priori with the same conservative criteria as previously used with the model in Figure 1(a).<sup>10</sup>

**Table 1.** Methods to develop and refine a palliative care goals model specific to dementia.

Round	Aim (month and year)	Design and methods; input	Contributors (numbers); expertise (numbers)	Main results and feeding into the next round
1	Developing a multidimensional palliative care goals model and preparation of evaluation items (June 2021)	Qualitative; the themes from the meta-qualitative synthesis of 14 literatures about experience of end of life. <sup>17</sup> The EAPC white paper care goals model and its open comments (Figure 2; Supplemental 2)	Project team (17) <sup>17</sup> ; Nurse (6) Physician (3) Epidemiologist (3) Psychologist (2) Sociologist (1) Anthropologist (1) Occupational therapist (1)	Project team proposed four care goals and drafted a figure as multidimensional palliative care goals model showing changing effort of the palliative care goals as the disease progresses (Figure 1(b)). The team also generated explanatory text about the model Table 3
2	Survey Delphi panel about multidimensional palliative care goals model and its explanatory texts (Figure 1) (September <sup>a</sup> –October 2021)	Quantitative and qualitative; 5-point agreement scale and open-ended items soliciting for comments; consensus determined by conservative criteria <sup>b</sup>	International panel (97/107 experts from 33 countries); experts in dementia care or advance care planning. Characteristics in Table 2	
3	Interim analysis and revising model (November, 2021)	Qualitative; discussion of consensus and comments in round 2	Project team (17); Same as round 1	Project team made minor revisions to the explanatory text, generating items to examine about life prolongation and use in practices Table 4
4	Survey Delphi panel about the model use (December 2021 <sup>c</sup> –January 2022) (1) relation of life-prolongation and the model (2) appropriate use of the model	Quantitative and qualitative; 5-point agreement scale and open-ended items soliciting for comments	International panel (86/107 experts from 33 countries) Same as round 2	
5	Interim analysis and finalising (January–February 2022)	Qualitative; discussion of agreement and comments in round 4  Synthesis; evaluation of results of all previous rounds, and open-ended comments in round 4	Core project team (6); Physician (2) Nurse (1) Epidemiologist (2) Occupational therapist (1) Project team (17); Same as round 1	Project team accepted diverse views about life prolongation and issues on use in practice.  Add minimal revision of the name of axis and finalised the model as the new multidimensional palliative care goals model for people with dementia and families (Figure 1(b))

<sup>a</sup>The survey was open for 23 days.

<sup>b</sup>Criteria for consensus (defined as high or very high (dis)agreement): very high agreement, a median of 5 and an IQR of 0 and  $\geq 80\%$  scoring a 4 or 5; high agreement, a median of 5 and an IQR  $\leq 1$  and  $\geq 80\%$  scoring a 4 or 5; moderate agreement, a median of 4–5 and an IQR  $\leq 2$  and  $\geq 60\%$  scoring a 4 or 5; low agreement, a median of 4–5 and an IQR  $\leq 2$  or  $\geq 60\%$  scoring a 4 or 5; no agreement, a median 4–5 otherwise or a median  $> 2$  and  $< 4$ . For consensus on disagreement, reverse median with the same IQR requirements and disagreement percentages (van der Steen et al., 2014).<sup>10</sup>


<sup>c</sup>The survey was open for 23–25 days.

**Round 3: Interim analysis and revising the model.** The results from Round 2 were tabulated and discussed in an interim analysis to inform revisions. A priori criteria for consensus including ‘high’ or ‘very high’ (dis)agreement are shown in Tables 3 and 4.<sup>25</sup> JTS and MaN summarised the comments from panellists using Microsoft Excel and proposed amendments to the project team. Unresolved issues were included in the next round.

**Round 4: Survey Delphi panel about the model use.** All panellists received the results of Round 2, and we

presented a revised model and additional questions based on the work undertaken in Round 3. The questions solicited after the model’s usefulness in the panellists’ own countries, and a goal of life prolongation related to the condition of the person with dementia.

**Round 5: Final analysis and conclusion of the study.** As in Round 3, we analysed the panellists’ evaluations and comments. A core project team (KHD, LS, EIOV, MaN, TN and JTS) discussed whether a further round would be helpful.




**Figure 1.** Existing (a) and newly developed final (b) models of dementia care goals. (a) A previous model - dementia progression and suggested prioritising of care goals in the European Association for Palliative Care dementia white paper. The goals of maintenance of function and maximisation of comfort are compatible with palliative care which aims to improve quality of life (van der Steen et al.<sup>10</sup> Copyright by the Authors. Reprinted by permission of SAGE Publications, Ltd.) (b) A model of palliative care goals towards the end of life for people living with dementia and family to support prioritising and detailing of effort to achieve individual care goals that contribute the most to quality of life until the end of life.

**Table 2.** Delphi expert panellists’ characteristics.

Characteristic	% or mean (SD)	n
Continent of residence (33 countries), %		
Europe (17 countries)	58.9	63
Americas (Northern, Southern, Middle; 4 countries)	18.7	20
Asia (6 countries)	10.3	11
Oceania/Australasia (2 countries)	5.6	6
Middle East (2 countries)	2.8	3
Africa (2 countries)	1.9	2
Alternating between continents	1.9	2
Gender, % <sup>a</sup>		
Woman	69.2	74
Man	30.8	33
Age, mean (SD)	52.0 (12.1)	104
Professional experience, mean number of years (SD)	24.4 (11.8)	103
Profession, %		
Medical (physician, physician assistant and nurse practitioner)	49.5	53
Non-medical (more possible), %	50.5	54
Nurse (any level)	19.6	21
Psychologist	11.2	12
Ethicist	8.4	9
Policy/administration	7.5	8
Social worker	4.7	5
Epidemiologist	3.7	4
Spiritual counsellor	2.8	3
Other, for example, sociologist and lawyer	11.2	12
Specific expertise, %		
Advance care planning in dementia specifically	57.7	60
Advance care planning or dementia care but not this combined	42.3	44
Personally experienced a family member or friend, %		
Having advanced dementia at the end of their life, yes	70.6	72
No such experience	29.4	30

n = 107.

<sup>a</sup>Other response options, not chosen, were ‘other’ and ‘prefer not to say.’



**Figure 2.** A process of development of an international multidimensional model.

a. The goal was relevant but control is less feasible (criterion 3 in the text) near the end of life. b. We rephrased for consistency, with the verb last. c. 'Life prolongation' was excluded because there was no consensus how to include it and to keep a focus on palliative goals exclusively. d. 'Satisfaction with life and spiritual wellbeing' was excluded because it is an overall outcome rather than a specific care goal that can be ensured.

### Secondary analysis

To examine if panellists with specific characteristics were more likely to agree with the model and other items, we compared the following subgroups of panellists (a) European versus non-European residence; (b) medical professionals as responsible for medical treatment and endorsing care plans versus other professionals; (c) expertise in advance care planning in dementia specifically versus no such

expertise specifically; (d) those with personal experience with advanced dementia care versus those without such experience personally.

### Ethical considerations

Model development was part of the meta-qualitative study approved by the Graduate School and Faculty of

**Table 3.** Delphi experts' evaluation of statements about the palliative care model for people with dementia.

Statement (n of expert respondents)	Median (IQR)	Percentage (dis) agreed <sup>a</sup>	Consensus; <sup>b</sup> level of agreement
<b>Overall</b>			
Having read the explanatory text <sup>c</sup> , I feel that the figure <sup>c</sup> represents appropriate palliative care goals and changes towards the end of life (n = 87, round 2)	5 (1)	80.5% agreed 10.3% disagreed	Yes; On high agreement
<b>Visualisation and interpretation</b>			
I feel that the palliative care goals should be listed only – with no visualisation of how they may change over time (n = 88, round 2)	2 (1)	10.2% agreed 78.4% disagreed	No; Moderate disagreement
I feel that comfort ensured should be the only palliative care goal with no trade off (n = 84, round 2)	2 (2)	20.2% agreed 65.5% disagreed	No; Moderate disagreement
I feel that the interpretation of the Figure above is problematic (n = 87, round 2)	2 (2)	24.1% agreed 58.6% disagreed	No; Low disagreement
<b>Compared with previous model (Figure 1(a))</b>			
This model replaces the previous model (n = 81, round 2)	4 (3)	60.5% agreed 27.2% disagreed	No; No agreement
This model complements the previous model (n = 78, round 2)	3 (2)	42.3% agreed 35.9% disagreed	No; No agreement
<b>Usefulness in own country</b>			
This model will help professionals with different background in my country to develop a shared understanding about palliative care for people with dementia (n = 83, round 4)	4 (1)	89.2% agreed 3.6% disagreed	No; Moderate agreement
This model will stimulate professionals in my country to consider goals aimed at the person's quality of life (n = 82, round 4).	4 (1)	90.2% agreed 2.4% disagreed	No; Moderate agreement
<b>Open ended comment from panellists on comparison two care goal models (n = 52)<sup>d</sup></b>			
Positive opinions about the new model			22
Valuing both models and could be used as needed			9
Preferring EAPC care goals model			8
Objections to not having included life-prolongation as a goal of care			7
Recommendation adding dementia stages in the new model			3
Recommendation adding other goals of care such as inclusion, having meaningful time, finding joy and activities especially in the early stages			3
Emphasising families' value on maintaining function			3
Objections to develop a visual model because it is considered to oversimplify palliative dementia care			3

IQR: inter-quartile range.

<sup>a</sup>Percent (dis)agreement refers to the percent combined strongly and moderately (dis)agree on the 1–5 agreement scale (percent answering with 4 or 5 for agreement; percent answering 1 or 2 for disagreement). <sup>b</sup>See footnotes to Tables 1 for criteria on consensus. <sup>c</sup>See the final explanatory text in Box 1 (minimally changed after round 3) and the previous iterations in Supplemental 2. <sup>d</sup>Some single comments included multiple opinions; therefore, the number of comments does not equal the total number of opinions.

Medicine Kyoto University (R1924-1), 25 April 2019. The advance care planning in dementia Delphi study protocol included evaluation of the model and was declared exempt from the Medical Research Involving Human Subjects Act by the Medical Ethical Committee Leiden Den Haag Delft, the Netherlands (N21.105). The protocol was registered at the WHO International Clinical Trials Registry Platform; NL9720, 7 September 2021. Confidentiality of individual responses was ensured by processing of coded data.

## Results

### Round 1

The initial model formulated by the project team consisted of four care goals, ranging from the individual level to the level of relationships and connectedness: Comfort ensured; Control over function; Identity protected personhood respected; Coping with grief and loss – person and family supported. To illustrate the changing emphasis on these four goals as dementia progresses, we created a

**Box 1.** Final version of the explanatory text that complements the care goals model.<sup>a</sup>

#### Purpose and use of the model of palliative care goals specific to dementia

The model aims to facilitate advance care planning discussions when, rather than length of life, quality of life is the main focus. The model may be applied from the early stages of dementia right up to the end of life. In order for the suggested goals to be achieved via the planned actions, the goals can be repeatedly selected (1), prioritised (2) and specified or refined for the individual (3).

#### Justification of four suggested palliative care goals

The four types of palliative care goals, indicated with four colours:

- are often important to persons with dementia and family in various cultures, whether more individual or relationship centred and in particular towards the end of life;
- represent care goals at macro-level for which preferences tend to differ between individuals, therefore cannot be taken for granted and the meaning of the care goals in the model need to be discussed for each individual;
- are feasible and realistic goals and be achieved with the appropriate effort from professionals and family;
- are goals that often require variable effort over time indicating the notion of being dynamic;
- can involve trade-offs with other, competing care goals which necessitates prioritisation.<sup>10,17,37,52</sup>

#### Coping with grief and loss – person and family supported and Bereavement support

Family emotional and practical support needs may change over time in various ways, and so may support needs in coping with grief and loss of the family and the person with dementia. The variability is indicated with a horizontal yet dashed line. Support needs may emerge that lead to, for example nursing home or hospital admission, or with difficult decisions in moderate dementia, or in the severe stage towards the end of life.<sup>10,38–45</sup>

#### Identity protected and personhood respected

Maintaining identity requires increasing effort and protection from those around the person with dementia. At times this may be important to both persons with dementia and family.<sup>17,37</sup> People change during their life, and persons with dementia and their relationships also change as a result of, for example, coping with changes in the condition.<sup>4</sup> Family may grieve the perceived loss of previous identity.<sup>4</sup> Further, aspects of identity such as personality may change. All of this should be accommodated in the care, respecting present identity and personhood and being treated as a fellow human being as a human right.<sup>46,47</sup> Some feel that personhood should be respected after life as well.

#### Control over function maintained

Functions, such as walking without help and the ability to communicate, decline due to dementia. However, such functions contribute to remaining connected and to quality of life, and many value control in terms of autonomy or maintaining activities for as long as possible.<sup>2,37,40</sup> At some point, however, efforts to maintain function may no longer be effective or present a burden as they cause distress or pain to the person and can become inappropriate.<sup>10,38</sup>

#### Comfort ensured

Ensuring comfort with little trade-off as regards other care goals may become more of a priority over time.<sup>10</sup>

Examples of trade-offs between the four goals

There are trade-offs between the care goals of functioning versus comfort as a means to achieve quality of life, and there are dilemmas in decisions around life-prolonging and potentially burdensome medical treatment.<sup>10</sup> However, there is little evidence on dilemmas created by competing palliative goals of care.

Some examples are:

- staying at home as this usually supports preserving or protecting **identity** the most, but this should be weighed against **family caregiver** burden;
- when people prioritise **identity** over **comfort** for choice of clothing that expresses an identity that is strongly adhered to maintaining identity, but wearing these clothes may be less comfortable.

Examples of competing comfort and functioning goals are:

- careful hand feeding for pleasure, irrespective of amount or nutritional value may become more important than feeding to preserve muscles or maintain **function**;
- maintaining **function** might threaten **comfort**, for example when encouraging mobility causes pain, while accepting a risk of falls might be appropriate;
- using opioids to decrease symptoms such as shortness of breath and improve **comfort** may decrease the **function** of communicating with loved ones.

<sup>a</sup>This text was shown to the panellists in round 2 and 4, except for a minor revision after round 3. In the section of 'Identity protected and personhood respected,' the project team added 'Some feel that personhood should be respected after life as well' based on comments from a panellist.



**Table 4.** Delphi experts' rating of statements about life prolongation and applicability of the care goals model.

Statement ( <i>n</i> of expert respondents, round 4)	Median (IQR)	Percentage (dis)agreed <sup>a</sup>	Consensus; <sup>a</sup> level of agreement
<b>How the model may relate to life prolongation</b>			
<b>Life prolongation in own country</b>			
In my country, too often, life prolongation is an inappropriate (implicit or explicit) goal of care and treatment ( <i>n</i> = 79)	4 (2)	60.8% agreed 15.2% disagreed	No; Moderate agreement
<b>Position of life prolongation</b>			
Applying this model requires a two-step approach: first, determine if life prolongation is (also) a care goal; next, discuss palliative care goals ( <i>n</i> = 81)	4 (2)	66.7% agreed 20.9% disagreed	No; Moderate agreement
This model should include 'life prolongation' as a goal competing with the palliative care goals (with palliative care not aiming to prolong or shorten life) ( <i>n</i> = 82)	3 (2)	35.4% agreed, 48.8% disagreed	No; No agreement
<b>Comorbidity</b>			
This model (without life prolongation added as a competing goal) is particularly helpful to consider care goals when dementia is the only relevant diagnosis (no comorbid conditions) ( <i>n</i> = 83)	4 (2)	60.2% agreed 16.9% disagreed	No; Moderate agreement
Life prolongation should be addressed as a competing goal (either as a first step or added to the model) depending on frequent relevant comorbid conditions of the person. . . .			
when the person has dementia and recurrent pneumonia due to dysphagia ( <i>n</i> = 75)	4 (3)	52.0% agreed 26.7% disagreed	No; No agreement
when the person has dementia and terminal cancer ( <i>n</i> = 77)	3 (3)	49.4% agreed 37.7% disagreed	No; No agreement
when the person has dementia and serious consequences from a stroke ( <i>n</i> = 76)	4 (3)	51.3% agreed 32.9% disagreed	No; No agreement
when the person has dementia and heart failure ( <i>n</i> = 76)	4 (2)	55.2% agreed 22.4% disagreed	No; Low agreement
Statement ( <i>n</i> = response per item)	Median (IQR)	Percentage (dis)agreed <sup>a</sup>	Consensus; <sup>a</sup> Level of agreement
<b>Setting; this model can be used in. . .</b>			
a. Own home with no healthcare professional involved ( <i>n</i> = 77)	4 (2)	53.3% agreed 31.2% disagreed	No; Low agreement
b. Home care ( <i>n</i> = 81)	4 (1)	85.2% agreed 7.4% disagreed	No; Moderate agreement
c. Group housing mainly offering social care and support ( <i>n</i> = 80)	4 (1)	81.3% agreed 10.0% disagreed	No; Moderate agreement
d. Nursing home offering medical and nursing care ( <i>n</i> = 83)	5 (1)	90.4% agreed 3.6% disagreed	Yes; On high agreement
e. Hospital ( <i>n</i> = 82)	5 (1)	90.3% agreed 6.1% disagreed	Yes; On high agreement
f. Hospice ( <i>n</i> = 81)	5 (1)	87.7% agreed 6.2% disagreed	Yes; On high agreement
<b>Professions; this model can be used by. . .</b>			
a. Physicians ( <i>n</i> = 81)	5 (0)	93.8% agreed 4.9% disagreed	Yes; on very high agreement
b. Trained non-physician facilitators including nurses, nurse assistants and social care professionals ( <i>n</i> = 81)	5 (1)	91.3% agreed 3.7% disagreed	Yes; On high agreement
c. Persons with dementia and their family in non-healthcare settings ( <i>n</i> = 78)	4 (2)	64.1% agreed 15.4% disagreed	No; Moderate agreement

<sup>a</sup>See explanatory text in Tables 1 and 3 for criteria on agreement and consensus.

model (Supplemental 2) and explanatory text (Figure 1). Several models were considered, including one that only listed the four care goals, one including life prolongation as a goal and one that emphasised comfort at the end as in the EAPC model. Delphi items were developed to rate the agreement level to the models and to invite open comments.

### Round 2

On September 2021, experts of dementia care and advance care planning were invited via email. Of 169 experts reached, 107 (63.3%) from 33 countries participated (response from Europe, 67.0%; from elsewhere, 58.7%). Half of respondents were medical professionals (53/107). Over half (60/104 – some missing values) had expertise in advance care planning in dementia specifically, and 72 out of 102 had personal experience (Table 2). Table 3 shows that the first iteration of the model achieved a consensus on representing appropriate palliative care goals and changes towards the end of life (80.5% agreed, median agreement rating 5, IQR 1). Two items that showed moderate disagreement were: (1) Not offering a visualisation of changes over time (10.2% agreed, 78.4% disagreed; median 2, IQR 1), and (2) Comfort should be the only palliative care goal with no trade off (20.2% agreed, 65.5% disagreed; median 2, IQR 2).

A total of 53 comments were made regarding comparing the new models with the EAPC care goals model of Figure 1(a). Of these (Table 3), 22 commented in a generally positive way, preferring the new model, while 8 preferred Figure 1(a), and 9 suggested both models are valuable and could be used as needed. Further, nine comments objected to omitting life prolongation as a goal of care (from Portugal, two; Netherlands, two; USA, Czech Republic, UK, Switzerland and Spain each one). In addition, three participants recommended adding dementia stages as in Figure 1(a) and two emphasised spiritual concerns after death.

### Round 3

Reviewing the panellists' feedback, the project team made minor amendments. We added the label 'After death' on the timeline and to the explanatory text, 'Some feel that personhood should be respected after life as well'.

Additionally, having already achieved a consensus on the model, the project team decided to move on to projected implementation and consult the panel's opinion on the applicability of this model, who and in which place of care it would be appropriate to use it. Other statements referred to panellists' degrees of agreement on usefulness of the model and the position of life prolongation in

their own country, and to applicability of the model depending on comorbidity, care settings and appropriate users of the model.

### Round 4

Table 4 shows that there was no consensus on the inclusion of life prolongation in this model, irrespective of comorbidities. Diverse views on life prolongation surfaced from 28 panellists' comments. Six panellists commented that life prolongation should always be included in the model for its usefulness in practice. In contrast, 12 others would not include life prolongation because 'it should not compete with other care goals' (10 panellists) or 'it is just beyond the scope of this palliative care goals model' (two panellists). Ten other panellists did not clearly express an opinion on whether to include it, but commented it should be considered: when the person with dementia wishes (four panellists); at an early stage of dementia (three panellists); when the person has heart failure because it is not part of advanced dementia (two panellists); when medical intervention is helpful but may threaten other goals (one panellist).

As for applying the model (Table 4), there was a consensus (high agreement) that the model could be implemented in nursing homes, hospitals and hospices. As well as a consensus with very high agreement on its use by physicians, there was a consensus with high agreement that it could be used by other trained healthcare professionals and social care staff.

There was no consensus on using the model in home care with no healthcare professional involved, which was consistent with the lack of consensus on the use of the model by persons with dementia and their families in non-healthcare settings. Suggestions included that a lay version with plain words would be needed if the person with dementia and their family were to use it on their own. Other comments referred to a need to clarify the labelling of the axes.

### Round 5

The core project team discussed various issues and whether any new issues had appeared in Round 4 from panellists' comments that would necessitate another round.

In summary, the main points raised and addressed were the following: (1) the X-axis label of 'emphasis and effort' was moved to the Y-axis so that the X-axis represented time only; (2) The lack of consensus in Round 4 among the panel on the position of life prolongation was also reflected within the project team; and (3) team discussions did not result in new insights which could feed into a next round. We agreed to accept the diversity in

opinions on the position of life prolongation and decided to not proceed with a further survey round.

### *Subgroup consensus*

The subgroup analyses (Supplemental 1) indicated that the model as presented the first time did not achieve a consensus in the subgroups of medical professionals, those with expertise in advance care planning in dementia specifically, and those with no personal experience. We found other subgroup differences, across the four subgroups, in consensus on the two statements about usefulness in the country of the panellists. The level of consensus about the use of the model in a home care setting differed across all four subgroups. Panellists from Europe did support the use of the model at home, as did panellists from non-medical professions, and no experience in advance care planning in dementia specifically or personally. In contrast, as in the total group, none of the items on life prolongation as a goal achieved a consensus among any of the subgroups.

## **Discussion**

### *Main findings*

We have developed a new palliative care goals model for people with dementia and their family based on a meta-qualitative study and a Delphi study. The model demonstrates how people can choose over time from four potentially competing goals of care regarding individuals, family and relationships. These include: Comfort ensured; Control over function maintained; Identity protected and personhood respected; Coping with grief and loss-person and family supported which concludes with 'bereavement support.'

### *What this paper adds?*

An EAPC white paper from 2014 defined optimal palliative care in dementia and presented a care goals model comprising medical care goals: Maintenance of function, Prolonging life; and Maximisation of comfort. The new model adds psycho-social-spiritual goals at the individual and relational level, and it has been supported by the diverse panel of experts from across the globe. The expert panel also agreed with its use by medical and social care professionals and applicability in other institutional settings.

The exploration of subgroups indicated that those with no medical or no professional advance care planning experience in dementia specifically were slightly less critical regarding the model. The most subgroup differences concerned the use of the models in home care, which may point to cultural, social or national policy differences in

the quantity or quality of home care or how it is being organised and funded.<sup>48–51</sup>

A possible reason why no consensus was reached on including life prolongation to the multidimensional model may relate to the definitions of palliative care by World Health Organization and the international Association for Hospice and Palliative Care, which states that palliative care 'intends neither to hasten nor postpone death.'<sup>52,53</sup> Other possible reasons include that this goal is the most medically oriented, or else how participants define futile treatment in the context of dementia is affected by their personal or cultural perspective. However, issues about prolonging life remain important; for example, when considering whether to withhold the treatment of comorbid, intercurrent disease or whether to use (intravenous) antibiotics to treat pneumonia. Clearly, different people have different perspectives on the meaning of life prolongation and its relevance as a care goal in the use of this model.<sup>54,55</sup>

In practice, this international multidisciplinary model may broaden the perspective of healthcare providers to consider goals of care that extend beyond mere medical care and that may change over time as dementia progresses. The model may promote a focus on care goals that are important to maintain good relationships and connectedness, which add to positive care experiences for persons with dementia and their family.<sup>56,57</sup> Using the model may be combined with existing decision-making tools.<sup>58–65</sup> while explaining that the way to achieving the competing goals of care will vary depending on their personal values and preferences.

Of note, the model does not force any decisions to be made in individual cases, nor gives simple answers of care goals in different cultural and social background. To consider individual care goals, the process of communication is still indispensable by actively listening to individual needs with guidance to encourage addressing of specific topics represented in the model. As Davies et al. pointed out, some people do not wish to think about the future and prefer to think about everyday life. Further research may consider how the model may support resilience of the wider networks in the environment around the person with dementia.<sup>66</sup> Understanding diversity in meaning of life prolongation as a care goal is important for the use both of this model and the white paper model that also included it. The usefulness of life prolongation as a goal of care should also be understood from the perspective of persons with dementia and their family to best serve their interests.

### *Strengths and weaknesses*

We did not include persons with dementia and family caregivers in the Delphi panel, but most experts had personally experienced a lived one with dementia at the end of

life. The model may not fully capture diversity, also because experts who could not communicate in English were not included. We used rigorous predetermined criteria, that may have led to overestimation of the degree of disagreement between subgroups in our subgroup analysis. In addition, there was no authorisation of the model by an external body (called 'external validation'<sup>22</sup>). Future studies should examine whether the international consensual model is applicable in practice and can improve the quality of care, and living and dying with dementia for those involved in diverse locations across the globe.

## Conclusion

A new palliative care goals model for people with dementia and their family with psycho-social-spiritual goals achieved consensus amongst a diverse expert panel. Controversy around life prolongation as a goal of care remained and needs further research.

## Acknowledgements

We thank EAPC and the Delphi panellists for their contributions and support in conducting the Delphi study. Also, we sincerely thank Professor Tom Dening, MA, MD, PhD, FRCPsych, Dementia Research, School of Medicine, University of Nottingham, UK, for his help in language editing of this manuscript.

## Author contributions

MaN, TN and JTS contributed to acquisition. MaN and JTvds jointly drafted the article. The core project team members (MaN, KHD, LS, EIOV, MiN, TN and JTS) contributed substantially to interpreting the survey findings and finalising the model. All authors made a substantial contribution to the concept or design of the work, and to analysis or interpretation of the data. Also, all authors revised the article critically for important intellectual content, and approved the version to be published. All authors have participated sufficiently in the work to take public responsibility for appropriate portions of the content.

## Declaration of conflicting interests

The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: This study was performed with the same expert panel as with EAPC's ACP Delphi study, but this study was not designed by the EAPC ACP in dementia task force. This separate study was not conducted as part of the remit of the EAPC task force. There are no conflicts of interest.


## Funding


The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: MaN and TN are supported by the Graduate School of Kyoto University, Japan. JTvds is supported by a European Research Council (ERC) Consolidator Grant (Agreement ID 771483).

## ORCID iDs

Mayumi Nishimura  <https://orcid.org/0000-0001-8025-5211>

Edison Iglesias de Oliveira Vidal  <https://orcid.org/0000-0002-1573-4678>

Nathan Davies  <https://orcid.org/0000-0001-7757-5353>

Mitsunori Miyashita  <https://orcid.org/0000-0002-7637-0409>

Jenny T van der Steen  <https://orcid.org/0000-0002-9063-7501>

## Data availability

Part of the data that support the findings of this study are available on request from the corresponding author, JTvds, for the data provided by panellists who provided informed consent for reuse for other research. There was no consent to provide open access to the data.

## Supplemental material

Supplemental material for this article is available online.

## References

1. Alzheimer's Disease International. Dementia statistics. Alzheimer's Disease International, <https://www.alzint.org/about/dementia-facts-figures/dementia-statistics/> (2021, accessed 20 October 2021).
2. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet* 2020; 396: 413–446.
3. Senderovich H and Retnasothie S. A systematic review of the integration of palliative care in dementia management. *Palliat Support Care* 2020; 18: 495–506.
4. Conway ER, Watson B, Tatangelo G, et al. Is it all bleak? A systematic review of factors contributing to relationship change in dementia. *Int Psychogeriatrics* 2018; 30: 1619–1637.
5. Alzheimer's Disease International. *World Alzheimer Report 2019 Attitudes to dementia*. London: Alzheimer's Disease International, 2019.
6. Volicer L and Simard J. Palliative care and quality of life for people with dementia: medical and psychosocial interventions. *Int Psychogeriatrics* 2015; 27: 1623–1634.
7. Lee EE, Chang B, Huege S, et al. Complex Clinical Intersection: Palliative Care In Patients With Dementia. *Am J Geriatr Psychiatry* 2018; 26: 224–234.
8. Lynn J and Adamson DM. *Living well at the end of life*. Santa Monica, CA: RAND, 2003.
9. Gillick MR. Adapting advance medical planning for the nursing home. *J Palliat Med* 2004; 7: 357–361.
10. van der Steen JT, Radbruch L, Hertogh CM, et al. White paper defining optimal palliative care in older people with dementia: a Delphi study and recommendations from the European Association for Palliative Care. *Palliat Med* 2014; 28: 197–209.
11. World Health Organization. *Global action plan on the public health response to dementia 2017 - 2025*. Geneva, Switzerland: World Health Organization. <https://www.who.int/dementia>

- who.int/publications/i/item/global-action-plan-on-the-public-health-response-to-dementia-2017---2025 (2017, accessed 29 February 2024).
12. Prince M, Ali G-C, Guerchet M, et al. Recent global trends in the prevalence and incidence of dementia, and survival with dementia. *Alzheimers Res Ther* 2016; 8: 23.
  13. Ezzati M, Pearson-Stuttard J, Bennett JE, et al. Acting on non-communicable diseases in low- and middle-income tropical countries. *Nature* 2018; 559: 507–516.
  14. Nishimura M, Kohno A, van der Steen JT, et al. Conceptualization of a good end-of-life experience with dementia in Japan: a qualitative study. *Int Psychogeriatrics* 2020; 32: 255–265.
  15. Takahashi Z, Yamakawa M, Nakanishi M, et al. Defining a good death for people with dementia: a scoping review. *Jpn J Nurs Sci* 2021; 18: e12402.
  16. Davies N, Rait G, Maio L, et al. Family caregivers' conceptualisation of quality end-of-life care for people with dementia: a qualitative study. *Palliat Med* 2017; 31: 726–733.
  17. Nishimura M, Denning KH, Sampson EL, et al. Cross-cultural conceptualization of a good end of life with dementia: a qualitative study. *BMC Palliat Care* 2022; 21: 1–18.
  18. Wehrmann H, Michalowsky B, Lepper S, et al. Priorities and preferences of people living with dementia or cognitive impairment – a systematic review. *Patient Prefer Adherence* 2021; 15: 2793–2807.
  19. Kim SK and Park M. Effectiveness of person-centered care on people with dementia: a systematic review and meta-analysis. *Clin Interv Aging* 2017; 12: 381–397.
  20. Wang J, Wu B, Bowers BJ, et al. Person-centered dementia care in China: a bilingual literature review. *Gerontol Geriatr Med* 2019; 5: 2333721419844349.
  21. Eriksen S, Engedal K and Grov EK. Lifeworld perspectives of people with dementia: a meta-aggregation of qualitative studies. *BMC Geriatr* 2022; 22: 1–8.
  22. Jünger S, Payne SA, Brine J, et al. Guidance on Conducting and REporting DELphi Studies (CREDES) in palliative care: recommendations based on a methodological systematic review. *Palliat Med* 2017; 31: 684–706.
  23. Hasson F, Keeney S and Mckenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs* 2000; 32: 1008–1015.
  24. Fitch K. *The Rand/UCLA appropriateness method user's manual*. Santa Monica, CA: RAND, 2001.
  25. van der Steen JT, Nakanishi M, Van den Block L, et al. Consensus definition of advance care planning in dementia: a 33- country delphi study. *Alzheimer's Dement*. Epub ahead of print 20 November 2023. DOI: 10.1002/alz.13526.
  26. Moore KJ, Davis S, Gola A, et al. Experiences of end of life amongst family carers of people with advanced dementia: Longitudinal cohort study with mixed methods. *BMC Geriatr* 2017; 17: 1–14.
  27. Lemos Dekker N. Moral frames for lives worth living: Managing the end of life with dementia. *Death Stud* 2018; 42: 322–328.
  28. Schmidt H, Eisenmann Y, Golla H, et al. Needs of people with advanced dementia in their final phase of life: a multi-perspective qualitative study in nursing homes. *Palliat Med* 2018; 32: 657–667.
  29. McCleary L, Thompson GN, Venturato L, et al. Meaningful connections in dementia end of life care in long term care homes. *BMC Psychiatry* 2018; 18: 307.
  30. Thompson G, Shindruk C, Wickson-Griffiths A, et al. "Who would want to die like that?" Perspectives on dying alone in a long-term care setting. *Death Stud* 2019; 43: 509–520.
  31. Harrison Denning K, Jones L, Sampson EL, et al. Preferences for end-of-life care: a nominal group study of people with dementia and their family carers. *Palliat Med* 2012; 27: 409–17.
  32. Bamford C, Lee R, McLellan E, et al. What enables good end of life care for people with dementia? A multi-method qualitative study with key stakeholders. *BMC Geriatr* 2018; 18: 302.
  33. Dempsey L, Dowling M, Larkin P, et al. Providing care for a person with late-stage dementia at home: what are carers' experiences? *Dementia* 2018; 19: 352–374.
  34. Lillo-Crespo M, Riquelme J, Macrae R, et al. Experiences of advanced dementia care in seven European countries: implications for educating the workforce. *Glob Health Action* 2018; 11: 1478686.
  35. Bolt SR, van der Steen JT, Schols JMGA, et al. Nursing staff needs in providing palliative care for people with dementia at home or in long-term care facilities: a scoping review. *Int J Nurs Stud* 2019; 96: 143–152.
  36. Watarai H, Takahashi Z, Yamakawa M, et al. A qualitative study to explore good death for people with dementia. In: *The 24th annual meeting of the Japanese society of palliative medicine*. Yokohama, Japan, 21-22 June 2019. Osaka: JSPM
  37. Tochel C, Smith M, Baldwin H, et al. What outcomes are important to patients with mild cognitive impairment or Alzheimer's disease, their caregivers, and health-care professionals? A systematic review. *Alzheimers Dement* 2019; 11: 231–247.
  38. Robinson L, Gemski A, Abley C, et al. The transition to dementia—individual and family experiences of receiving a diagnosis: a review. *Int Psychogeriatr* 2011; 23(7): 1026–1043.
  39. Lee K, Puga F, Pickering CEZ, et al. Transitioning into the caregiver role following a diagnosis of Alzheimer's disease or related dementia: a scoping review. *Int J Nurs Stud* 2019; 96: 119–131.
  40. Davison TE, Camões-Costa V and Clark A. Adjusting to life in a residential aged care facility: Perspectives of people with dementia, family members and facility care staff. *J Clin Nurs* 2019; 28(21–22): 3901–3913.
  41. Afram B, Verbeek H, Bleijlevens MHC, et al. Needs of informal caregivers during transition from home towards institutional care in dementia: a systematic review of qualitative studies. *Int Psychogeriatr* 2015; 27(6): 891–902.
  42. Saragosa M, Jeffs L, Okrainec K, et al. Using meta-ethnography to understand the care transition experience of people with dementia and their caregivers. *Dementia* 2022; 21: 153–180.
  43. Prorok JC, Horgan S and Seitz DP. Health care experiences of people with dementia and their caregivers: a meta-ethnographic analysis of qualitative studies. *CMAJ* 2013; 185(14): E669–E680.
  44. Sellars M, Chung O, Nolte L, et al. Perspectives of people with dementia and carers on advance care planning and end-of-life care: a systematic review and thematic synthesis of qualitative studies. *Palliat Med* 2019; 33(3): 274–290.

45. Cresp SJ, Lee SF and Moss C. Substitute decision makers' experiences of making decisions at end of life for older persons with dementia: a systematic review and qualitative meta-synthesis. *Dementia* 2020; 19(5): 1532–1559.
46. Gove D, Sparr S, Dos Santos Bernardo AM, et al. Recommendations on end-of-life care for people with dementia. *J Nutr Health Aging* 2010; 14(2): 136–139.
47. Peisah C, Sampson EL, Rabheru K, et al. The human rights of older people with mental health conditions and psychosocial disability to a good death and dying well. *Am J Geriatr Psychiatry* 2021; 29: 1041–1046.
48. Arias-Casais N, Garralda E, Sánchez-Cárdenas MA, et al. Evaluating the integration of palliative care in national health systems: an indicator rating process with EAPC task force members to measure advanced palliative care development. *BMC Palliat Care* 2021; 20: 1–7.
49. Maetens A, Beernaert K, Deliens L, et al. Policy measures to support palliative care at home: a cross-country case comparison in three european countries. *J Pain Symptom Manage* 2017; 54: 523–529.e5.
50. Tolson D, Holmerova I, Macrae R, et al. Improving advanced dementia care: an interprofessional palliative learning framework. *J Am Med Dir Assoc* 2017; 18: 561–563.
51. Abu-Odah H, Molassiotis A and Liu JYW. Gathering policy-makers' perspectives as an essential step in planning and implementing palliative care services at a national level: an example from a resource-limited country. *BMC Palliat Care* 2022; 21: 1–13.
52. Radbruch L, De Lima L, Knaul F, et al. Redefining palliative care—a new consensus-based definition. *J Pain Symptom Manage* 2020; 60: 754–764.
53. World Health Organization. *National Cancer Control Programmes. Policies and managerial guidelines*. 2nd ed. Geneva: World Health Organization, 2002.
54. Malhotra C, Mohamad H, Østbye T, et al. Discordance between dementia caregivers' goal of care and preference for life-extending treatments. *Age Ageing* 2021; 50: 1382–1390.
55. Lord K, Livingston G and Cooper C. A systematic review of barriers and facilitators to and interventions for proxy decision-making by family carers of people with dementia. *Int Psychogeriatrics* 2015; 27: 1301–1312.
56. Monin JK, Valeika S and Gil H. A dementia positive, family-centered question geriatricians can ask caregivers and persons living with dementia. *J Am Geriatr Soc* 2023; 71: 2049–2051.
57. Sideman AB, Merrilees J, Dulaney S, et al. "Out of the clear blue sky she tells me she loves me": Connection experiences between caregivers and people with dementia. *J Am Geriatr Soc* 2023; 71: 2172–2183.
58. Tolson D, Fleming A, Hanson E, et al. achieving prudent dementia care (palliare): an international policy and practice imperative. *Int J Integr Care* 2016; 16: 1–11.
59. Holmerová AI, Waugh A, Macrae R, et al. *Dementia palliare best practice statement*. Paisley, Scotland: University of the West of Scotland, 2016.
60. Hanson E, Hellström A, Sandvide Å, et al. The extended palliative phase of dementia – an integrative literature review. *Dementia* 2019; 18: 108–134.
61. Harding AJE, Doherty J, Bavelaar L, et al. A family carer decision support intervention for people with advanced dementia residing in a nursing home: a study protocol for an international advance care planning intervention (mySupport study). *BMC Geriatr* 2022; 22: 1–10.
62. Brazil K, Walshe C, Doherty J, et al. MySupport Study Group. Implementation of an advance care planning intervention in nursing homes: an international multiple case study. *Gerontologist*. 2024 Feb 13:gnae007. doi: 10.1093/geront/gnae007. Epub ahead of print.
63. Davies N, de Souza T, Rait G, et al. Developing an applied model for making decisions towards the end of life about care for someone with dementia. *PLoS ONE* 2021; 16: e0252464.
64. Davies N, Aker N, Vickerstaff V, et al. A feasibility study of a decision aid to support family carers of people with severe dementia or those towards the end-of-life. *Palliat Med* 2022; 36: 1432–1439.
65. Anantapong K, Bruun A, Walford A, et al. Co-design development of a decision guide on eating and drinking for people with severe dementia during acute hospital admissions. *Heal Expect* 2023; 26: 613–629.
66. Peeters G, Kok A, de Bruin SR, et al. Supporting resilience of older adults with cognitive decline requires a multi-level system approach. *Gerontology* 2023; 69: 866–874.